

Optical counterparts of ROSAT X-ray sources in two selected fields at low vs. high Galactic latitudes

J. Greiner¹, G.A. Richter²

¹ Max-Planck-Institut für extraterrestrische Physik, 85740 Garching, Germany

² Sternwarte Sonneberg, 96515 Sonneberg, Germany

Received 14 October 2013 / Accepted 12 August 2014

ABSTRACT

Context. The optical identification of large number of X-ray sources such as those from the ROSAT All-Sky Survey is challenging with conventional spectroscopic follow-up observations.

Aims. We investigate two ROSAT All-Sky Survey fields of size $10^\circ \times 10^\circ$ each, one at galactic latitude $b = 83^\circ$ (26 Com), the other at $b = -5^\circ$ (γ Sge), in order to optically identify the majority of sources.

Methods. We used optical variability, among other more standard methods, as a means of identifying a large number of ROSAT All-Sky Survey sources. All objects fainter than about 12 mag and brighter than about 17 mag, in or near the error circle of the ROSAT positions, were tested for optical variability on hundreds of archival plates of the Sonneberg field patrol.

Results. The present paper contains probable optical identifications of altogether 256 of the 370 ROSAT sources analysed. In particular, we found 126 AGN (some of them may be misclassified CVs), 17 likely clusters of galaxies, 16 eruptive double stars (mostly CVs), 43 chromospherically active stars, 65 stars brighter than about 13 mag, 7 UV Cet stars, 3 semiregular resp. slow irregular variable stars of late spectral type, 2 DA white dwarfs, 1 Am star, 1 supernova remnant and 1 planetary nebula. As expected, nearly all AGN are found in the high-galactic latitude field, while the majority of CVs is located at low galactic latitudes. We identify in total 72 new variable objects.

Conclusions. X-ray emission is, expectedly, tightly correlated with optical variability, and thus our new method for optically identifying X-ray sources is demonstrated to be feasible. Given the large number of optical plates used, this method was most likely not more efficient than e.g. optical spectroscopy. However, it required no telescope time, only access to archival data.

Key words. X-rays: stars – stars: variables – techniques: photometric – surveys

1 Introduction

In the past, many attempts were made to investigate the stellar content in the Solar neighbourhood, as well as in the whole Galaxy, and from that deduce the structure of our Galaxy by studying the spatial distribution and the dynamics of different types of objects within the Galaxy. An early summary of such analyses based on optical data is given by Hoffmeister et al. (1985). Variable stars played an important role right from the beginning; e.g., Richter (1968) investigated the structure of our Galaxy by means of the statistics of the variable stars of the Sonneberg field patrol.

Since that time, many objects have been found in other spectral regions by new ground-based (e.g. 2MASS, Pan-STARRS, PTF, SDSS) or space-based (ROSAT, GALEX, WISE, Fermi) surveys, and their spatial distribution is of interest. In this connection the X-ray sources found by ROSAT are very important because of its large number of about 200 000 sources, and their all-sky distribution (note that pointed Chandra, XMM-Newton or Swift/XRT observations together cover only about 7–8% of the sky). Beyond the sheer number, these surveys at other wavelengths have the additional advantage of suffering from completely different selection biases, thus improving our understanding of the limits of optical surveys. Examples include not only source populations dominated by X-ray emission like single neu-

tron stars (Haberl 2005), but also classical optical populations like chromospherically active stars (Schmitt & Liefke 2004).

Before these surveys can be used for this kind of study, rather complete optical identifications are necessary. In general, standard practice of optical identification of X-ray sources is via optical spectroscopy. Crude identifications can also be made based on optical colours and on the ratio of optical-to-X-ray flux (Maccacaro et al. 1982, Stocke et al. 1991, Beuermann et al. 1999), since different object classes have different efficiencies in producing X-ray emission. For the selection of blazars, the correlation of X-ray and radio surveys have proven very efficient (e.g. Brinkmann et al. 1995). However, for the purpose of the present paper we have chosen a different approach, namely to use archival data available from the Sonneberg optical sky survey. The underlying idea was that the majority of X-ray-emitting objects is expected to be variable and that the chance coincidence of a variable inside the ROSAT error box is small.

The Sonneberg sky survey consists of two major parts: the sky patrol and the field patrol (Bräuer et al. 1999). The sky patrol records the entire northern sky since 1926 with 14 short-focus cameras in two colours down to a limiting magnitude of $\sim 14^m$. The field patrol monitored 80 selected $10^\circ \times 10^\circ$ fields between 1926 and 1995 in one colour down to a limiting magnitude of $\sim 18^m$ at the plate centre.

Here we present the results of the classification of selected ROSAT X-ray point sources and their optical counterparts based on archival photographic plates, complemented by the standard identification methods of a low- (γ Sge, $b = -5^\circ$) and a high-

(26 Com, $b = 83^\circ$) galactic latitude field, respectively, in order to make statements about the population characteristics of various types of these objects and, after all, about the population differences at different galactic latitudes.

2. The data

2.1. The X-ray data from the ROSAT all-sky survey

The ROSAT all-sky-survey was performed between August 1990 and January 1991 with the position-sensitive proportional counter (PSPC), sensitive in the energy range 0.1–2.4 keV (Trümper 1983). The two fields were scanned by ROSAT over a time period of nearly three to four weeks each: the Com field during Dec. 4–22, 1990 (TJD 48229–48247), and the Sge field during Oct. 7 – Nov. 3, 1990 (TJD 48171–48198). The exposure times in the two fields are 250–500 sec (Com) and 300–590 sec (Sge), respectively, and are typically distributed over 20–30 individual scans. There is an exposure gradient over the fields according to ecliptic latitude.

After extracting the data of the two $10^\circ \times 10^\circ$ fields, the EXSAS package (Zimmermann et al. 1994) was used for the data reduction. The adapted source detection technique consists of several steps: first, all possible sources are identified by means of a “sliding window” technique and removed from the data. This procedure was applied twice with two different sizes of the sliding window in order to account for different coverage of sky regions at different off-axis angles, which leads to varying merged point-spread functions across the fields. Also, the likelihood threshold was set at a very low value to allow many source candidates to join the list. Second, a background map is produced with a bi-cubic spline fit to the resulting image. Finally, a maximum likelihood algorithm is applied to the background-subtracted data (e.g., Cruddace, Hasinger & Schmitt 1988) in three separate pulse-height channel ranges. Each candidate source of the sliding-window list is tested in this way with a likelihood threshold of 10. If a source is detected in more than one energy band, the detection corresponding to a higher likelihood value is considered. As a result, we have detected 238 (26 Com) and 132 (γ Sge) X-ray point sources in the above two fields. The ROSAT source positions provided here are typically accurate to less than a $30''$ radius. The error is dominated by systematic effects, hence independent of the brightness of the sources.

Tables 1a,1b contain the main X-ray data of the ROSAT sources in 26 Com and γ Sge, respectively. Subitem “a” refers to the Com field and subitem “b” to the Sge field.

Column 1: Running number.

Column 2: Coordinates (2000.0) of the ROSAT position in right ascension ($^h, ^m, ^s$) and declination ($^\circ, ', ''$).

Column 3: Name of source when contained in the ROSAT all-sky survey catalogue (RXS; Voges et al. 1999).

Column 4: Statistical error of the RXS position (from the RXS catalogue).

Column 5: Vignetting corrected mean X-ray intensity in PSPC counts/sec.

Column 6+7: X-ray spectral shape, expressed in terms of two hardness ratios HR1 and HR2. The hardness ratio HR1 is defined as the normalized count difference $(N_{50-200} - N_{10-40}) / (N_{10-40} + N_{50-200})$, where N_{a-b} denotes the number of counts in the PSPC between channels a and b. Similarly, the hardness ratio HR2 is defined as $(N_{91-200} - N_{50-90}) / N_{50-200}$. HR1 is sensitive to the Galactic foreground absorbing column.




Fig. 1. Distribution of the X-ray hardness ratios of all detected sources in the two fields (top: Com; bottom: Sge). Sources with more than 20 cts are shown with filled symbols, and spectral parameters are given in Tables 2a,2b, while sources with fewer than 20 cts are shown with open triangles and their error bars. In the bottom panel note the substantial difference in the distribution which is the effect of larger absorption in the Sge field due to its low galactic latitude, i.e. scarcity of sources with negative HR1 and/or with negative HR2. See the text for the definition of HR1 and HR2.

Column 8: The observed (not extinction corrected!) X-ray flux in the 0.1–2.4 keV band in units of 10^{-13} erg/cm²/s. This is a gross underestimate for objects identified as clusters of galaxies (or candidates), as this is derived from point-source PSF-fitting.

Column 9: Most likely optical identification according to various criteria (see text). The symbol is that of the corresponding object in column 1 of Table 5a,5b, except for the Sge “sources” 87-91,93-94 which are detections of flux enhancements of an extended supernova remnant (SNR). A dash means that no optical identification can be proposed, and a question mark after the symbol denotes some doubts due

to inconclusive data or other alternatives. “Cluster” denotes cases where the X-ray emission is more likely associated with the galaxy cluster gas emission rather than individual galaxies in that cluster.

We note that there is not a one-to-one correspondence of sources detected with the above procedure and those published in the ROSAT all-sky survey (RXS) catalogue. This is due to the fact that this program, and in particular the source detection in the ROSAT all-sky survey, had been started in 1993 with the processing state of the data of that time. In contrast, the RXS survey was prepared several years later with various processing improvements. We have no indication that our version suffers any systematic problems which post-facto is proven by the very similar identification rates (see Table 7). The major difference in the source detection is, though, that we used a slightly lower maximum likelihood threshold (8) as compared to that of the RXS catalogue (10), and therefore the present source list is more extensive than the RXS catalogue in these two areas. A noteworthy difference in approach is that the RXS catalogue lists the statistical error for each source, some of which are as small as $8''$. At this size, the positional error is affected by systematic effects of at least a similar level. We therefore start out identification process with a generic $30''$ error circle for all sources.

The distribution of hardness ratios of all sources is shown in Fig. 1, and shows the significant differences caused by the different galactic foreground absorbing column in the two fields: The fraction of soft sources is much lower in the Sge field.

For the brighter X-ray sources (more than 20 counts) in both fields, we have performed spectral fitting using three different models: a power law, a thermal bremsstrahlung and a blackbody model. Together with the absorption by neutral hydrogen which was always left as a free variable, a total of three parameters were fit to the X-ray spectrum. The selection of the “most appropriate” spectral fit was based on the goodness of the fit (reduced χ^2) and the consistency with the expected spectral shape for the type of object according to the most likely identification - thus it was an iterative process. The fit parameters of the selected “most appropriate” model for each X-ray source are reported in Table 2a,2b, and the unfolded photon spectra are shown in Fig. 4a,4b.

Rough distance estimates can be made for the brighter X-ray sources based on the measured hydrogen column density. For the Com field, at galactic latitude $b = +83^\circ$, the foreground column density is in the range $(0.10\text{--}0.26) \times 10^{21} \text{ cm}^{-2}$ (corresponding to $A_V = 0.06\text{--}0.15 \text{ mag}$), and sources with less than this column should be at less than 100 pc distance. In contrast, sources in the Sge field suffer an absorbing column in the range of $(1.3\text{--}12.7) \times 10^{21} \text{ cm}^{-2}$ (corresponding to $A_V = 0.7\text{--}7 \text{ mag}$). While this larger dynamic range allows a rough distance estimate, it also implies that more distant sources are easily absorbed, thus falling below the sensitivity threshold of the ROSAT telescope during the survey exposure.

2.2. The optical plate material

For the field γ Sge we have altogether 239 plates of the 400/1600 mm quadruplet astrographs with $10^\circ \times 10^\circ$ and 87 plates taken with the 400/1950 mm quadruplet astrograph with 8.5° by 8.5° . They cover the time interval 1935 – 1995 more or less continuously. Moreover, for some overlapping zones we have 227 plates of the neighbouring field γ Aql, 192 plates for β Del. Furthermore, for stars brighter than 16 mag we have 193 plates of the Zeiss triplet 170/1400 mm.




Fig. 2. The X-ray to optical flux is plotted over the $B - V$ colour for all objects of the Com field (Table 5a). The straight line marks the upper boundary of the region populated by stars (Stoeck et al. 1991), and the areas populated by various spectral types are taken from Beuermann et al. (1999). Thus, triangles denote secure star identifications. Accretion-powered systems (CVs, AGN) can reach high f_X/f_{opt} values. Since most of those systems have blue optical colours, they populate the upper left corner. The high number of objects reflects the fact that the Com field is at high galactic latitude where the fraction of AGN is high. No extinction correction has been applied, since the correction factor is smaller than a factor of two (Fig. 3). This and the error in $B - V$ are visualized by the cross in the lower right corner.

For the field 26 Com, we have 226 plates taken with the 400/1600 mm astrographs, 204 plates taken with the 400/1950mm astrograph, partly with displaced centre, and 294 plates (1400/1600 mm) of the overlapping neighbouring field 5 Com, covering the time interval 1960–1995. The best Sonneberg plates reach a sensitivity of nearly 18 mag at the plate centre.

Finally, we used the POSS Sky Survey prints, and in some cases plates of the Tautenburg 2 m Schmidt telescope with a limiting magnitude of $\sim 21 \text{ mag}$, at the best.

2.3. Swift X-ray observations

In the process of this study it became clear that there was some fraction of sources for which a unique identification was not possible. This was particularly true for the Sge field, where often more than one optically variable object was located within the ROSAT error circle. We therefore proposed and obtained short Swift (Gehrels et al. 2004) X-ray observations to obtain a more precise X-ray position. Typically, the X-ray telescope (XRT; Burrows et al. 2005) onboard Swift provides positions at the $5''$ accuracy level for bright sources, which degrades somewhat for faint sources. The exposure time has been chosen to provide a secure detection even if the source dropped by a factor of three in X-ray intensity. In general, however, the number of photons detected for these sources is not sufficient for deriving a proper X-ray spectrum.

2.4. *SDSS-III spectra*

At a very late stage of this paper, when practically all analysis was already finished, the DR9 (IIIrd phase of the Sloan Digital Sky Survey, SDSS) release contained spectra of about 130 sources in the Coma field. Obviously, this resource has been used.

Table 1a. ROSAT X-ray sources in the Com field. All positions have an error radius of 30''. The ID column provides cross-reference to tables 5a,5b and Figs. 40a,40b.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl (° ' ")	RXS J Name	RXS Err (")	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
1	12 33 55.9 +26 02 41	123356.4+260249	16	0.040	-0.31±0.34	0.12±0.63	3.15	a?
2	12 36 17.8 +25 59 00	-	-	0.042	0.80±0.19	0.37±0.23	4.64	a
3	12 54 09.1 +25 55 26	-	-	0.137	-0.60±0.12	0.09±0.33	11.50	a
4	12 23 08.9 +25 50 53	122308.9+255058	18	0.051	0.42±0.27	0.39±0.27	12.50	a
5	12 24 02.4 +25 51 05	-	-	0.012	0.27±0.47	0.49±0.43	1.28	a
6	12 22 30.7 +25 50 25	-	-	0.680	0.12±0.06	0.09±0.08	57.47	a
7	12 41 36.0 +25 52 44	124136.3+255244	15	0.050	-0.86±0.14	-1.00±0.00	3.24	a
8	12 28 12.6 +25 49 38	-	-	0.068	-0.39±0.19	0.01±0.37	1.97	a
9	12 20 44.7 +25 46 05	122045.4+254609	19	0.021	-0.42±0.36	-1.00±0.00	1.63	a
10	12 24 11.1 +25 43 27	122411.4+254330	13	0.043	-0.54±0.25	-1.00±0.00	2.37	a
11	12 35 16.7 +25 42 44	123517.8+254233	18	0.015	0.11±0.40	-0.05±0.46	1.63	—
12	12 53 27.5 +25 34 05	125328.0+253407	17	0.045	-0.68±0.19	-0.61±0.58	3.51	a
13	12 25 01.8 +25 33 37	-	-	0.615	0.05±0.06	-0.11±0.08	49.80	a
14	12 17 55.9 +25 29 06	121756.9+252919	18	0.013	0.64±0.57	1.00±0.00	1.43	b
15	12 59 54.9 +25 27 59	125955.5+252805	13	0.036	-0.03±0.29	0.62±0.49	2.80	a
16	12 51 39.3 +25 30 46	125139.3+253043	12	0.039	-0.24±0.31	0.20±0.42	3.04	a
17	12 50 27.2 +25 30 12	-	-	0.038	-0.79±0.22	-0.38±1.63	2.97	a
18	12 39 43.1 +25 30 52	123943.6+253054	16	0.027	-0.57±0.31	-1.00±0.00	2.11	a
19	12 54 14.6 +25 24 55	125415.0+252457	15	0.039	-0.38±0.25	-0.21±0.46	3.07	a?
20	12 40 10.8 +25 21 33	124011.4+252133	17	0.012	1.00±0.00	0.25±0.48	1.33	Cluster?
21	12 59 23.8 +25 15 32	125924.8+251531	22	0.026	-0.43±0.33	0.18±0.68	2.03	a?
22	12 30 14.0 +25 18 01	-	-	0.241	-0.10±0.11	0.14±0.15	22.53	a
23	12 38 45.6 +25 17 20	-	-	0.012	1.00±0.00	-0.82±0.78	1.34	—
24	12 30 43.5 +25 14 50	123043.3+251455	17	0.032	-0.21±0.32	-0.66±0.63	2.54	a
25	12 46 39.8 +25 12 29	124639.8+251247	27	0.027	0.21±0.31	-0.02±0.32	2.99	b
26	12 26 04.6 +25 06 44	122604.6+250648	13	0.033	-0.56±0.31	-0.22±0.70	2.62	a
27	12 36 50.9 +25 07 49	-	-	0.114	-0.20±0.15	0.12±0.24	11.27	a
28	13 00 44.0 +24 58 31	-	-	0.026	-0.61±0.27	0.53±0.57	2.06	a?
29	12 40 54.1 +25 00 53	124054.3+250059	15	0.041	0.01±0.31	0.64±0.37	4.61	a
30	12 25 39.9 +24 58 33	-	-	0.147	-0.33±0.13	0.11±0.25	15.11	a
31	12 56 03.5 +24 51 42	125603.7+245147	16	0.031	-0.36±0.33	0.05±0.58	2.47	a
32	12 49 58.0 +24 52 58	-	-	0.012	1.00±0.00	-0.21±0.53	1.31	a
33	12 45 20.0 +24 52 46	124518.3+245257	26	0.017	1.00±0.00	0.38±0.40	1.88	Cluster
34	12 30 15.0 +24 51 57	123015.2+245155	13	0.032	-0.61±0.27	1.00±0.00	2.50	a
35	12 42 06.0 +24 52 26	124206.5+245224	16	0.041	-0.89±0.17	1.00±0.00	3.23	a
36	12 48 47.7 +24 50 27	124848.1+245031	10	0.142	-0.52±0.12	0.17±0.27	4.40	a
37	12 43 08.7 +24 47 16	124310.2+244714	22	0.032	-0.16±0.36	0.28±0.48	2.53	a
38	12 25 57.5 +24 38 11	122557.5+243817	10	0.099	-0.72±0.13	-0.40±0.49	2.68	a
39	12 35 33.7 +24 39 11	123533.9+243915	12	0.051	0.08±0.27	0.29±0.33	0.99	a
40	12 24 24.0 +24 36 08	-	-	0.172	0.01±0.13	0.00±0.17	12.00	a
41	12 55 39.1 +24 30 14	125539.2+243021	13	0.040	-0.25±0.32	0.66±0.61	3.13	a
42	12 56 04.8 +24 29 19	125607.0+242909	17	0.048	-0.31±0.25	-0.45±0.38	8.69	a
43	12 29 41.6 +24 31 17	-	-	0.197	-0.21±0.11	-0.24±0.17	20.85	a
44	12 58 46.7 +24 27 37	125847.1+242740	11	0.047	-0.56±0.20	-0.04±0.52	4.97	a
45	12 39 18.3 +24 31 43	-	-	0.260	-0.75±0.07	-0.26±0.30	17.17	a
46	12 53 07.4 +24 27 49	125307.1+242749	23	0.020	-0.82±0.29	1.00±0.00	1.61	a
47	12 22 14.4 +24 23 58	122215.4+242405	24	0.012	1.00±0.00	0.65±0.32	1.37	a
48	12 40 09.1 +24 25 32	-	-	0.042	0.90±0.30	-0.05±0.25	4.66	a
49	12 24 30.2 +24 20 53	122430.5+242103	15	0.025	-0.05±0.39	-0.06±0.55	1.98	a
50	12 22 18.5 +24 19 01	124434.2+242009	11	0.043	0.79±0.26	0.52±0.21	3.14	Cluster?
51	12 44 34.2 +24 20 11	-	-	0.008	-0.39±0.31	0.48±0.46	0.66	a
52	12 57 31.4 +24 12 44	-	-	0.929	-0.09±0.05	-0.03±0.07	70.28	b
53	12 56 36.2 +24 12 45	-	-	0.021	-1.00±0.00	0.00±0.00	1.67	a
54	12 48 39.4 +24 13 02	124838.0+241314	32	0.027	-1.00±0.00	0.00±0.00	2.09	a
55	12 27 10.5 +24 09 35	122711.0+240940	14	0.039	-0.47±0.28	-0.01±0.58	3.04	a
56	12 51 42.0 +24 04 34	125141.9+240444	12	0.052	0.03±0.23	0.28±0.30	8.64	a
57	12 58 51.0 +23 55 33	-	-	0.436	-0.38±0.07	0.05±0.13	28.75	a
58	12 28 19.0 +23 57 49	122819.4+235758	11	0.033	-0.24±0.31	-0.00±0.47	2.59	a

Table 1a. continued.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl ([°] ' ")	RXS J Name	RXS Err (")	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
59	12 21 59.7 +23 54 21	122159.9+235434	25	0.020	0.30±0.39	0.68±0.53	2.20	Cluster
60	12 37 38.3 +23 52 43	123738.4+235242	20	0.019	-1.00±0.00	0.00±0.00	1.54	–
61	12 30 54.8 +23 51 56	123055.4+235202	14	0.037	-0.07±0.31	0.11±0.41	2.93	a
62	12 34 38.7 +23 50 10	-	-	0.186	-0.13±0.12	0.14±0.18	14.24	e
63	12 32 40.7 +23 47 58	-	-	0.084	0.04±0.20	0.40±0.26	15.00	a
64	12 52 59.0 +23 43 22	125259.1+234320	19	0.027	-0.82±0.19	1.00±0.00	2.16	a?
65	12 33 50.5 +23 37 45	123350.7+233745	20	0.030	-0.15±0.39	0.04±0.51	2.32	a
66	12 25 32.3 +23 35 39	122532.4+233551	12	0.030	0.01±0.33	0.31±0.44	3.30	–
67	12 35 16.8 +23 34 41	123516.9+233447	14	0.087	-1.00±0.00	0.00±0.00	4.63	a
68	12 58 07.3 +23 29 41	125807.3+232902	18	0.048	-0.94±0.12	-1.00±0.00	2.40	a
69	12 56 52.4 +23 29 52	-	-	0.071	-0.38±0.21	-0.04±0.39	7.30	a?
70	12 43 14.7 +23 31 41	124315.4+233141	18	0.026	-0.41±0.35	-0.13±0.63	2.05	b
71	12 37 58.1 +23 31 21	-	-	0.015	1.00±0.00	-0.94±0.87	1.67	Cluster?
72	12 37 29.5 +23 29 18	-	-	0.033	-0.72±0.25	0.77±0.73	2.62	–
73	12 46 23.5 +23 27 22	124625.6+232741	20	0.023	0.27±0.31	-0.00±0.32	2.59	a?
74	13 00 33.1 +23 20 48	130033.6+232050	17	0.038	-0.32±0.29	0.31±0.61	3.02	a
75	12 25 59.2 +23 21 28	122559.4+232135	10	0.081	-0.03±0.20	0.45±0.25	9.86	a
76	12 20 50.9 +23 19 16	-	-	0.012	0.36±0.50	-0.38±0.43	1.39	a?
77	12 41 20.2 +23 22 03	-	-	0.026	-0.28±0.38	0.16±0.65	2.08	a
78	12 41 48.9 +23 19 35	124149.3+231939	12	0.031	-0.41±0.28	-0.07±0.56	2.41	b
79	12 38 52.2 +23 16 36	123852.0+231647	13	0.027	-0.63±0.26	0.35±0.87	2.13	a
80	12 25 22.5 +23 13 43	122522.5+231348	14	0.040	-0.31±0.27	0.39±0.40	3.16	a
81	12 46 45.3 +23 12 35	124645.7+231240	14	0.019	-0.36±0.40	-1.00±0.00	1.50	a?
82	12 40 43.1 +23 10 09	124043.3+231021	24	0.033	-0.16±0.35	1.00±0.00	2.58	a
83	12 41 27.1 +23 09 56	124127.7+231004	17	0.027	-0.59±0.29	-1.00±0.00	2.15	a
84	12 36 45.1 +23 09 48	123646.0+230955	22	0.031	-1.00±0.00	0.00±0.00	2.41	a
85	12 57 30.1 +23 01 54	125733.4+230213	32	0.027	-1.00±0.00	0.00±0.00	2.14	–
86	12 34 43.3 +23 01 29	-	-	0.030	0.74±0.36	-0.26±0.29	3.30	–
87	12 29 26.4 +22 59 40	-	-	0.058	0.31±0.19	-0.08±0.20	2.40	a
88	13 00 31.8 +22 55 22	-	-	0.009	0.66±0.49	-0.14±0.58	1.06	a
89	12 17 50.1 +22 55 25	121748.5+225527	30	0.023	1.00±0.00	0.30±0.30	2.50	Cluster?
90	12 18 04.0 +22 48 26	121803.5+224758	38	0.019	1.00±0.00	0.13±0.39	2.06	Cluster?
91	12 40 07.2 +22 52 18	-	-	0.013	1.00±0.00	0.45±0.44	1.47	Cluster?
92	12 21 21.7 +22 46 57	122121.1+224729	19	0.018	0.39±0.42	-0.35±0.41	2.03	a
93	12 29 12.7 +22 49 03	-	-	0.012	0.77±0.49	-0.11±0.50	1.38	a?
94	12 53 36.2 +22 47 37	-	-	0.075	-0.12±0.21	0.15±0.30	8.13	a
95	12 31 56.1 +22 48 06	123156.5+224814	14	0.033	-0.19±0.34	-0.14±0.54	2.62	a
96	12 23 53.5 +22 45 41	-	-	0.013	0.53±0.66	0.25±0.52	1.46	a
97	12 30 36.6 +22 41 58	123036.5+224209	20	0.025	-0.78±0.32	-1.00±0.00	1.94	a
98	12 24 00.1 +22 39 14	122400.3+223920	19	0.028	-1.00±0.00	0.00±0.00	2.21	b
99	12 47 05.1 +22 37 35	124707.7+223743	39	0.025	-1.00±0.00	0.00±0.00	1.95	a
100	12 55 05.0 +22 32 56	125505.2+223302	17	0.016	0.62±0.60	0.22±0.43	1.79	a
101	12 52 30.4 +22 33 32	125230.3+223329	25	0.033	1.00±0.00	0.04±0.28	3.63	b
102	12 26 29.5 +22 33 21	122631.6+223321	28	0.024	0.22±0.37	-0.29±0.33	2.63	b
103	12 52 11.3 +22 33 09	125211.6+223315	12	0.047	0.20±0.24	0.45±0.29	6.62	–
104	12 51 46.5 +22 32 36	-	-	0.413	-0.20±0.08	0.00±0.12	26.57	a
105	12 40 00.0 +22 33 27	124000.4+223335	9	0.130	-0.22±0.15	-0.46±0.20	12.33	a
106	12 43 07.4 +22 26 43	124308.1+222659	28	0.015	0.98±0.53	0.20±0.41	1.67	–
107	12 35 15.2 +22 25 53	123516.0+222604	16	0.029	0.09±0.33	0.32±0.44	3.19	a
108	12 25 19.0 +22 20 14	122520.0+222011	18	0.016	1.00±0.00	-0.04±0.42	1.81	a
109	12 27 33.6 +22 14 17	122734.1+221423	12	0.045	-0.83±0.15	-1.00±0.00	4.56	a
110	12 25 14.0 +22 13 07	122514.1+221315	18	0.015	0.25±0.42	0.53±0.50	1.62	a
111	12 43 12.2 +22 12 44	124312.6+221255	12	0.014	0.46±0.38	-0.34±0.59	1.60	a
112	12 54 39.3 +22 06 30	-	-	0.216	-0.30±0.10	-0.05±0.18	20.66	a
113	12 57 01.9 +22 01 50	-	-	5.662	-1.00±0.00	0.00±0.00	280.80	a
114	12 28 08.0 +22 03 22	122808.4+220329	16	0.015	1.00±0.00	-0.47±0.38	1.71	a
115	12 28 36.9 +22 01 04	122837.7+220044	20	0.013	0.51±0.51	0.39±0.45	1.44	a?
116	12 36 26.7 +22 01 54	123627.0+220200	16	0.033	-0.33±0.29	-0.08±0.49	2.56	a
117	12 44 22.2 +21 59 33	124422.6+215938	8	0.134	-0.15±0.14	-0.11±0.20	11.03	a
118	12 30 12.5 +21 58 00	123013.7+215812	13	0.035	0.45±0.24	-0.10±0.28	3.85	a

Table 1a. continued.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl ([°] ['] ^{''})	RXS J Name	RXS Err (^{''})	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
119	12 26 58.0 +21 57 03	122658.2+215706	22	0.031	-0.23±0.36	-0.24±0.72	2.43	a
120	12 26 48.9 +21 52 57	122649.3+215301	14	0.063	-0.11±0.24	0.08±0.33	5.89	—
121	12 35 35.5 +21 53 48	123535.7+215403	28	0.013	1.00±0.00	-0.51±0.49	1.41	a
122	12 40 08.9 +21 46 14	124008.1+214630	17	0.020	0.63±0.32	0.40±0.34	2.19	a?
123	12 56 43.0 +21 41 07	-	-	0.034	0.46±0.35	-0.04±0.35	3.74	a
124	12 22 12.1 +21 37 41	122212.3+213747	10	0.108	0.14±0.16	-0.23±0.20	9.12	a
125	13 00 16.4 +21 33 08	130016.4+213313	16	0.024	0.65±0.33	0.16±0.32	2.68	a
126	12 29 33.9 +21 37 03	122934.1+213707	16	0.037	-0.48±0.25	1.00±0.00	2.89	a
127	12 32 13.5 +21 37 07	123213.9+213710	12	0.051	-0.27±0.26	0.23±0.41	11.98	a
128	12 56 28.0 +21 31 09	125628.4+213114	11	0.033	1.00±0.00	0.48±0.22	3.71	a
129	12 34 15.7 +21 28 52	123416.3+212858	20	0.023	-0.84±0.19	0.60±0.39	1.84	—
130	12 51 20.0 +21 25 24	125120.0+212526	13	0.012	1.00±0.00	0.80±0.41	1.30	a
131	12 44 31.2 +21 25 53	124431.7+212559	17	0.020	0.93±0.45	0.01±0.35	2.28	a
132	12 33 20.5 +21 25 24	123320.9+212531	11	0.035	0.30±0.26	-0.80±0.19	3.88	a
133	12 37 35.4 +21 25 21	-	-	0.018	0.89±2.23	0.90±0.10	2.05	a
134	12 24 54.6 +21 22 45	-	-	0.180	0.04±0.12	-0.81±0.15	22.05	a
135	12 58 59.3 +21 12 52	-	-	0.029	-0.46±0.32	0.35±0.71	2.28	a
136	12 53 18.1 +21 14 22	125318.2+211425	15	0.023	-0.64±0.30	1.00±0.00	1.77	a
137	12 44 57.4 +21 14 10	124457.6+211416	14	0.029	1.00±0.00	-0.12±0.53	3.23	a
138	12 38 09.8 +21 14 03	-	-	0.014	1.00±0.00	0.63±0.18	1.57	a?
139	12 19 05.3 +21 06 30	121905.3+210633	10	0.113	0.27±0.17	-0.67±0.26	11.23	a
140	12 34 28.1 +21 09 34	123428.5+210942	21	0.028	-0.59±0.34	1.00±0.00	2.17	a
141	12 39 31.3 +21 08 16	-	-	0.051	1.00±0.00	0.11±0.27	6.42	a
142	12 32 30.1 +21 05 59	123230.5+210602	11	0.028	1.00±0.00	0.50±0.32	3.07	a
143	12 53 40.4 +21 01 36	125340.8+210137	15	0.018	0.60±0.51	0.86±0.14	1.95	a
144	12 25 42.0 +20 55 00	-	-	0.327	-0.67±0.07	-0.21±0.23	21.37	a
145	12 32 09.1 +20 55 19	123209.5+205523	12	0.029	0.77±0.23	-1.00±0.00	3.23	a
146	12 20 29.5 +20 46 48	-	-	0.010	1.00±0.00	-0.60±0.65	1.08	—
147	12 40 08.5 +20 48 06	124008.9+204814	24	0.027	-0.13±0.35	-0.16±0.40	2.10	a
148	12 42 44.2 +20 46 36	124244.6+204638	14	0.013	1.00±0.00	0.18±0.47	1.41	a
149	12 18 03.1 +20 38 21	-	-	0.014	1.00±0.00	0.45±0.34	1.54	Cluster?
150	12 32 46.9 +20 37 12	123247.0+203730	20	0.027	0.26±0.30	-0.39±0.40	3.05	a
151	12 50 46.0 +20 35 58	125046.9+203601	24	0.015	0.75±0.37	-0.12±0.38	1.70	c
152	12 23 59.0 +20 31 38	122359.7+203146	25	0.016	1.00±0.00	0.65±0.21	1.78	a
153	12 59 25.6 +20 29 15	125926.1+202914	12	0.146	-0.21±0.15	-0.50±0.33	17.13	a
154	12 50 42.8 +20 32 13	125043.5+203216	17	0.035	-0.39±0.29	0.36±0.42	2.75	a
155	12 30 52.1 +20 30 02	123052.8+203006	16	0.050	-0.23±0.25	0.26±0.40	3.46	a
156	12 57 40.6 +20 24 03	125741.1+202359	14	0.017	1.00±0.00	0.12±0.45	1.86	a
157	12 51 51.5 +20 22 47	125151.8+202253	10	0.045	-0.37±0.22	0.38±0.26	6.42	a
158	12 32 26.2 +20 23 29	-	-	0.130	-0.23±0.15	0.58±0.12	2.44	a
159	12 32 03.6 +20 09 27	123203.6+200930	8	0.607	-0.35±0.06	-0.67±0.13	45.74	a
160	12 21 05.3 +20 05 38	122105.7+200541	11	0.050	0.24±0.25	-0.43±0.46	5.98	a
161	12 18 52.2 +19 58 50	121852.6+195853	21	0.029	0.20±0.30	-1.00±0.00	3.22	a
162	12 35 31.8 +19 54 36	-	-	0.018	-0.29±0.42	0.42±0.32	1.44	a
163	12 39 44.7 +19 54 18	123945.2+195426	22	0.020	0.45±0.43	-1.00±0.00	2.24	a
164	12 49 29.0 +19 52 40	124929.0+195244	19	0.022	-1.00±0.00	1.00±0.00	1.76	a?
165	12 26 44.0 +19 50 07	122644.2+195009	25	0.020	1.00±0.00	-0.16±0.37	2.26	Cluster
166	12 41 47.6 +19 49 13	-	-	0.019	1.00±0.00	-0.55±0.64	2.10	Cluster
167	12 36 19.2 +19 48 18	-	-	0.028	1.00±0.00	0.26±0.20	3.15	Cluster
168	12 42 22.6 +19 47 07	124223.0+194709	12	0.095	0.07±0.18	-0.75±0.24	16.54	a
169	12 19 57.9 +19 41 55	121958.0+194154	10	0.103	-0.70±0.13	0.20±0.27	11.60	a
170	12 53 02.9 +19 36 15	125303.4+193618	8	0.133	-0.06±0.15	-0.70±0.22	9.10	a
171	12 55 28.5 +19 33 24	125528.5+193326	26	0.049	-0.03±0.30	-0.12±0.36	3.28	a
172	12 28 54.3 +19 26 06	122852.0+192616	13	0.017	1.00±0.00	-0.05±0.44	1.93	Cluster
173	12 41 19.9 +19 25 16	124119.0+192525	28	0.013	1.00±0.00	0.64±0.27	1.47	b?
174	12 50 14.8 +19 23 53	125015.2+192357	9	0.177	-0.66±0.10	-1.00±0.00	15.13	a
175	12 45 38.4 +19 20 50	-	-	0.021	-0.86±0.42	0.83±0.32	1.68	a
176	12 36 07.5 +19 19 42	123607.9+191935	22	0.021	1.00±0.00	0.05±0.53	2.37	a
177	12 28 54.7 +19 17 34	122855.4+191740	21	0.017	1.00±0.00	-0.05±0.39	1.89	—
178	12 50 43.8 +19 14 34	125044.3+191437	15	0.024	-0.07±0.40	-1.00±0.00	1.90	a?

Table 1a. continued.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl ([°] ' ")	RXS J Name	RXS Err (")	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
179	12 18 47.2 +19 10 03	121847.5+190959	16	0.027	-0.74±0.24	0.38±0.45	2.13	a
180	12 56 32.1 +19 07 48	125632.8+190750	17	0.013	1.00±0.00	-1.00±0.00	1.49	a?
181	12 47 49.3 +19 01 23	124749.5+190127	11	0.056	-0.53±0.19	0.23±0.29	2.90	a
182	12 25 45.7 +18 58 48	122546.6+185839	17	0.025	0.48±0.32	0.37±0.36	2.80	c
183	12 43 47.3 +18 48 52	124347.3+184855	11	0.092	-0.45±0.17	0.14±0.38	3.56	a
184	12 52 37.7 +18 44 48	125238.1+184449	15	0.036	-0.44±0.31	1.00±0.00	2.81	a?
185	12 18 29.7 +18 35 28	121829.6+183526	13	0.028	-0.17±0.45	-1.00±0.00	2.20	a
186	12 49 17.8 +18 38 21	124918.1+183822	16	0.031	-0.42±0.28	-0.18±0.39	2.48	a?
187	12 48 07.1 +18 38 12	-	-	0.048	-0.45±0.27	0.88±0.10	3.66	a
188	12 41 17.1 +18 34 44	-	-	0.099	0.49±0.16	-0.88±0.09	37.73	b
189	12 46 59.6 +18 33 32	124659.6+183329	22	0.015	1.00±0.00	-0.30±0.42	1.68	Cluster?
190	12 18 00.5 +18 27 12	121800.8+182711	15	0.031	-0.03±0.48	0.60±0.44	2.41	—
191	12 21 55.2 +18 23 45	122155.5+182349	21	0.013	1.00±0.00	0.75±0.19	1.39	—
192	12 19 05.9 +18 21 21	-	-	0.073	0.45±0.16	-1.00±0.00	14.49	a
193	12 23 49.2 +18 20 16	-	-	0.018	1.00±0.00	-0.08±0.32	2.05	—
194	12 38 10.9 +18 22 00	-	-	0.021	0.30±0.38	0.72±0.15	2.29	—
195	12 29 40.2 +18 16 48	-	-	0.433	-0.35±0.07	-0.36±0.14	33.47	—
196	12 48 14.1 +18 16 44	124813.8+181648	11	0.061	0.35±0.18	0.31±0.19	12.37	a
197	12 47 21.8 +18 14 34	124721.8+181436	11	0.178	-0.04±0.12	-0.92±0.18	20.89	a
198	12 41 42.1 +18 11 01	124142.3+181101	28	0.029	-0.46±0.34	0.67±0.24	2.26	—
199	12 35 50.0 +18 02 47	123549.0+180252	53	0.099	-0.27±0.17	0.20±0.33	4.78	a
200	12 48 03.5 +18 00 05	124803.8+180009	11	0.023	0.89±0.35	0.23±0.43	2.52	a
201	12 54 47.3 +17 56 07	125447.5+175617	22	0.017	1.00±0.00	-1.00±0.00	1.90	b
202	12 50 35.9 +17 57 08	125036.2+175704	12	0.012	1.00±0.00	0.27±0.30	1.32	a
203	12 44 07.3 +17 47 59	124407.5+174757	14	0.026	0.73±0.39	0.57±0.19	2.87	a?
204	12 44 58.7 +17 32 59	-	-	0.073	0.30±0.17	-0.70±0.18	6.67	a
205	12 35 00.1 +17 32 07	-	-	0.015	1.00±0.00	-0.47±0.62	1.69	a
206	12 43 29.4 +17 29 58	-	-	0.035	-0.57±0.34	0.30±0.66	2.74	a?
207	12 31 21.1 +17 21 48	-	-	0.027	-0.58±0.39	0.00±0.00	2.14	a?
208	12 24 22.9 +17 19 45	122423.2+171936	20	0.020	-1.00±0.00	1.00±0.00	1.59	—
209	12 44 10.4 +17 21 08	124410.2+172050	12	0.039	-0.29±0.28	0.82±0.33	3.11	a
210	12 38 50.2 +17 16 29	-	-	0.032	0.33±0.28	-0.63±0.36	3.54	—
211	12 42 24.6 +17 14 24	-	-	0.022	-0.30±0.42	0.36±0.47	1.76	a?
212	12 46 10.9 +17 10 45	124613.9+171048	33	0.014	1.00±0.00	0.23±0.47	1.59	a?
213	12 52 22.0 +17 06 28	125222.5+170629	10	0.019	0.52±0.41	0.45±0.24	2.11	a
214	12 28 29.5 +17 05 13	122829.4+170515	14	0.035	0.75±0.25	0.77±0.12	3.94	a
215	12 33 05.4 +17 01 34	-	-	0.272	0.13±0.10	-0.96±0.17	17.28	a
216	12 35 28.5 +17 00 27	-	-	0.024	0.06±0.42	0.20±0.36	2.68	a
217	12 54 40.3 +16 56 07	125441.8+165617	19	0.016	1.00±0.00	0.61±0.28	1.81	Cluster
218	12 32 04.5 +16 55 29	-	-	0.015	0.09±0.35	1.00±0.00	1.66	a
219	12 41 32.9 +16 54 44	-	-	0.020	0.66±0.34	0.34±0.38	2.20	—
220	12 44 03.9 +16 52 16	-	-	0.019	1.00±0.00	-1.00±0.00	2.10	Cluster
221	12 37 26.6 +16 50 33	-	-	0.044	-0.88±0.14	0.27±0.54	10.98	—
222	12 48 02.8 +16 48 11	-	-	0.017	0.76±0.46	0.58±0.19	1.90	a
223	12 29 44.0 +16 40 01	122943.5+164007	14	0.064	0.40±0.19	0.32±0.18	12.48	c
224	12 19 06.2 +16 32 53	-	-	0.259	-0.26±0.10	-0.79±0.18	18.86	a
225	12 59 52.0 +16 30 32	125956.0+163127	14	0.035	0.08±0.62	0.80±0.57	3.93	b?
226	12 19 46.7 +16 30 57	121947.7+163103	14	0.022	1.00±0.00	-0.48±0.55	2.45	a
227	12 57 37.6 +16 30 36	125737.6+163038	25	0.030	-0.13±0.30	0.62±0.31	2.38	a
228	12 41 47.3 +16 32 33	124155.4+163056	19	0.030	1.00±0.00	0.37±0.37	3.31	Cluster
229	12 34 37.0 +16 31 59	123439.0+163149	64	0.135	-0.47±0.14	0.21±0.24	10.96	b?
230	12 36 24.2 +16 32 05	-	-	0.056	0.44±0.22	-0.41±0.27	26.31	a?
231	12 43 57.3 +16 22 55	-	-	0.030	0.82±0.47	-0.35±0.26	3.35	a
232	12 55 01.4 +16 15 40	125502.3+161542	14	0.031	-0.07±0.37	1.00±0.00	2.45	—
233	12 42 52.6 +16 18 03	-	-	0.025	-0.46±0.36	0.48±0.55	2.00	a?
234	12 34 39.1 +16 06 06	-	-	0.015	0.66±0.47	-0.26±0.52	1.71	a?
235	12 35 45.5 +16 06 05	-	-	0.038	-0.56±0.25	0.67±0.26	3.01	a
236	12 39 44.8 +16 06 02	123946.4+160603	33	0.052	0.35±0.30	-1.00±0.00	5.77	d
237	12 23 45.9 +16 00 34	-	-	0.090	0.41±0.77	0.25±0.41	3.13	—
238	12 32 36.4 +16 02 44	-	-	0.114	-0.28±0.20	0.00±0.00	1.36	a

Table 1b. ROSAT X-ray sources in the Sge field. All positions have an error radius of 30".

No	Coordinates (2000.0) RA (^h ^m ^s) Decl ([°] ['] ["])	RXS J Name	RXS Err (["])	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
1	19 37 42.7 +24 15 13	193742.4+241452	11	0.037	0.65±0.24	-0.42±0.26	4.09	b
2	19 56 28.8 +23 54 57	195628.7+235445	17	0.014	1.00±0.00	0.87±0.41	1.57	a
3	19 58 00.1 +23 54 05	195759.6+235356	17	0.060	0.37±0.18	0.48±0.17	4.10	a?
4	20 07 02.5 +23 52 46	200702.4+235242	13	0.010	0.21±0.41	0.05±0.46	1.16	a
5	19 53 13.8 +23 51 08	195314.1+235110	18	0.023	1.00±0.00	0.16±0.36	2.59	a
6	20 20 03.8 +23 38 11	202003.8+233818	10	0.135	-0.30±0.12	-0.40±0.20	11.55	a
7	19 45 51.9 +23 30 27	194552.5+233015	16	0.048	0.45±0.20	-0.03±0.26	8.37	a
8	19 57 26.3 +23 28 43	195726.6+232844	10	0.027	0.26±0.31	0.10±0.31	2.96	—
9	19 38 39.0 +23 20 45	193838.7+232037	20	0.012	-1.00±0.00	0.00±0.00	0.96	a?
10	19 43 12.7 +23 19 48	-	-	0.025	1.00±0.00	1.00±0.00	2.78	e
11	20 13 52.5 +23 17 37	201351.6+231724	23	0.025	0.18±0.42	1.00±0.00	2.76	a
12	20 15 32.8 +23 12 56	-	-	0.012	1.00±0.00	1.00±0.00	1.34	a?
13	20 17 39.5 +23 09 48	201739.1+230950	11	0.044	0.20±0.23	0.40±0.29	5.17	a
14	20 18 05.7 +23 04 02	-	-	0.014	1.00±0.00	-0.10±0.42	1.56	a
15	20 10 06.9 +23 04 16	-	-	0.022	-0.37±0.38	1.00±0.00	1.76	a
16	20 19 49.2 +22 56 21	-	-	0.363	-0.24±0.07	-0.25±0.12	26.12	a
17	20 03 23.1 +22 59 32	-	-	0.016	0.77±0.35	-0.07±0.45	1.80	c
18	19 52 12.7 +22 52 41	-	-	0.022	0.34±0.42	-0.30±0.44	2.47	a?
19	19 59 36.4 +22 43 06	195936.2+224309	9	0.328	-1.00±0.00	-1.00±0.00	11.42	a
20	20 00 43.3 +22 42 41	200043.3+224240	13	0.115	-0.60±0.14	-0.37±0.50	9.74	a
21	20 05 41.4 +22 39 49	200542.0+223955	10	9.255	-0.98±0.00	0.01±0.14	257.10	a
22	20 07 31.3 +22 34 29	-	-	0.011	1.00±0.00	-0.42±0.44	1.24	b?
23	19 47 35.0 +22 28 58	194735.5+222849	10	0.125	-0.04±0.15	0.05±0.21	8.20	a
24	20 02 48.3 +22 28 33	-	-	0.021	1.00±0.00	0.65±0.27	2.28	a
25	20 07 48.6 +22 26 01	200748.9+222601	18	0.022	-0.25±0.34	-0.37±0.44	1.70	—
26	20 10 48.7 +22 21 41	201049.0+222142	12	0.032	0.66±0.24	0.08±0.27	3.55	a?
27	20 01 51.4 +22 17 00	-	-	0.012	1.00±0.00	0.14±0.45	1.29	a
28	20 14 04.7 +22 13 24	201405.1+221333	11	0.069	0.24±0.22	0.32±0.24	7.06	a
29	20 16 19.6 +22 06 15	-	-	0.035	-0.74±0.22	-0.03±0.99	2.77	a
30	19 49 59.2 +22 04 26	195000.5+220334	40	0.015	1.00±0.00	0.65±0.49	1.67	—
31	20 01 26.2 +22 03 38	-	-	0.023	0.40±0.44	-0.09±0.41	2.50	a?
32	20 11 36.9 +21 58 51	-	-	0.009	1.00±0.00	-0.42±0.47	1.02	b?
33	20 08 22.0 +21 47 55	-	-	0.020	-0.15±0.40	-0.25±0.51	1.54	a?
34	20 09 24.2 +21 42 55	-	-	0.012	1.00±0.00	1.00±0.00	1.33	a
35	19 59 30.4 +21 33 35	195930.4+213336	14	0.062	0.21±0.24	0.32±0.27	14.28	a
36	19 55 37.8 +21 29 51	-	-	0.016	1.00±0.00	-0.47±0.44	1.78	—
37	20 19 06.5 +21 21 12	-	-	0.062	0.20±0.23	0.24±0.29	7.59	a?
38	19 43 55.5 +21 17 53	194356.1+211731	27	0.066	0.91±0.12	1.00±0.00	26.13	a?
39	20 07 55.2 +21 16 01	-	-	0.012	1.00±0.00	0.37±0.44	1.29	—
40	19 53 05.4 +21 14 31	-	-	0.024	1.00±0.00	-0.17±0.39	2.70	a
41	20 13 43.3 +21 07 25	-	-	0.012	0.21±0.48	-0.20±0.45	1.31	a
42	20 06 24.8 +21 07 11	-	-	0.019	1.00±0.00	0.64±0.28	2.06	a
43	20 08 30.6 +21 05 00	-	-	0.015	1.00±0.00	0.35±0.38	1.72	a?
44	20 12 03.5 +20 57 43	-	-	0.021	-0.23±0.39	-1.00±0.00	1.66	a?
45	20 09 56.1 +20 54 43	-	-	0.064	-0.52±0.18	-0.51±0.33	5.27	a
46	19 59 21.0 +20 55 30	-	-	0.015	1.00±0.00	0.66±0.71	1.64	—
47	20 18 59.5 +20 51 15	-	-	0.028	0.94±0.33	0.17±0.31	3.08	a
48	19 53 11.0 +20 44 13	-	-	0.023	-1.00±0.00	0.00±0.00	1.84	h
49	20 06 48.2 +20 41 15	200647.7+204135	20	0.067	0.19±0.22	0.13±0.27	5.28	a
50	19 41 08.2 +20 08 52	194108.7+200848	16	0.027	0.23±0.35	0.60±0.32	3.02	b
51	20 17 42.7 +20 08 02	-	-	0.016	1.00±0.00	1.00±0.00	1.83	—
52	19 57 02.6 +20 05 14	-	-	0.039	-0.35±0.31	0.42±0.64	3.04	b
53	19 53 54.3 +19 58 47	-	-	0.152	0.03±0.15	0.14±0.20	21.06	a
54	20 13 08.8 +19 56 49	-	-	0.015	1.00±0.00	0.13±0.39	1.67	c?
55	20 02 01.0 +19 55 41	-	-	0.014	1.00±0.00	-0.20±0.44	1.50	a
56	19 40 27.7 +19 38 48	194028.3+193835	16	0.023	-0.63±0.27	0.16±0.89	1.83	a
57	19 56 58.4 +19 35 06	-	-	0.020	1.00±0.00	-0.45±0.46	2.21	e?
58	19 44 32.7 +19 32 18	194430.4+193255	52	0.117	-0.03±0.19	-0.26±0.27	11.03	b
59	19 40 37.9 +19 29 55	194038.2+192951	16	0.027	-0.26±0.31	-0.85±0.58	2.12	—

Table 1b. continued.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl ([°] ' ")	RXS J Name	RXS Err (")	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
60	20 19 18.7 +19 23 52	-	-	0.016	0.38±0.51	0.31±0.46	1.73	–
61	20 11 12.4 +19 21 43	-	-	0.032	-0.49±0.28	-0.79±0.72	2.52	a
62	20 17 43.5 +19 14 23	-	-	0.310	0.13±0.09	-0.06±0.12	25.46	a?
63	20 18 15.9 +19 04 08	-	-	0.018	0.67±0.37	0.37±0.35	2.00	b
64	19 53 00.6 +18 51 16	-	-	0.016	1.00±0.00	0.86±0.48	1.77	–
65	20 07 40.2 +18 38 40	-	-	0.045	-0.16±0.30	0.20±0.40	2.91	a
66	20 05 56.3 +18 38 35	-	-	0.029	-0.56±0.25	-0.33±0.55	2.26	a?
67	19 47 22.7 +18 32 32	-	-	0.016	1.00±0.00	-0.61±0.38	1.81	a?
68	20 11 13.2 +18 30 34	-	-	0.046	0.03±0.25	0.59±0.26	5.24	f
69	19 49 29.9 +18 28 47	-	-	0.018	0.16±0.46	0.21±0.59	1.96	–
70	19 55 57.0 +18 28 06	-	-	0.044	0.30±0.32	-0.44±0.32	4.89	–
71	20 05 07.4 +18 25 26	-	-	0.011	0.74±0.58	1.00±0.00	1.26	c
72	20 02 34.4 +18 22 02	-	-	0.024	-0.54±0.30	1.00±0.00	1.93	a
73	20 06 31.3 +18 18 03	200629.7+181813	44	0.031	0.68±0.27	0.60±0.25	3.42	–
74	20 07 20.5 +18 10 26	-	-	0.020	0.21±0.54	0.23±0.58	2.25	a?
75	19 54 55.0 +18 04 29	-	-	0.022	1.00±0.00	-0.10±0.38	2.43	–
76	19 40 06.5 +18 00 43	194006.8+180038	14	0.014	1.00±0.00	-0.06±0.45	1.54	a
77	20 09 22.8 +18 00 42	-	-	0.021	-0.07±0.41	-0.39±0.81	1.62	a?
78	20 15 23.5 +17 58 36	-	-	0.018	0.47±0.38	0.69±0.30	1.97	–
79	20 17 39.2 +17 56 30	-	-	0.020	1.00±0.00	0.20±0.37	2.25	–
80	20 07 36.5 +17 42 09	-	-	0.214	0.11±0.11	0.24±0.14	11.79	a
81	20 04 35.5 +17 40 22	200434.5+174014	16	0.034	0.38±0.35	0.25±0.37	3.79	a
82	19 42 58.5 +17 37 08	-	-	0.016	1.00±0.00	0.41±0.35	1.82	c?
83	20 05 43.1 +17 35 49	200542.5+173543	31	0.047	0.60±0.20	0.46±0.22	10.31	a
84	19 50 42.4 +17 31 33	-	-	0.023	-0.09±0.39	-0.11±0.54	1.84	–
85	20 14 17.4 +17 27 38	201417.7+172744	10	0.044	0.31±0.27	-0.29±0.30	3.27	a
86	19 48 48.6 +17 26 04	-	-	0.021	1.00±0.00	0.84±0.59	2.30	a?
87	19 38 17.8 +17 18 29	193816.5+171850	21	0.216	1.00±0.00	0.34±0.10	27.78	SNR
88	19 38 39.9 +17 18 03			0.094	0.98±0.09	-0.02±0.14		SNR
89	19 38 54.1 +17 15 00			0.067	1.00±0.00	0.39±0.16		SNR
90	19 38 31.5 +17 14 32			0.146	1.00±0.00	0.18±0.12		SNR
91	19 39 01.7 +17 08 17			0.024	1.00±0.00	-0.04±0.31		SNR
92	20 05 25.1 +17 10 14	200524.9+171021	15	0.017	0.52±0.35	0.54±0.34	1.88	e
93	19 38 35.7 +17 04 58			0.035	1.00±0.00	0.89±0.21		SNR
94	19 38 57.9 +17 04 57			0.054	1.00±0.00	-0.14±0.24		SNR
95	20 03 31.0 +17 07 47	200330.8+170753	23	0.025	-0.10±0.40	0.03±0.62	1.94	–
96	20 09 29.4 +17 06 02	200929.4+170612	24	0.016	1.00±0.00	0.53±0.38	1.74	–
97	20 04 08.1 +17 04 19	200407.7+170432	14	0.067	-1.00±0.00	-1.00±0.00	4.91	a
98	20 03 18.0 +17 02 41	200317.9+170250	12	0.033	0.83±0.18	0.38±0.30	3.63	a
99	20 15 20.9 +16 58 07	-	-	0.008	1.00±0.00	0.18±0.77	0.84	–
100	19 46 08.8 +16 51 03	194608.2+165108	35	0.067	-0.21±0.22	0.07±0.35	4.96	a?
101	20 14 53.1 +16 46 32	201453.1+164635	10	0.046	0.78±0.19	0.02±0.26	5.07	a
102	19 41 57.5 +16 44 41	-	-	0.028	0.91±0.18	0.03±0.29	3.11	a
103	20 10 40.7 +16 42 17	201040.5+164223	21	0.017	1.00±0.00	-0.08±0.41	1.93	a
104	19 59 57.5 +16 43 14	195957.3+164329	17	0.039	0.21±0.34	-0.44±0.36	4.30	a?
105	20 00 59.9 +16 36 24	-	-	0.016	1.00±0.00	-0.49±0.55	1.76	a
106	20 18 38.1 +16 27 54	201838.4+162801	14	0.017	0.15±0.45	-0.14±0.53	1.90	a?
107	20 13 54.2 +16 25 23	201354.3+162526	8	0.045	0.83±0.19	0.15±0.23	4.18	a
108	20 04 42.2 +16 26 57	200441.5+162704	16	0.015	0.82±0.37	0.55±0.41	1.69	a
109	20 15 02.2 +16 22 08	201503.1+162211	18	0.015	0.79±0.29	-0.66±0.31	1.61	b
110	19 56 40.6 +16 21 48	-	-	0.019	1.00±0.00	1.00±0.00	2.11	–
111	19 56 06.9 +16 19 58	195606.1+162007	14	0.024	-0.14±0.37	0.56±0.51	1.86	b?
112	20 07 59.3 +16 09 53	200759.3+160955	8	0.704	0.55±0.05	0.22±0.06	74.58	a
113	20 10 13.3 +16 08 57	201013.4+160900	16	0.014	1.00±0.00	-0.76±0.30	1.61	a?
114	20 17 25.5 +16 03 20	201725.6+160325	10	0.081	1.00±0.00	0.66±0.13	28.94	f?
115	19 54 15.8 +15 59 51	-	-	0.022	-0.20±0.37	1.00±0.00	1.75	a
116	20 09 51.9 +15 57 35	200951.8+155743	15	0.019	0.70±0.32	0.52±0.34	2.11	a
117	20 18 45.9 +15 50 54	201846.1+155058	13	0.019	-0.46±0.30	0.26±0.60	1.53	a
118	20 04 48.1 +15 54 05	200448.1+155413	21	0.088	-0.42±0.17	-0.01±0.37	1.75	a
119	19 46 12.2 +15 49 02	194613.1+154856	13	0.026	0.59±0.33	0.41±0.41	2.90	a

Table 1b. continued.

No	Coordinates (2000.0) RA (^h ^m ^s) Decl (° ' ")	RXS J Name	RXS Err (")	CR (c/s)	HR1	HR2	F _X /10 ⁻¹³ erg/cm ² /s	ID
120	20 13 57.5 +15 44 33	201358.0+154436	30	0.025	1.00±0.00	0.61±0.27	2.74	a
121	19 58 37.1 +15 43 40	195836.9+154353	10	0.129	0.87±0.10	0.26±0.15	24.29	a
122	20 08 26.6 +15 40 29	200826.7+154033	8	0.176	-0.17±0.12	-0.46±0.19	12.63	a
123	19 43 11.8 +15 28 42	194312.2+152834	12	0.010	1.00±0.00	0.62±0.67	1.14	–
124	19 42 50.0 +15 22 33	194250.6+152229	18	0.018	-0.29±0.36	0.18±0.52	1.40	a
125	19 54 01.9 +15 17 37	195402.4+151719	28	0.063	0.04±0.21	0.44±0.25	3.95	a
126	20 18 54.4 +15 07 14	201855.0+150717	14	0.050	0.92±0.12	0.72±0.16	28.34	b?
127	20 17 05.3 +15 02 36	201705.5+150236	21	0.023	0.22±0.44	0.64±0.42	2.51	b
128	19 43 02.2 +14 55 58	194302.3+145550	20	0.019	0.58±0.45	0.46±0.44	2.08	c
129	20 14 55.6 +14 50 55	201455.3+145054	14	0.016	0.77±0.45	0.36±0.35	1.83	a
130	20 19 24.9 +14 36 58	-	-	0.026	0.16±0.20	-0.05±0.27	2.90	b?
131	19 52 54.9 +14 36 21	-	-	0.041	0.38±0.27	0.09±0.37	4.54	a
132	20 10 13.2 +14 32 14	201013.0+143152	16	0.020	1.00±0.00	0.90±0.37	2.20	b

3. Optical identifications of the ROSAT sources

3.1. Optical variability

As mentioned above, we will use the detection of optical variability and the course of the brightness changes as an additional essential criterion for the identification of the ROSAT sources. Given the number of “only” about 100 000 known optically variable objects on the sky, i.e. 2.5 per square degree, a crude likelihood of finding an optically variable object within a 30'' ROSAT position is 2×10^{-4} . A more thorough estimate which takes into account the brightness and amplitude of variables, has been made by Richter (1968), according to whom the percentage of variables with amplitude >0.3 mag among all stars up to 17^m is about 0.23%. The total number of stars ($<17^m$) in the Sge area is estimated to be 7.25×10^5 . This implies a likelihood of finding any optically variable object within a 30'' ROSAT position of about 3.8×10^{-2} for the Sge field. Concerning the other field (26 Com), we deduce from Richter & Greiner (1999) that it contains about 1/20 of the number of stars existing in the Sge field. If we assume that the percentage of variables is also 0.23% (which may not be the case due to the different galactic latitude), we arrive at a 2×10^{-3} chance coincidence for a variable object being within a 30'' ROSAT position. We note in passing that in a more recent paper Vogt et al. (2004) estimated the percentage of variable stars among all stars to be as large as 7.9% – a factor 34 more than the above mentioned value. This estimate is based on two selection effects: (i) the sample of Richter includes stars between 12 and 17 mag, while that of Vogt contains stars brighter than about 11.5 mag. The brighter the stars, the larger the percentage of the (more frequent variable) giant stars. (ii) the sample by Richter is limited to brightness amplitudes >0.3 mag, while that of Vogt includes everything down to ≥ 0.1 mag. It is known that the number of variables increases steeply with smaller amplitude, though a quantitative estimate covering different types of variables is still missing.

Four additional criteria were applied in order to support the optical identification of the X-ray sources:

1. Using the objective prism spectra taken with the Hamburg Schmidt telescope on Calar Alto, which mainly selected stars and AGN. In the Hamburg objective prism survey (Hagen et al. 1995, Bade et al. 1998) spectra are taken in the 3400–5400 Å range with a dispersion of 1390 Å/mm down to 17–18th mag covering the whole northern hemisphere except the galactic plane ($|b| > 20^\circ$). A systematic identification of X-ray sources from the Bright Source Catalog of the ROSAT all-sky survey has been published by Zickgraf et al. (2003).
2. Including the positional correlation with the X-ray positions, i.e. giving lower weight to more distant sources.
3. For the brighter sources and multiple optical counterpart candidates: Obeying consistency between X-ray absorption as determined from the X-ray spectra and visual extinction.
4. Evaluating the X-ray to optical intensity ratio for known populations. It is long known (Maccacaro et al. 1982, Stocke et al. 1991) that stars populate only some sub-phase-space in the $\log(f_X/f_{\text{opt}})$ vs. $B - V$ diagram, while accretion-powered systems can reach much larger f_X/f_{opt} values. We have used these properties to select the more likely optical counterpart (see Fig. 2 which shows, for the Com field, the stars (filled triangles) according to Stocke et al. (1991) together with the upper boundary of values reachable by stars (straight line)). We used the X-ray fluxes in the 0.1–2.4 keV band as determined either from spectral fitting (bright sources) or count-to-flux conversion (for faint sources; more specif-




Fig. 3. The ratio of absorption-corrected X-ray (0.1–2.4 keV) to extinction-corrected optical (V band) luminosity as a function of visual extinction, with the insert showing the range up to $A_V = 12$ mag. The ordinate corresponds to the correction which has to be applied to the X-ray to optical luminosity ratio if the observed values without extinction correction are used. Shown are two blackbody (bb) models with 50 and 150 eV, two thermal bremsstrahlung (tb) models with 100 eV and 900 eV, and two power law (po) models with photon index 1.5 and 2.0, respectively. Except for very soft X-ray spectra (50 eV blackbody and 100 eV thermal bremsstrahlung) this ratio is within a factor of two with respect to the uncorrected luminosity values. This justifies the decision to use observed, not extinction-corrected values to determine the X-ray to optical luminosity ratio (see text).

ically: for sources with $HR1 \geq 0$ we use $f_X = 0.9 \times 10^{11}$ cts $\text{cm}^2 \text{erg}^{-1}$ matching a power law spectrum of photon index 2, while for sources with $HR1 < 0$ we use $f_X = 1.27 \times 10^{11}$ cts $\text{cm}^2 \text{erg}^{-1}$ matching a blackbody spectrum with temperature of a few hundred eV and the V magnitudes (with $f_{\text{opt}} = 4.26 \times 10^{-6-0.4 \times V}$) from USNO-A2 (if not available, then B). We did not apply extinction correction, neither to the X-ray fluxes nor to the visual magnitudes, as these corrections are uncertain and the net effect is within a factor of 2 for most sources (Fig. 3). This allows to select the most probable stellar counterpart, but not to discriminate between e.g. star and quasar (since then a large f_X/f_{opt} range is allowed).

An unequivocal optical identification of the ROSAT sources was not possible in some cases because the accuracy of the ROSAT X-ray positions is not better than 30'', and in most cases there are several possible optical counterpart candidates within the error circle and therefore Swift/XRT observations were obtained. This is particularly true in the low galactic latitude field γ Sge. Moreover, the real counterpart may be invisible in some cases, i.e. fainter than our limiting magnitude. Therefore we can only estimate the reliability of a supposed optical identification by statistical methods (see, e.g., Richter & Greiner 1999). But anticipating the results, we can say that the probability of a positive optical identification is very high if we are dealing either

with a bright object (as a rule an object with an HD or a NGC number), or an active galactic nucleus (quasar, BL Lac object, Seyfert galaxy). As a new element for optical identification we also consider an object with brightness variability as a very likely optical counterpart since the chance probability of a variable object to be inside the error circle of the ROSAT position is rather small (see above).

Thus, we have tested the objects in or near the error circles of the ROSAT positions for variability with the following exceptions: 1. Very bright objects, mostly HD stars, which are too bright to be tested on astrographic plates. Empirically, most of them will be BY Dra stars with amplitudes rarely more than 0.1 mag and therefore their variability is hard to discover on photographic plates. 2. Objects fainter than about 17.5 mag near the plate centre and about 16.5 mag near the edge of a plate. 3. galaxies (with exception of some AGN).

All magnitudes, if not noticed otherwise, are photographic and not far from the B magnitude in the system of Johnson and Morgan (typically $B = m_{\text{phot}} + 0.1$ mag).

The results are given in Tables 5a,5b and 8, and Figures 40a,40b contain the finding charts of all ROSAT sources, with each chart having a size of 2.5×2.5 . The labelling is the same as in column 2 of Table 5a,5b.

Tables 5a,5b contain the data of optical objects inside or near the error circles of the ROSAT positions. As a rule, all objects brighter than about 18 mag are listed. In some cases, if of interest, also fainter objects are added. The columns contain the following informations:

Column 1: Running number of the ROSAT objects from Table 1a,1b, appended with an alphabetical specification of all optical objects inside or near the error circle of about $30''$ of the ROSAT position. These letters are also used as labels in Fig. 40a,40b.

Column 2: Coordinates (2000.0) of the optical objects in right ascension ($^{\text{h}},^{\text{m}},^{\text{s}}$) and declination ($^{\circ},',''$). A dash means that no object visible on our plates (with some exceptions) is within the error circle of the ROSAT source.

Column 3: Distance D between ROSAT position and optical position in arcseconds, unless the optical object is extended.

Column 4: Number in usual star or nebular object catalogues.

Column 5: If the object is variable: Name from the General Catalogue of Variable Stars and its supplements (GCVS, Moscow). If not named, the preliminary designation of newly discovered Sonneberg variables is given by the usual S-number (some prominent cases among the of order 70 new variables discovered in this work have already been published by us separately, and already received an IAU variable star name). In few cases, the number in the New Catalogue of Suspected Variable Stars (NSV catalogue, Moscow 1982) is given. If the object was found to be constant on all plates, a "C" is given while a "C?" means that there could be small amplitude variations which are marginal in our data. A void place means that the object was not tested for variability, for example bright stars (mostly HD stars), and very faint stars below the plate limit.

Column 6: Type of object and type of variability corresponding to the nomenclature of the GCVS. G = galaxy. GCl = cluster of galaxies. AGN = active galactic nucleus. AGN? = supposed AGN only by reason of its blue colour, though in single cases it may be a white dwarf or a cataclysmic variable. If possible, further sub-classification of AGN: QSO = quasistellar object, BLL = BL Lacertae object, SY = Seyfert galaxy. ULX = ultra-luminous X-ray source. CV = cataclysmic variable. Further specification: UGSU = SU UMa

type, AM = AM Her type, NC = very slow nova. E = eclipsing variable, EA = Algol type, EB = Beta Lyrae type. CA = chromospherically active star. If possible, further subclasses are specified: RS = RS CVn, BY = BY Dra. UV = UV Cet type. LB resp. SRB = slowly irregular resp. semiregular variable of late spectral type. PN = planetary nebula.

Column 7: Spectral type. FG means F or G star according to the objective prism spectra taken with the Hamburg Schmidt telescope on Calar Alto (Bade et al. 1998).

Columns 8-10: RBV magnitudes; if existing, mostly from the USNO-A2 catalogue.

Column 11: Brightness amplitude, generally in the blue.

Column 12: Logarithm of the ratio of X-ray to visual flux. This ratio is arbitrary for variable sources, since the catalogued V (or B) band value is taken which is not contemporaneous to the X-ray observation.

Dates or light curves of already known variables can be found in the literature (see notes on individual objects). For newly discovered variable objects we give light curves either for the whole time of observation (Figs. 41a,41b), or for interesting time intervals (Figs. 6 - 37; see above section).

The *Simbad* and *NED* catalogues were used to search for coincidences, and matches are listed either in Table 5a, 5b, or mentioned in sections 3.4 and 3.5.

3.2. Pointed X-ray observations

In addition to the ROSAT all-sky survey and optical data, we have used the ROSAT pointed observations with both the position-sensitive proportional counter (PSPC) and the high-resolution imager (HRI), the *Chandra* and *XMM-Newton* archives, and dedicated Swift/XRT observations. This provided more accurate source positions which were particularly helpful in the Sge field due to the heavy crowding in the galactic plane. It also provided a second epoch X-ray observation which allows us to assess X-ray variability for a number of sources. The results of the dedicated Swift/XRT observations are summarized in Tables 3a and 3b. Improved positions and variability are always mentioned in the notes for the individual objects, sections 3.4 and 3.5.

3.3. SDSS spectra

All 313 objects in Table 5a have been cross-correlated with the DR9 spectral data release, and 139 matches were found. Interestingly, while it provided optical identifications for hitherto 21 unclassified objects, this affected only 2 of our optical identifications of X-ray sources (see below). The more important impact, however, was the provision of redshifts for all these 139 objects which are listed in Table 4.

The two objects where SDSS-spectra helped in the identification were: (i) *Com097*: without spectra, no distinction was possible between *Com097a* and *Com097b*, while SDSS-III provided a spectrum for both, thus solving the ambiguity. (ii) *Com109*: similarly, the distinction between the 5 optical objects was resolved by the SDSS-III spectrum of *Com109a*.

Table 2a. Spectral fit parameters for Com sources

No	N_{H} (10^{21} cm^{-2})	Power law model		χ_{red}	Flux ph/cm ² /s
		Norm (ph/cm ² /s/keV)	photon index		
003	1.02E-01±2.65E-01	1.43E-04±2.05E-04	-3.00E+00±1.89E+00	1.22	6.00E-04
007	1.00E-03±4.57E-01	4.87E-05±1.28E-04	-2.26E+00±3.75E+00	0.04	-2.91E-04
008	9.89E-02±8.95E-01	7.42E-05±2.01E-04	-2.60E+00±4.78E+00	2.88	1.87E-05
010	1.62E-01±1.19E+00	4.76E-05±1.79E-04	-3.24E+00±5.75E+00	2.20	8.78E-05
022	1.15E-01±2.41E-01	6.74E-04±2.31E-04	-1.91E+00±1.04E+00	0.49	1.55E-03
027	1.00E-03±2.08E-01	3.51E-04±1.72E-04	-1.37E+00±1.23E+00	0.64	8.15E-04
030	2.86E-01±4.85E-01	3.44E-04±1.98E-04	-3.06E+00±1.40E+00	0.65	9.67E-04
038	5.29E-02±4.19E-01	1.00E-05±1.16E-04	-4.02E+00±8.21E+00	0.41	2.06E-04
040	3.31E-01±7.16E-01	3.21E-04±2.10E-04	-2.91E+00±1.89E+00	0.72	7.49E-04
042	1.00E-03±3.97E-01	7.92E-05±1.29E-04	-2.00E+00±2.98E+00	2.23	3.86E-04
044	6.83E-01±2.68E+00	3.35E-05±1.52E-04	-5.74E+00±6.11E+00	2.39	2.83E-04
045	8.39E-01±1.64E+00	1.04E-05±6.80E-05	-8.00E+00±1.88E+01	0.87	7.37E-04
050	1.00E-03±6.17E-01	2.43E-04±9.44E-04	-6.41E-01±1.42E+00	1.78	1.96E-04
052	2.14E-01±1.32E-01	2.08E-03±3.50E-04	-2.41E+00±4.88E-01	1.24	4.72E-03
056	2.31E-01±7.93E-01	2.81E-04±2.37E-04	-2.23E+00±2.82E+00	0.28	5.39E-04
057	9.54E-02±1.50E-01	7.20E-04±2.26E-04	-2.21E+00±7.49E-01	1.18	1.74E-03
063	7.50E-02±4.53E-01	3.98E-04±5.07E-04	-1.23E+00±1.02E+00	0.95	9.36E-04
068	1.00E-03±4.73E-01	6.89E-05±1.19E-04	-1.92E+00±3.39E+00	1.68	3.37E-04
075	1.20E+00±3.16E+00	2.81E-05±1.71E-04	-7.73E+00±2.27E+01	1.37	6.44E-04
103	1.05E-02±4.22E-01	2.03E-04±2.12E-04	-1.32E+00±2.23E+00	0.70	4.13E-04
105	1.24E-01±3.78E-01	1.78E-04±1.65E-04	-2.75E+00±1.92E+00	1.13	7.67E-04
109	3.18E-01±1.42E+00	1.24E-04±1.99E-04	-3.10E+00±4.06E+00	0.70	2.70E-04
120	2.33E-01±1.13E+00	1.88E-04±1.54E-04	-2.19E+00±4.15E+00	0.29	3.85E-04
124	7.39E-01±1.46E+00	4.17E-04±2.27E-04	-3.08E+00±3.05E+00	1.23	5.69E-04
127	7.11E-01±1.29E+00	1.99E-04±1.80E-04	-4.28E+00±2.39E+00	1.14	9.20E-04
134	4.39E-01±5.29E-01	7.06E-04±2.71E-04	-2.66E+00±1.53E+00	0.75	1.38E-03
139	2.09E-01±5.82E-01	3.46E-04±2.10E-04	-2.19E+00±2.14E+00	0.50	7.01E-04
144	2.67E-01±3.15E-01	3.69E-04±1.82E-04	-3.33E+00±9.95E-01	0.89	1.39E-03
153	1.85E-01±3.05E-01	3.64E-04±2.01E-04	-2.79E+00±1.23E+00	0.43	1.07E-03
155	1.00E-03±1.87E-01	1.16E-04±1.53E-04	-2.15E+00±1.57E+00	1.87	5.84E-04
157	2.92E+01±6.97E+04	2.93E-03±7.66E+00	-1.00E+00±1.78E+03	1.28	7.11E-04
159	5.83E-01±4.38E-01	6.19E-04±3.30E-04	-4.46E+00±9.71E-01	1.51	3.12E-03
168	1.00E-03±2.98E-01	3.79E-04±4.75E-04	-1.08E+00±1.22E+00	1.10	1.03E-03
169	1.58E-01±4.77E-01	2.48E-04±1.66E-04	-2.54E+00±2.07E+00	1.23	8.00E-04
171	9.75E+00±7.76E+01	3.34E-03±8.08E-03	-8.00E+00±1.69E+02	0.93	1.21E-04
174	6.62E-01±1.39E+00	1.01E-04±2.15E-04	-5.38E+00±3.02E+00	1.15	8.66E-04
183	1.35E-01±9.01E-01	4.67E-05±1.78E-04	-3.33E+00±5.12E+00	0.49	1.88E-04
187	1.00E-03±4.40E-01	1.00E-05±1.74E-04	-3.46E+00±7.40E+00	1.05	2.59E-05
188	1.32E-01±1.87E-01	1.37E-03±3.63E-04	-1.78E+00±7.60E-01	0.84	2.36E-03
197	1.72E-01±2.96E-01	6.18E-04±1.29E-03	-1.71E+00±1.48E+00	0.95	1.43E-03
199	1.00E-03±3.80E-01	1.49E-04±2.02E-04	-1.64E+00±2.60E+00	0.54	4.05E-04
204	8.84E+00±3.66E+01	3.98E-03±1.01E-02	-8.00E+00±7.66E+01	0.28	4.16E-04
215	2.65E-01±1.94E-01	8.78E-04±1.93E-03	-1.57E+00±1.50E+00	1.06	1.08E-03
223	6.98E-01±5.25E-01	6.27E-04±2.11E-03	-2.14E+00±2.57E+00	0.7	7.79E-04
229	1.99E-02±1.63E-01	2.55E-04±2.03E-04	-2.13E+00±1.15E+00	0.74	8.43E-04
230	2.16E-01±5.09E-01	8.18E-04±5.68E-04	-1.60E+00±1.27E+00	0.34	1.64E-03
236	1.11E+00±1.83E+01	3.33E-04±1.93E-03	-1.87E+00±8.96E+00	0.81	3.60E-04

No	N_{H} (10^{21} cm^{-2})	Bremsstrahlung model		χ_{red}	Flux ph/cm ² /s
		Norm (ph/cm ² /s/keV)	kT (keV)		
006	1.48E-01±9.19E-02	1.81E-03±8.19E-04	1.03E+00±3.86E-01	1.85	3.54E-03
013	2.17E-01±1.13E-01	1.46E-03±3.77E-04	5.53E-01±1.83E-01	2.51	3.04E-03
036	4.19E-02±4.83E-01	3.94E-07±1.87E-05	1.00E-01±6.02E-01	1.1	3.21E-04
039	1.00E-03±4.52E+00	3.25E-08±1.90E-05	9.54E-02±1.43E+00	1.14	1.05E-04
069	1.09E-02±1.93E-01	1.94E-04±2.48E-04	8.65E-01±1.31E+00	0.97	4.79E-04
087	2.91E+01±9.18E+02	4.83E-02±4.70E+00	3.02E-02±4.66E+00	0.56	1.70E-04
094	2.13E-01±3.15E-01	3.15E-04±3.04E-04	1.02E+00±3.34E+00	0.6	5.36E-04
104	1.32E-01±1.42E-01	7.87E-04±4.66E-04	6.59E-01±3.84E-01	0.57	1.70E-03
112	3.47E-02±1.17E-01	4.80E-04±3.77E-04	9.05E-01±7.24E-01	1.77	1.25E-03
117	1.48E-01±3.07E-01	2.30E-04±2.15E-04	3.96E-01±4.72E-01	1.61	6.67E-04
141	7.17E+01±4.16E+02	3.24E+03±1.15E+05	4.84E-02±3.87E+00	1.98	3.90E-04
181	1.00E-03±1.31E+00	5.47E-08±1.13E-05	9.29E-02±4.91E-01	1.75	1.83E-04
196	1.52E-01±4.08E-01	4.01E-04±1.10E-03	5.57E+00±5.39E+01	0.24	7.67E-04
224	5.51E-02±1.51E-01	5.30E-04±5.82E-04	9.75E-01±9.08E-01	1.18	1.17E-03

Table 2a. cont.: Spectral fit parameters for Com sources

No	Blackbody model				
	N_{H} (10^{21} cm^{-2})	Norm (ph/cm ² /s)	kT (keV)	χ_{red}	Flux ph/cm ² /s
004	5.72E+00±1.81E+01	1.66E-01±2.37E+00	9.99E-02±4.70E-01	0.88	7.00E-04
043	1.00E-03±1.60E-01	2.81E-03±2.42E-03	1.53E-01±2.56E-02	0.75	1.30E-03
062	1.00E-03±2.12E-01	2.16E-03±2.56E-03	1.48E-01±3.35E-02	0.79	8.89E-04
067	1.73E-01±3.22E+00	2.63E+00±3.04E+02	1.59E-02±4.95E-01	3.16	2.89E-04
113	1.00E-03±6.06E-02	2.34E+00±8.49E+00	1.78E-02±4.67E-03	2.66	1.75E-02
158	5.55E-02±9.94E-01	1.47E-03±1.04E-02	6.76E-02±4.61E-01	1.35	1.52E-04
160	2.84E+01±1.46E+02	5.26E+03±4.52E+05	6.04E-02±1.99E+00	0.53	3.47E-04
170	1.43E+01±1.28E+02	2.26E+04±2.91E+06	4.19E-02±2.49E+00	0.51	5.68E-04
192	1.00E-03±2.60E-01	1.41E-03±1.27E-03	2.18E-01±7.17E-02	1.19	8.81E-04
195	5.57E-02±1.56E-01	5.70E-03±4.36E-03	1.42E-01±1.78E-02	1.69	2.09E-03
221	8.70E-01±3.59E+00	1.44E-01±2.44E+00	4.99E-02±5.25E-01	0.44	6.85E-04
237	1.66E+00±4.25E+02	1.53E-03±7.74E-01	1.00E-01±9.56E+00	0.43	8.60E-05
238	1.00E-03±1.59E+02	1.96E-03±1.15E+01	2.67E-02±2.10E+01	0.84	8.49E-05

Table 2b. Spectral fit parameters for Sge sources

No	Bremsstrahlung model				
	N_{H} (10^{21} cm^{-2})	Norm (ph/cm ² /s/keV)	kT (keV)	χ_{red}	Flux ph/cm ² /s
003	6.63E-01±1.16E+01	1.95E-06±1.04E-04	1.00E-01±2.04E+00	2.72	2.07E-04
013	1.98E+01±1.55E+02	4.80E-02±3.13E-01	1.00E-01±3.96E+00	0.59	2.08E-04
016	1.00E-03±5.43E-02	6.30E-04±3.94E-04	1.21E+00±7.22E-01	0.98	1.56E-03
020	1.00E-03±1.29E-01	1.79E-04±2.17E-04	6.79E-01±7.71E-01	0.47	5.82E-04
045	1.00E-03±5.91E-01	1.16E-07±1.02E-05	9.84E-02±2.84E-01	2.92	2.39E-04
065	1.00E-03±3.84E+00	2.47E-08±1.38E-05	9.68E-02±1.31E+00	0.57	9.15E-05
068	1.47E-01±2.09E+00	2.79E-07±2.71E-05	9.98E-02±1.24E+00	1.19	9.37E-05
080	1.52E-01±1.89E-01	6.23E-04±3.60E-04	1.46E+00±3.86E+00	1.07	7.48E-04
083	1.07E+00±3.62E+01	3.89E-04±1.63E-03	1.00E+02±1.72E+04	0.12	6.23E-04
100	4.08E-01±1.61E+00	1.24E-04±3.13E-04	2.71E-01±5.16E-01	1.55	2.48E-04
125	1.00E-03±1.90E+00	1.11E-08±3.72E-06	8.26E-02±4.33E-01	2.56	8.44E-05
126	9.74E-01±6.37E+00	1.15E-03±1.37E-03	1.93E+01±4.81E+02	0.5	1.76E-03

No	Blackbody model				
	N_{H} (10^{21} cm^{-2})	Norm (ph/cm ² /s)	kT (keV)	χ_{red}	Flux ph/cm ² /s
006	1.00E-03±1.59E-01	1.88E-03±1.89E-03	1.36E-01±4.07E-02	1.11	7.21E-04
007	1.07E-01±4.84E-01	1.24E-03±2.06E-03	1.77E-01±6.42E-02	0.73	4.69E-04
019	1.00E-03±4.15E-01	3.02E-02±5.67E-01	2.40E-02±4.75E-02	1.05	7.13E-04
021	7.03E-02±8.81E-02	2.07E+00±5.51E+00	2.63E-02±1.43E-02	1.29	1.61E-02
023	1.00E-03±3.14E-01	1.05E-03±1.71E-03	1.77E-01±7.69E-02	1.88	5.12E-04
028	1.00E-03±6.02E-01	5.51E-04±7.91E-04	2.57E-01±1.71E-01	1.85	4.17E-04
035	3.76E+00±2.98E+01	6.04E+02±5.19E+04	2.98E-02±6.67E-01	0.56	1.04E-03
037	4.07E+01±2.95E+02	1.22E+03±1.27E+05	7.51E-02±2.96E+00	1.64	2.51E-04
038	2.47E+01±9.19E+01	5.50E-02±9.02E-01	2.52E-01±2.83E+00	0.45	1.59E-03
049	1.00E-03±5.21E-01	6.66E-04±1.94E-03	1.57E-01±1.05E-01	1.03	2.72E-04
053	1.00E-03±1.87E-01	2.03E-03±1.69E-03	2.00E-01±4.92E-02	2.58	1.12E-03
058	8.92E-02±4.57E-01	2.57E-03±5.84E-03	1.32E-01±4.58E-02	0.66	6.88E-04
062	1.00E-03±1.12E-01	3.19E-03±1.76E-03	1.78E-01±3.08E-02	0.93	1.59E-03
085	1.00E-03±5.11E-01	6.06E-04±1.66E-03	1.65E-01±9.48E-02	0.42	2.28E-04
097	1.00E-03±1.48E+00	2.67E-02±2.35E+00	1.89E-02±1.35E-01	1.44	3.07E-04
101	1.30E+01±5.62E+01	6.71E-01±2.13E+01	9.83E-02±1.13E+00	0.86	2.65E-04
107	3.33E+01±1.91E+02	1.67E+04±1.83E+06	5.84E-02±2.38E+00	0.55	2.26E-04
112	1.00E-03±7.96E-02	5.71E-03±1.46E-03	2.68E-01±3.31E-02	0.93	4.28E-03
114	3.68E-01±3.42E+00	1.91E-03±1.47E-03	5.50E-01±4.80E-01	0.78	1.75E-03
118	1.73E-01±6.65E+00	1.09E-02±1.34E+00	3.53E-02±1.74E+00	1.78	1.09E-04
121	2.30E-01±9.66E-01	2.01E-03±1.64E-03	2.69E-01±1.09E-01	0.29	1.35E-03
122	1.77E-03±1.83E-01	1.92E-03±1.97E-03	1.42E-01±6.65E-02	0.49	7.88E-04




Fig. 4a. ROSAT survey spectra of Com sources. For each source, the photon spectrum is shown. The choice of the model for a given source was made based on the lowest reduced χ^2 and consistency with the optical properties (see Table 2a for model and spectral parameters).




Fig. 4a. contd. ROSAT survey spectra of Com sources.




Fig. 4a. contd. ROSAT survey spectra of Com sources.




Fig. 4a. contd. ROSAT survey spectra of Com sources.




Fig. 4b. ROSAT survey spectra of Sge sources. For each source, the photon spectrum is shown. The choice of the model for a given source was made based on the lowest reduced χ^2 and consistency with the optical properties (see Table 2b for model and spectral parameters).




Fig. 4b. contd. ROSAT survey spectra of Sge sources.

Table 3a. Swift/XRT observations of ROSAT sources in the Com field

Name	Date of observation	Exposure (sec)	XRT coordinate (2000.0)		Position error (″)	CR (cts/s)
			RA	Decl.		
Com11	2007-07-17	3230	–		–	<0.002
Com12	2008-12-01	1653	12 53 28.00	+25 34 08.9	6.1	0.015±0.003
Com15	2008-02-22	747	12 59 54.90	+25 28 09.0	7.5	0.013±0.005
Com15	2008-02-23	558	12 59 55.25	+25 28 09.2	13.5	<0.006±0.005
Com25	2008-04-20	1487	12 46 38.18	+25 13 24.6	5.8	0.010±0.003
Com28	2007-07-19	1583	–		–	<0.006
Com28	2008-05-11	1498	–		–	<0.006
Com31	2008-08-15	2674	12 56 04.38	+24 51 39.5	7.7	0.003±0.001
Com33	2009-02-15	2297	12 45 14.9	+24 53 25	– ^a	0.024±0.002
Com37	2009-11-29	3499	12 43 09.45	+24 47 03.7	5.4	0.008±0.002
Com46	2009-02-15	4666	12 53 08.27	+24 27 48.0	5.6	0.006±0.001
Com47	2009-02-26	1489	12 22 14.52	+24 24 20.3	7.5	0.023±0.003
Com49	2007-07-01	1068	12 24 31.05	+24 21 05.8	10	0.003±0.003
Com50	2009-02-26	1489	12 22 17.42	+24 19 05.0	– ^a	0.035±0.004
Com60	2008-02-18	3626	–		–	<0.002
Com61	2007-06-14	1726	12 30 54.37	+23 52 01.2	6.2	0.011±0.003
Com65	2009-03-07	1221	–		–	<0.006
Com66	2007-06-14	2079	–		–	<0.003
Com74	2007-12-27	986	13 00 32.75	+23 20 25.4	7.7	0.008±0.003
Com84	2008-02-18	2662	–		–	<0.003
Com89	2009-11-29	3140	–		–	<0.005
Com91	2010-02-15	2835	–		–	<0.003
Com95	2007-07-13	4046	12 31 55.85	+22 48 03.2	5.1	0.011±0.002
Com101	2007-07-10	1862	12 52 32.69	+22 33 33.9	5.3	0.017±0.003
Com102	2007-07-10	3877	12 26 31.77	+22 33 18.1	4.4	0.021±0.002
Com103	2007-07-10	1862	12 52 10.53	+22 33 42.9	5.1	0.015±0.003
Com104	2007-07-10	1862	12 51 47.30	+22 32 38.73	4.1	0.072±0.007
Com106	2007-07-13	597	–		–	<0.01
Com120	2007-06-13	1956	–		–	<0.003
Com126	2007-12-04	1805	12 29 35.48	+21 37 20.6	7.0	0.004±0.002
Com139	2007-12-04	1025	12 19 05.18	+21 06 37.0	5.6	0.033±0.006
Com146	2009-11-29	5182	–		–	<0.002
Com149	2009-11-29	5723	–		–	<0.003
Com151	2012-08-11	540	12 50 47.79	+20 36 21.9	10.2	0.010±0.005
Com155	2007-12-03	1374	12 30 53.50	+20 30 10.3	7.1	0.010±0.003
Com165	2007-07-21	2630	12 26 44	+19 50 40	– ^a	0.018±0.004
Com166	2007-12-02	5199	12 41 51	+19 49 40	– ^a	0.020±0.005
Com167	2007-12-03	2735	12 36 19	+19 48 11	– ^a	0.010±0.004
Com168	2007-12-02	5199	12 42 23.29	+19 47 21.8	3.8	0.040±0.005
Com171	2007-12-02	1425	12 55 27.86	+19 33 26.0	5.2	0.033±0.005
Com180	2007-07-25	1561	–		–	<0.004
Com194	2013-03-11	373	–		–	<0.02
Com195	2007-12-02	917	–		–	<0.007
Com201	2008-08-14	2038	12 54 47.50	+17 56 22.7	6.5	0.009±0.002
Com208	2007-06-14	4603	12 24 05.13	+17 15 22.8	6.8	0.002±0.001
Com212	2007-07-22	4467	12 46 13.51	+17 10 19.9	5.4	0.003±0.001
Com217	2007-11-12	4083	12 54 42.3	+16 56 20	– ^a	0.009±0.003
Com219	2007-11-09	3500	–		–	<0.002
Com220	2008-01-25	3166	12 44 01	+16 53 51	– ^a	0.018±0.004
Com221	2007-12-04	1619	–		–	<0.005
Com225	2007-07-28	2446	–		–	<0.003
Com228	2007-12-05	4357	12 41 52.5	+16 33 22	– ^a	0.042±0.005
Com232	2007-11-12	1454	–		–	<0.006
Com233	2007-11-09	968	12 42 50.98	+16 18 20.5	11	0.003±0.002
Com234	2007-12-04	3255	12 34 38.23	+16 05 44.4	5.4	0.009±0.002
Com236	2007-12-05	1377	12 39 45.04	+16 05 56.5	5.0	0.022±0.004
Com237	2007-12-04	1054	–		–	<0.006

^{a)} diffuse emission.

Table 3b. Swift/XRT observations of ROSAT sources in the Sge field

Name	Date of observation	Exposure (sec)	XRT coordinate (2000.0)		Position error (″)	CR (cts/s)
			RA	Decl.		
Sge5	2007-07-10	3654	19 53 14.97	+23 50 43.7	6.0	0.004±0.001
Sge8	2007-11-24	8318	19 57 26.30	+23 28 46.9	4.1	0.018±0.002
Sge9	2008-02-17	3658	—	—	—	<0.002
Sge10	2008-02-17	6489	19 43 14.15	+23 20 15.9	5.3	0.003±0.001
Sge11	2007-11-28	3514	20 13 51.56	+23 17 51.8	6.3	0.004±0.001
Sge12	2007-08-07	666	20 15 34.24	+23 12 47.8	8.0	0.016±0.006
Sge12	2008-01-08	5874	20 15 34.44	+23 12 57.8	4.2	0.021±0.002
Sge14	2007-07-25	2187	20 18 05.27	+23 03 48.5	6.0	0.009±0.002
Sge14	2007-10-17	1622	20 18 30.81	+22 56 21.2	9.1	0.004±0.002
Sge15	2007-11-29	2163	—	—	—	<0.003
Sge15	2008-02-29	1675	—	—	—	<0.004
Sge17	2007-11-29	3363	20 03 24.91	+22 59 53.9	5.3	0.010±0.002
Sge17	2008-06-17	3123	20 03 24.77	+22 59 48.9	6.1	0.007±0.002
Sge25	2008-02-28 ^a	3865	—	—	—	<0.002
Sge26	2008-03-01	3533	20 10 48.93	+22 21 37.4	4.4	0.026±0.003
Sge29	2007-12-24	8271	20 16 17.90	+22 06 11.2	4.4	0.010±0.001
Sge30	2007-07-24	3745	19 49 58.55	+22 04 34.6	5.3	0.010±0.002
Sge30	2008-05-15	6717	19 49 58.52	+22 04 34.5	4.5	0.012±0.002
Sge32	2007-12-28	4500	—	—	—	<0.001
Sge34	2008-06-17	2728	20 09 23.53	+21 42 57.0	6.5	0.006±0.002
Sge36	2007-07-24	5355	—	—	—	<0.001
Sge39	2008-02-29	4695	20 07 53.41	+21 15 19.1	5.3	0.007±0.001
Sge46	2008-02-28	4191	—	—	—	<0.002
Sge48	2007-07-25	2736	19 53 10.45	+20 43 53.6	7.5	0.003±0.001
Sge48	2007-10-19	2462	19 53 10.23	+20 43 50.9	6.4	0.006±0.002
Sge48	2007-10-21	2456	19 53 10.36	+20 43 51.2	5.1	0.015±0.003
Sge50	2007-12-19	2946	19 41 08.43	+20 08 45.0	5.0	0.019±0.003
Sge51	2007-10-17	2775	—	—	—	<0.002
Sge52	2009-02-23	2677	19 57 01.70	+20 05 54.3	5.7	0.010±0.002
Sge53	2008-02-23	1379	19 53 54.78	+19 59 04.9	5.4	0.023±0.004
Sge54	2007-12-27	4522	20 13 05.46	+19 56 46.1	4.7	0.004±0.001
Sge56	2007-12-18	2291	19 40 28.14	+19 38 25.0	5.8	0.010±0.003
Sge57	2007-12-19	5116	19 56 55.39	+19 35 49.8	4.5	0.016±0.002
Sge58	2007-12-18	993	19 44 32.95	+19 31 44.7	5.6	0.034±0.007
Sge59	2007-12-18	2291	—	—	—	<0.003
Sge60	2007-11-23	3205	20 19 19.16	+19 24 53.1	4.4	0.026±0.003
Sge60	2008-04-23	1830	20 19 18.93	+19 24 52.0	4.7	0.034±0.005
Sge63	2007-11-22	3466	20 18 14.55	+19 04 54.4	6.1	0.006±0.002
Sge64	2007-07-18	4054	19 53 03.30	+18 51 58.0	4.3	0.025±0.003
Sge65	2008-12-09	1384	20 07 41.11	+18 39 17.2	5.9	0.020±0.004
Sge65	2009-03-02	692	20 07 40.95	+18 39 25.2	7.7	0.017±0.006
Sge68	2008-03-03	2081	20 11 11.23	+18 31 03.1	5.9	0.012±0.003
Sge69	2007-07-18	4075	—	—	—	<0.001
Sge70	2007-12-27	2292	19 55 56.98	+18 28 57.2	5.2	0.018±0.003
Sge71	2007-12-29	5536	20 05 08.25	+18 25 33.6	4.6	0.013±0.002
Sge72	2007-11-25	4068	20 02 33.39	+18 22 11.9	6.2	0.005±0.001
Sge73	2008-06-17	2998	—	—	—	<0.002
Sge75	2007-12-16 ^a	2368	—	—	—	<0.003
Sge77	2008-03-02	4183	—	—	—	<0.002
Sge78	2007-12-20	4487	—	—	—	<0.001
Sge78	2007-12-21	3559	—	—	—	<0.002
Sge79	2007-07-23	4292	—	—	—	<0.002
Sge81	2007-12-27	2966	20 04 36.19	+17 40 43.7	5.0	0.014±0.002
Sge82	2007-12-02	4547	19 42 57.08	+17 36 52.3	6.8	0.002±0.001
Sge83	2007-12-03	1439	20 05 44.92	+17 35 51.7	6.2	0.012±0.003
Sge84	2007-07-18 ^a	1613	—	—	—	<0.003
Sge92	2007-12-16	4203	20 05 25.83	+17 10 01.7	8.6	0.002±0.001
Sge95	2007-12-17	2869	—	—	—	<0.001
Sge96	2007-12-07	3799	20 09 30.18	+17 06 28.3	6.0	0.005±0.001
Sge99	2007-12-17	1854	—	—	—	<0.003
Sge109	2007-12-09	4517	20 15 04.12	+16 22 27.7	4.1	0.024±0.003
Sge110	2007-12-14	2945	19 56 41.46	+16 25 58.5	6.4	0.003±0.001
Sge115	2007-11-27	1075	19 54 25.08	+15 57 41.8	8.0	0.009±0.003
Sge117	2007-11-22	4517	20 18 45.07	+15 50 47.6	6.7	0.003±0.002
Sge120	2007-12-08	1366	20 13 58.79	+15 44 32.7	4.8	0.049±0.007
Sge120	2007-12-18	1919	20 13 58.79	+15 44 28.3	4.2	0.078±0.007
Sge124	2007-11-28	3995	19 42 50.13	+15 22 53.6	5.5	0.007±0.002
Sge127	2007-11-22	6215	20 17 03.96	+15 02 41.1	11	0.002±0.001
Sge128	2007-11-25	6977	19 43 05.12	+14 56 13.9	4.1	0.021±0.002
Sge132	2007-11-11	4098	20 10 14.08	+14 31 58.8	4.2	0.029±0.003

^a A second, shorter observation did also not reveal any source.

Table 4. Optical objects in the Com field with existing SDSS-III spectra where the source ID refers to Table 5a.

Source ID	Plate	MJD	Fiber	Ident.	z
003a	2661	54505	550	QSO	0.1957
007a	2660	54504	454	QSO	0.6680
008a	2658	54502	617	GALAXY	0.0668
010a	2658	54502	464	QSO	0.1598
012a	2661	54505	516	QSO	0.1417
014b	2657	54502	543	QSO	0.6789
020a	2660	54504	431	GALAXY	0.1834
024a	2659	54498	464	QSO	0.3408
025b	2660	54504	640	GALAXY	0.0844
026a	2658	54502	111	QSO	0.0637
027a	2659	54498	591	QSO	0.5447
027b	2660	54504	350	STAR	0.0000
029a	2660	54504	183	QSO	0.5865
030a	2658	54502	154	QSO	0.2679
032a	2661	54505	213	GALAXY	0.2465
035a	2660	54504	161	QSO	0.2184
038a	2658	54502	107	QSO	0.5471
040a	2658	54502	164	GALAXY	0.2187
042b	2661	54505	47	QSO	1.2078
044a	2662	54505	392	QSO	0.3710
045a	2660	54504	278	QSO	0.1856
045a	3374	54948	507	QSO	0.1856
047a	2658	54502	213	QSO	0.1167
048a	2660	54504	226	QSO	0.8282
049a	2658	54502	98	QSO	1.0444
051a	2660	54504	58	QSO	0.4647
052b	2662	54505	311	GALAXY	0.1406
054a	2661	54505	319	QSO	0.5071
055a	2658	54502	11	QSO	0.3795
056a	2661	54505	197	QSO	0.1884
059a	2658	54502	242	GALAXY	0.1318
061a	2659	54498	207	QSO	0.3985
062e	2647	54495	450	QSO	0.1382
065a	2647	54495	407	AGN	0.1377
067a	2647	54495	525	STAR	0.0001
068a	2662	54505	293	QSO	0.2583
070a	2660	54504	93	GALAXY	0.1947
070b	2648	54485	406	QSO	1.1775
072b	3374	54948	178	STAR	0.0005
074a	2662	54505	212	QSO	0.2728
075a	2646	54479	531	QSO	0.1606
077a	2648	54485	379	QSO	0.6667
078b	2648	54485	364	QSO	1.3866
079a	2647	54495	618	QSO	0.3412
081a	2648	54485	580	QSO	0.9669
082a	2648	54485	334	QSO	0.7227
083a	2648	54485	367	QSO	0.5299
084a	2647	54495	547	QSO	0.2034
089b	2645	54477	481	GALAXY	0.1405
090a	2645	54477	519	GALAXY	0.0760
095a	2647	54495	426	QSO	0.1603
097a	2647	54495	394	QSO	0.5538
097b	2646	54479	627	STAR	0.0003
098b	2646	54479	432	GALAXY	0.4821
100a	2649	54212	484	QSO	0.0862
101b	2649	54212	421	QSO	0.2113
102b	2646	54479	506	QSO	0.1296
105a	2648	54485	344	QSO	0.1019
107a	2647	54495	160	QSO	0.5230
108a	2646	54479	173	QSO	0.4365

Table 4. continued.

Source ID	Plate	MJD	Fiber	Ident.	z
109a	2646	54479	113	QSO	0.3392
110a	2646	54479	171	QSO	1.0040
115a	2647	54495	318	STAR	0.0000
115c	2646	54479	68	GALAXY	0.2257
116a	2647	54495	101	QSO	0.1798
119a	2646	54479	141	QSO	0.4363
121b	2647	54495	145	STAR	0.0006
122a	2648	54485	290	GALAXY	0.1114
124a	2646	54479	283	QSO	0.3478
126a	2647	54495	288	QSO	0.3484
127a	2647	54495	256	QSO	0.2336
128a	2649	54212	44	QSO	0.0757
130a	2649	54212	249	QSO	0.2354
131a	2648	54485	99	GALAXY	0.1156
132a	2647	54495	204	QSO	0.1897
133a	2647	54495	10	QSO	1.4783
134a	2646	54479	204	QSO	0.4338
138a	2613	54481	453	GALAXY	0.1089
139a	2610	54476	575	GALAXY	0.0514
139b	2611	54477	329	STAR	0.0002
139b	3172	54863	523	STAR	0.0002
140a	2647	54495	128	QSO	0.1391
141a	2613	54481	523	STAR	0.0001
144a	2612	54480	330	QSO	0.3338
146a	2611	54477	369	GALAXY	0.0851
148a	2613	54481	617	QSO	1.3540
149a	2610	54476	515	GALAXY	0.1963
150a	2612	54480	590	QSO	0.5678
151c	2614	54481	623	QSO	0.8401
152a	2611	54477	496	QSO	0.6332
153a	2616	54499	367	QSO	0.0756
155a	2612	54480	504	QSO	0.4287
156a	2615	54483	570	QSO	0.0806
156a	2616	54499	398	GALAXY	0.0801
159a	2613	54481	342	QSO	0.0636
163a	2614	54481	350	QSO	0.2394
164a	2615	54483	342	STAR	0.0001
165a	2611	54477	80	GALAXY	0.2231
167a	2613	54481	476	GALAXY	0.2139
168a	2614	54481	279	QSO	0.1603
168c	2613	54481	623	STAR	0.0000
169a	2610	54476	120	QSO	0.2246
171a	2615	54483	462	QSO	0.1313
172a	2612	54480	170	GALAXY	0.1605
173b	2614	54481	308	QSO	0.5417
174a	2615	54483	266	QSO	0.2790
175a	2614	54481	167	QSO	0.6930
178a	2615	54483	318	QSO	0.4994
178b	2614	54481	21	QSO	0.9963
183a	2614	54481	255	QSO	0.2223
184a	2615	54483	257	QSO	0.5992
185a	2610	54476	43	QSO	0.1970
187a	2614	54481	1	QSO	0.7224
188c	2600	54243	404	GALAXY	0.0711
193c	2597	54231	575	QSO	1.4015
194b	2599	54234	538	QSO	0.2502
195a	2598	54232	577	GALAXY	0.1341
197c	2601	54144	372	GALAXY	0.0667
199a	2599	54234	478	QSO	0.2540
200a	2601	54144	433	QSO	0.0753
202a	2601	54144	516	QSO	0.3526

Table 4. continued.

Source ID	Plate	MJD	Fiber	Ident.	z
203a	2601	54144	344	GALAXY	0.1670
204a	2601	54144	271	QSO	0.2630
206b	2601	54144	306	QSO	0.5478
207a	2598	54232	148	QSO	0.4067
209a	2600	54243	560	QSO	1.2822
210a	2599	54234	119	GALAXY	0.0720
212a	2600	54243	32	GALAXY	0.0804
215a	2599	54234	268	STAR	0.0003
216a	2599	54234	191	GALAXY	0.3806
218a	2599	54234	276	QSO	0.2085
223a	2598	54232	140	QSO	2.4468
225b	2602	54149	50	GALAXY	0.0763
226b	2597	54231	212	STAR	0.0001
227b	2602	54149	134	GALAXY	0.0635
231a	2600	54243	55	QSO	0.5552
235a	1768	53442	620	QSO	0.0711
236b	1769	53502	444	GALAXY	0.0702
236d	2600	54243	207	GALAXY	0.2808

Table 5a. Optical objects in or near the ROSAT X-ray error circle in the Com field.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	$\log(f_X/f_V)$
	R.A	Decl.	"					mag	mag	mag	mag	
001 a	12 33 56.7	+26 02 33	13			AGN?		18.6	18.7	18.6		+0.31
002 a	12 36 02.0	+25 59 20	40	NGC 4565	ULX-4	ULX			10.4			–
003 a	12 54 09.1	+25 55 24	2		S10944 Com	QSO		16.9	16.7	16.8	16.6–18:	+0.15
004 a	12 23 08.5	+25 51 06	14	HD 107793			F8V	8.8	9.6	9.1		-2.89
005 a	12 24 03.5	+25 51 06	15	HD 107935			Am	-	7.0	6.7		-4.84
006 a	12 22 30.3	+25 50 46	23	HD 107700			F2	-	5.4	4.9		-3.91
007 a	12 41 36.0	+25 52 51	8		S10945 Com	QSO		17.2	17.3	17.2	16.9–18.5	-0.24
008 a	12 28 12.8	+25 49 34	5	GSC 1989 2197	S10966 Com	AGN		15.6	16.4	15.9	16.6–18:	-0.97
009 a	12 20 45.6	+25 45 57	14	HD 107399			F8V	8.6	9.7	9.0		-3.82
010 a	12 24 11.2	+25 43 43	16			QSO		17.6	18.7	18.0		-0.06
011 a	12 35 15.2	+25 42 24	27	GSC 1990 2034				14.1	17.1	16.0		-1.02
b	12 35 15.3	+25 42 19	30	GSC 1990 2062				14.1	16.9	15.5		-1.19
012 a	12 53 28.1	+25 34 09	9			QSO		14.7	17.2	15.6		-0.84
013 a	12 25 02.6	+25 33 43	13	HD 108102	IL Com	RS	F8	7.9	8.4	7.9	0.04	-2.77
014 a	12 17 54.7	+25 29 21	22					15.9	17.0	16.3		-0.95
b	12 17 56.8	+25 29 27	26	4C 25.39		QSO		17.5	17.5	17.5		-0.47
015 a	12 59 55.4	+25 28 11	14	GSC 1993 2297	C			11.5	13.7	12.3		-2.26
b	12 59 54.5	+25 27 57	7		V?			18.7	18.9	18.8		+0.34
016 a	12 51 38.9	+25 30 34	12	HD 111813			K1V	8.4	10.0	9.0		-3.55
017 a	12 50 26.6	+25 30 04	11	NGC 4725		SY2		10.1	11.6	10.7		-2.88
018 a	12 39 43.5	+25 30 46	8	GSC 1990 541	C		M4.5	13.2	16.2	14.3		-1.58
019 a	12 54 15.1	+25 24 40	15			G		17.5	19.0	18.1		+0.10
020 a	12 40 10.7	+25 21 31	2			GCI?		16.4	18.6	17.2		-0.63
021 a	12 59 23.6	+25 15 06	25			AGN?		18.2	18.6	18.4		+0.04
022 a	12 30 14.1	+25 18 07	6	GSC 1989 2067		BLL		16.1	16.8	16.4		+0.28
023 a	12 38 43.5	+25 17 33	31		V?		FG	15.3	16.5	15.8		-1.18
024 a	12 30 43.8	+25 14 53	5			QSO		19.1	19.2	19.1		+0.42
025 a	12 46 38.8	+25 13 04	38					13.6	14.6	–		-2.18
b	12 46 38.4	+25 13 18	52			G		18.3	18.6	–		-0.06
026 a	12 26 04.2	+25 06 37	8	1E 1223.5 +2522		AGN		14.4	15.3	14.7		-1.33
027 a	12 36 51.6	+25 07 51	10		S10967 Com	QSO		17.3	17.4	17.3	16.7–17.5	+0.34
b	12 36 52.5	+25 07 51	22					17.9	18.9	18.3		+0.74
028 a	13 00 42.7	+24 58 28	18			AGN?		19.2	19.5	–		+0.26
029 a	12 40 55.2	+25 00 59	15			QSO		17.5	17.3	17.4		-0.01
b	12 40 54.5	+25 00 32	22			AGN?		16.9	18.4	17.5		+0.03
030 a	12 25 39.5	+24 58 36	6	2E 1223.1 +2515	V	SY 1.5		16.0	16.6	16.2	16.6–17.4	+0.03
031 a	12 56 04.4	+24 51 34	15			AGN?		19.5	–	–		+0.62
032 a	12 49 58.9	+24 52 33	27			AGN		18.0	18.2	18.1		-0.27
033	–	–				GCI						–
034 a	12 30 16.2	+24 51 56	16	GSC 1989 2380	S10968 Com	RS,E	M	12.9	15.3	13.8	15.0–15.4	-1.71
035 a	12 42 06.6	+24 52 23	8			QSO		18.6	18.8	18.7		+0.36
036 a	12 48 48.4	+24 50 28	9	HR 4864			G7V	-	7.2	6.4		-4.43
037 a	12 43 09.5	+24 47 06	15	GSC 1990 1861	S10969 Com	CA	K/M	12.6	15.3	13.6	15.3–15.6	-1.79
b	12 43 07.3	+24 47 17	19			AGN?		18.9	19.1	19.0		+0.37
038 a	12 25 57.9	+24 38 16	8			QSO		17.8	17.5	17.7		-0.12
039 a	12 35 33.9	+24 39 19	9	GSC 1990 2631	S10970 Com	CA?	M	12.2	14.9	13.2	14.7–15.2	-2.35
040 a	12 24 24.2	+24 36 24	16	1E 1221+2452	S10971 Com	BLL		16.7	16.9	16.8	16.6–17.8	+0.17
041 a	12 55 39.3	+24 30 26	12			AGN?		18.4	18.1	18.3		+0.19
042 a	12 56 05.8	+24 29 19	12			QSO		19.0	18.3	18.7		+0.79
b	12 56 03.0	+24 29 23	27			QSO		19.0	18.9	19.0		+0.91
043 a	12 29 41.0	+24 31 15	9	GSC 1989 3015	HZ Com	RS	G8V	9.3	10.5	9.8	9.7–9.8	-2.39
044 a	12 58 46.7	+24 27 39	3			QSO		17.7	17.8	17.7		+0.15
045 a	12 39 18.8	+24 31 44	7		S10940 Com	SY1		17.7	17.6	17.7	17–18.5	+0.69
046 a	12 53 08.2	+24 27 50	15			AGN?		19.1	19.2	19.1		+0.22
b	12 53 07.2	+24 28 05	25					19.1	20.3	18.0		+0.22
047 a	12 22 14.6	+24 24 21	23			QSO		17.1	17.5	17.3		-0.57
048 a	12 40 09.1	+24 25 31	0			QSO		16.6	16.8	16.7		-0.28
049 a	12 24 30.7	+24 21 06	14			QSO		19.0	19.2	19.1		+0.31
b	12 24 31.7	+24 20 55	20					19.0	19.4	19.2		+0.35

Table 5a. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R mag	B mag	V mag	Ampl. mag	$\log(f_X/f_V)$
	R.A	Decl.	"									
c	12 24 28.4	+24 20 39	28				K	16.1	16.7	16.3		-0.81
050	—	—				GCl						—
051 a	12 44 33.3	+24 20 05	14		S10972 Com	QSO		18.4	18.4	18.4	17.2-18.4	-0.45
052 a	12 57 31.1	+24 12 34	11					17.8	18.3	16.5		+0.82
b	12 57 31.9	+24 12 40	8		S10941 Com	BLL		15.7	17.7	17.8	17-18	+1.40
053 a	12 56 38.8	+24 12 54	32			AGN?		18.6	18.8	18.0		-0.21
054 a	12 48 40.4	+24 12 41	26			QSO		17.8	18.0	17.9		-0.15
055 a	12 27 10.6	+24 09 47	13			QSO		17.6	17.6	17.6		-0.11
056 a	12 51 42.3	+24 04 35	4		S10973 Com	SY		17.6	17.7	17.6	16.5-18	+0.35
057 a	12 58 51.5	+23 55 26	9	PGC 44513		SY		13.8	15.3	14.4		-0.41
058 a	12 28 20.1	+23 57 51	15	GSC 1989 369			K	11.9	13.1	12.4		-2.26
059 a	12 21 59.8	+23 54 43	22			G		14.2	17.5	15.4		-1.13
060	—	—										—
061 a	12 30 54.5	+23 52 03	8			AGN		17.6	17.3	17.5		-0.16
b	12 30 55.1	+23 51 48	9			AGN?		18.5	18.6	18.5		+0.24
062 a	12 34 38.6	+23 50 16	6			AGN?		17.3	18.0	17.6		+0.56
b	12 34 37.2	+23 50 08	21			G		18.0	19.1	18.4		+0.88
c	12 34 38.3	+23 49 47	24			AGN?		15.7	18.0	16.6		+0.16
d	12 34 37.1	+23 50 17	23			AGN?		17.4	19.1	—		+0.94
e	12 34 39.0	+23 50 06	6			QSO		15.4	18.4	17.1		+0.36
063 a	12 32 40.7	+23 48 06	8	GSC 1989 2738				10.3	11.1	10.3		-2.33
064 a	12 52 58.5	+23 43 13	12			UV?		20.0	22.0	-	15.5?-21	+0.77
b	12 52 56.8	+23 43 36	33	GSC 1993 604	C		M	14.4	15.8	14.9		-1.34
065 a	12 33 51.9	+23 37 45	19			AGN						—
066 a	12 25 33.0	+23 35 34	10					18.9	19.5	19.1		+0.53
b	12 25 31.5	+23 35 28	16			G		11.3	15.0	12.7		-2.03
067 a	12 35 16.4	+23 34 24	18	PG 1232+238	C		DA	17.0	16.6	16.9		-0.20
b	12 35 15.4	+23 34 20	29	GSC 1990 2552	C			13.2	14.6	13.7		-1.48
068 a	12 58 07.5	+23 29 22	20			QSO		16.9	17.3	17.1		-0.41
069 a	12 56 52.4	+23 29 50	2	GSC 1993 2793	KX Com	UV	M	11.9	14.9	13.0	13.5-16	-1.57
b	12 56 52.0	+23 29 50	6	GSC 1993 2417				12.0	15.4	13.3		-1.45
070 a	12 43 14.7	+23 31 44	4			G		17.5	18.8	18.0		-0.12
b	12 43 15.5	+23 31 22	21			QSO		18.0	18.2	18.1		-0.08
071	—	—				GCl						—
072 a	12 37 29.9	+23 29 29	12			G		18.0	19.3	18.5		+0.19
b	12 37 27.3	+23 29 11	31		C?		A0	18.3	18.1	18.2		+0.07
073 a	12 46 22.7	+23 27 09	16					18.0	19.7	18.6		+0.22
074 a	13 00 33.2	+23 20 18	29			AGN		18.8	19.1	—		+0.26
075 a	12 25 59.4	+23 21 35	7			QSO		16.8	17.9	17.2		+0.24
076 a	12 20 51.7	+23 19 15	11		S10974 Com	CA?	M	15.5	17.5	16.3	17.3-17.8	-0.97
b	12 20 51.5	+23 19 09	11					17.0	18.8	17.7		-0.41
077 a	12 41 20.9	+23 21 53	13			AGN?		18.6	18.8	-		-0.02
078 a	12 41 48.3	+23 19 49	16	GSC 1990 1120	C		K	14.4	15.8	14.9		-1.29
b	12 41 49.8	+23 19 35	13		S10975 Com	QSO		17.7	18.1	17.9	17.3-18.2	-0.09
079 a	12 38 52.2	+23 16 38	2			QSO		17.1	19.1	17.9		-0.14
b	12 38 53.5	+23 17 22	49				FG	15.5	16.7	16.0		-0.90
080 a	12 25 22.5	+23 13 45	2	HD 108154			F5		9.1	8.6		-3.69
081 a	12 46 45.2	+23 12 40	6		V?	AGN?		17.7	18.0	17.8		-0.33
b	12 46 42.9	+23 12 43	34		C?			17.2	18.0	17.5		-0.45
082 a	12 40 44.5	+23 10 46	42		C	QSO		18.0	18.0	18.0		-0.02
083 a	12 41 27.7	+23 10 07	13			QSO		18.6	19.4	18.9		+0.26
084 a	12 36 46.1	+23 10 05	22			QSO		18.2	18.8	18.4		+0.11
085	—	—										—
086 a	12 34 45.7	+23 02 11	53	GSC 1448 3631	C		K	11.8	13.9	12.6		-2.07
b	12 34 40.6	+23 00 50	55					15.9	17.5	16.5		-0.51
087 a	12 29 27.7	+22 59 47	19	GSC 1989 2450	S10976 Com	CA	M3.5V	12.9	15.2	13.8	14.9-15.5	-2.32
088 a	13 00 33.2	+22 55 41	28		KZ Com	UV	M	14.5	17.3	15.6	14.5-17.5	-1.37
089 a	12 17 49.4	+22 55 13	16			G		13.8	17.6	15.2		-1.15
b	12 17 48.9	+22 55 28	16			G		14.1	17.6	15.4		-1.07
090 a	12 18 05.2	+22 48 45	25			G		13.8	15.1	14.3		-1.60

Table 5a. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R mag	B mag	V mag	Ampl. mag	$\log(f_X/f_V)$
	R.A	Decl.	"									
091	—											—
092 a	12 21 21.1	+22 47 25	25			QSO		16.6	18.5	17.3		-0.40
093 a	12 29 12.5	+22 49 43	40	GSC 1989 1936	S10977 Com	RS+E?	FG	15.1	15.9	15.4	15.8-16.3	-1.33
094 a	12 53 36.2	+22 47 35	2	GSC 1993 876	KW Com	UV+E	M	13.7	15.8	14.5	15.2-16.6	-0.92
095 a	12 31 55.7	+22 48 05	5			AGN		17.4	17.8	17.6		-0.17
096 a	12 23 54.9	+22 45 46	19	GSC 1989 693			K0V	11.6	13.2	12.2		-2.58
b	12 23 51.7	+22 45 30	29					16.8	18.2	17.3		-0.54
097 a	12 30 36.2	+22 42 07	11			AGN		18.7	19.3	18.9		+0.22
b	12 30 37.6	+22 41 58	14				F5	17.3	18.1	17.6		-0.30
098 a	12 23 56.9	+22 39 43	53	GSC 1989 943	S10978 Com	E		14.8	15.8	15.2	15.8-16.2	-1.21
b	12 24 01.0	+22 39 39	28			BLL		18.6	19.1	18.0		-0.09
099 a	12 47 06.7	+22 37 01	41	HD 111154			G5	8.3	8.9	8.5		-3.94
100 a	12 55 04.0	+22 33 13	22			SY		15.9	17.7	16.6		-0.74
101 a	12 52 30.8	+22 33 26	8		S10979 Com	CV?		18.2	18.5	18.0	17.3-18.5	+0.13
b	12 52 32.7	+22 33 38	31			QSO		18.3	19.2	—		+0.38
102 a	12 26 30.9	+22 33 18	20					18.3	19.2	18.6		+0.23
b	12 26 31.9	+22 33 21	33			QSO		16.7	18.5	17.4		-0.25
103	—											—
104 a	12 51 47.1	+22 32 40	10	BD +23 2508	C?	CA?	M0	9.9	12.3	10.8		-1.89
105 a	12 40 00.1	+22 33 26	1		S10980 Com	QSO		16.4	17.2	16.7	16.8-18.0	+0.14
106 a	12 43 03.8	+22 26 29	51					16.8	17.6	17.1		-0.57
107 a	12 35 15.6	+22 26 02	10		S10981 Com	QSO		17.4	17.1	17.3	16.2-18.0	-0.21
b	12 35 14.6	+22 25 27	28	GSC 1448 1401	C		FG	13.4	14.2	13.7		-1.65
108 a	12 25 18.0	+22 20 19	14			QSO		17.3	17.1	17.2		-0.49
b	12 25 17.7	+22 20 31	25	GSC 1447 1748			G6V	13.8	14.4	14.0		-1.77
109 a	12 27 34.2	+22 14 31	16		C?	AGN		19.1	19.7	19.3		+0.75
b	12 27 30.4	+22 14 29	47	GSC 1447 1361	C			15.0	15.5	15.2		-0.89
c	12 27 33.7	+22 13 50	28			G		17.2	18.2	17.6		+0.07
d	12 27 33.2	+22 13 42	36			G		18.4	18.5	18.4		+0.39
e	12 27 28.8	+22 15 00	79		S10982 Com	AGN/CV		18.5	18.4	18.5	18.5-20.5	+0.43
110 a	12 25 14.8	+22 13 25	21			QSO		18.1	18.2	18.1		-0.18
111 a	12 43 13.1	+22 12 48	13		S10983 Com	AGN/CV		18.8	18.4	18.7	18.4-20:	+0.06
112 a	12 54 40.3	+22 06 24	10	HD 112196			F8V	7.1	7.6	7.1		-3.47
113 a	12 57 02.4	+22 02 02	14	GD 153	C		DA	13.4	13.3	13.4		+0.18
114 a	12 28 08.7	+22 03 22	9	GSC 1447 1989	S10984 Com	CA?	K	12.5	13.9	13.0	14.0-14.2	-2.20
115 a	12 28 37.0	+22 01 04	2				M3	16.9	19.0	17.7		-0.39
b	12 28 37.1	+22 01 14	10					18.1	19.3	18.6		-0.03
c	12 28 40.1	+22 01 27	50			G		17.2	18.7	17.8		-0.35
116 a	12 36 26.2	+22 01 48	9		S10985 Com	QSO		16.9	17.3	17.1	16.4-17.5	-0.38
b	12 36 26.6	+22 02 27	33	GSC 1448 1268	C		FG	13.6	14.7	14.0		-1.62
117 a	12 44 23.2	+21 59 45	18	HD 110789			F6V	7.7	8.2	7.7		-3.51
118 a	12 30 13.9	+21 58 12	23	BD+22 2477			G0	9.7	10.5	10.0		-3.04
119 a	12 27 00.1	+21 56 57	30		C?	AGN		18.7	18.5	18.6		+0.20
120 a	12 26 45.7	+21 52 27	54		C?			15.9	17.6	16.5		-0.26
121 a	12 35 34.3	+21 53 53	17		S10986 Com	AGN?		16.8	17.7	17.1	16.8-17.7	-0.64
b	12 35 33.1	+21 54 13	42		C		A0	16.3	16.7	16.5		-0.88
122 a	12 40 10.7	+21 46 23	27			G		16.1	17.1	16.5		-0.69
123 a	12 56 43.8	+21 40 57	13	NGC4826=M64		SY		-	9.3	8.4		-3.70
124 a	12 22 12.5	+21 37 40	5		S10987 Com	QSO		16.8	16.7	16.8	16.4-17.8:	+0.05
125 a	13 00 15.6	+21 33 12	13	HD 112974			F7V	-	8.0	7.5		-4.20
126 a	12 29 35.6	+21 37 17	29			QSO		18.7	18.5	18.6		+0.27
127 a	12 32 14.0	+21 36 53	16			QSO		17.7	18.3	17.9		+0.61
128 a	12 56 27.4	+21 31 18	14			QSO		14.7	16.1	15.2		-0.98
129	—											—
130 a	12 51 20.4	+21 25 26	6		S10988 Com	QSO		17.3	17.5	17.4	17-18.5	-0.56
131 a	12 44 31.0	+21 26 00	9			AGN		17.5	16.3	15.8		-0.95
b	12 44 33.2	+21 25 17	45					17.0	18.0	17.4		-0.31
132 a	12 33 21.0	+21 25 30	9			AGN		17.6	19.2	18.2		+0.24
133 a	12 37 35.4	+21 25 26	5			QSO		18.6	18.3	18.5		+0.08
b	12 37 36.1	+21 25 20	10			G		17.6	19.1	18.2		-0.04

Table 5a. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	$\log(f_X/f_V)$
	R.A	Decl.	"					mag	mag	mag	mag	
134 a	12 24 54.5	+21 22 47	3	PG 1222+216	S10989 Com	QSO		17.9	17.5	17.8	16-17.5	+0.83
135 a	12 59 00.3	+21 12 56	15		KY Com	UV+E	M	14.4	16.8	15.3	16.1-17.5	-1.15
136 a	12 53 17.8	+21 14 42	22	ADS 8695 ABC		double*	G8	-	6.1	5.0		-5.38
137 a	12 44 58.4	+21 14 14	14	GSC 1448 2849	S10990 Com	CA?	K	11.2	13.0	11.9	12.9-13.2	-2.36
138 a	12 38 09.9	+21 14 01	2			G		16.4	18.2	17.1		-0.59
b	12 38 11.1	+21 13 56	20	GSC 1448 1932			FG	14.2	15.4	14.7		-1.55
139 a	12 19 05.2	+21 06 32	3			AGN		12.5	14.7	13.3		-1.26
b	12 19 05.1	+21 06 54	25				A0	18.0	17.8	17.9		+0.58
140 a	12 34 27.5	+21 09 35	10			QSO		16.6	17.1	16.8		-0.57
141 a	12 39 32.1	+21 08 06	15		IR Com	CV+E		16.7	16.8	16.7	13.4-18.5	-0.14
142 a	12 32 31.1	+21 06 01	15	BD +21 2429	KK Com	EB	F8	8.9	9.8	9.2	9.54-9.77	-3.46
143 a	12 53 42.3	+21 01 45	28	GSC 1455 1009	C?			11.5	12.9	12.0		-2.54
144 a	12 25 41.9	+20 55 04	5		S10942 Com	SY1		16.0	16.1	16.0	16.1-16.6	+0.10
145 a	12 32 09.9	+20 55 10	14	HD 109118	IZ Com	L	K2	7.7	10.0	8.6	8.66-8.80	-3.68
146 a	12 20 32.9	+20 46 45	48			G		16.6	16.9	16.7		-0.92
147 a	12 40 06.4	+20 48 49	52	SAO 82428			K0	8.6	10.2	9.2		-3.63
148 a	12 42 45.1	+20 46 39	13		S10991 Com	QSO		17.8	18.3	18.0	17.2-18.5	-0.28
b	12 42 43.8	+20 46 10	27	GSC 1448 2426	C			14.8	17.3	15.7		-1.20
149 a	12 18 02.8	+20 38 56	35			GCl?		17.7	18.4	18.0		-0.24
150 a	12 32 47.4	+20 37 45	35			QSO		18.2	17.7	18.0		+0.05
151 a	12 50 46.8	+20 35 39	22					19.4	20.5	-		+0.57
b	12 50 46.3	+20 35 36	22					19.5	20.4	-		+0.53
c	12 50 47.4	+20 36 24	32			QSO		19.5	20.5	-		+0.57
152 a	12 24 00.9	+20 32 06	39		C?	QSO		17.6	17.5	17.6		-0.34
153 a	12 59 26.5	+20 29 27	17			QSO		15.4	16.4	15.8		-0.08
154 a	12 50 42.3	+20 32 06	9	SAO 82529			K5	8.4	10.3	9.1		-3.55
155 a	12 30 53.9	+20 30 06	26			QSO		18.9	19.7	19.2		+0.59
156 a	12 57 41.1	+20 23 50	14	IC 3929		AGN		13.5	15.0	14.1		-1.72
157 a	12 51 52.0	+20 22 54	10	GSC 1455 859	S10992 Com	RS+E?	M	13.3	15.8	14.2	15.6-16.0	-1.14
158 a	12 32 26.2	+20 23 27	1	GSC 1448 2731			K	11.6	14.2	12.6		-2.20
159 a	12 32 03.6	+20 09 30	3	PG 1229+204	V	SY		11.9	13.5	12.5		-0.97
b	12 32 02.0	+20 09 18	30			AGN?		18.5	18.9	18.7		+1.51
160 a	12 21 05.7	+20 05 48	12	BD+20 2710			K5	9.7	11.6	10.4		-2.69
161 a	12 18 52.3	+19 59 10	20	FAO 82233			G5	8.5	9.6	8.9		-3.56
162 a	12 35 33.0	+19 54 43	17	GSC 1448 255	KV Com	UV+E	K	13.4	15.6	14.2	15.1-16.0	-1.79
163 a	12 39 45.0	+19 54 26	9		S10993 Com	QSO		16.8	17.0	16.9	16.8-17.6	-0.52
b	12 39 44.4	+19 53 55	24					18.9	19.6	19.2		+0.40
164 a	12 49 28.7	+19 52 18	21		S10994 Com	CA?	F9	15.1	16.0	15.4	15.9-16.2	-1.22
b	12 49 27.1	+19 52 51	29	GSC 1452 250	C		FG	13.5	14.7	14.0		-1.78
165 a	12 26 42.5	+19 50 26	29			G		16.5	18.7	17.3		-0.35
166	—	—				GCl						-
167 a	12 36 19.3	+19 48 21	3	Abell 1570 041		G		17.3	19.0	17.9		+0.03
168 a	12 42 23.4	+19 47 21	18		C	QSO		15.9	16.2	16.0		-0.01
b	12 42 22.5	+19 47 26	19		C?			17.8	17.8	17.8		+0.71
c	12 42 21.6	+19 47 05	17				F9	15.9	16.2	16.0		-0.01
169 a	12 19 58.2	+19 41 55	4		S10995 Com	AGN		17.0	17.1	17.0	16.5-17.9	+0.24
170 a	12 53 04.2	+19 36 19	19	HD 111997			F7	7.4	8.1	7.7		-3.59
171 a	12 55 28.0	+19 33 26	8	ACO 1548		QSO		15.9	17.5	16.5		-0.51
172 a	12 28 54.3	+19 25 55	11			G		14.2	16.5	15.1		-1.30
173 a	12 41 20.2	+19 25 28	14		C?	AGN		17.8	18.1	17.9		-0.30
b	12 41 17.3	+19 25 30	39		S10996 Com	QSO		18.1	18.1	18.1	17.7-18.6	-0.22
174 a	12 50 15.0	+19 23 50	3			SY		-	19.0	-		+0.92
175 a	12 45 38.7	+19 20 47	6			QSO		18.5	19.8	-		+0.05
176 a	12 36 07.0	+19 19 38	7	NGC 4561		G-Pair		16.6	18.1	17.2		-0.38
177	—	—										-
178 a	12 50 43.9	+19 14 36	4			QSO		18.4	18.0	18.3		-0.03
b	12 50 43.4	+19 14 21	13			QSO		18.7	18.6	18.7		+0.13
179 a	12 18 46.5	+19 10 07	13	HD 107083			G0V	7.6	8.6	8.0		-4.10
180 a	12 56 30.7	+19 07 22	32		V?		K5	15.7	17.3	16.3		-0.94
181 a	12 47 49.1	+19 01 28	6	HD 111238			F6	7.6	8.4	7.9		-4.01

Table 5a. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R mag	B mag	V mag	Ampl. mag	$\log(f_X/f_V)$
	R.A	Decl.	"									
182 a	12 25 46.7	+18 58 46	13			G?		19.1	20.5	17.6		+0.40
b	12 25 44.4	+18 59 02	24		V?	AGN?		15.9	16.2	16.0		-0.78
c	12 25 47.4	+18 58 43	24		C	AGN		18.1	19.1	18.5		+0.22
183 a	12 43 47.5	+18 48 57	7		S10997 Com	QSO		17.6	17.5	17.6	16.8-17.6	-0.04
184 a	12 52 40.1	+18 45 05	37		S10998 Com	QSO		17.2	16.9	17.1	16.4-17.4	-0.34
185 a	12 18 30.8	+18 34 59	33		S10999 Com	QSO		16.6	17.0	16.8	16.8-18:	-0.57
186 a	12 49 19.1	+18 38 11	21	GSC 1452 292	S11000 Com	E?	M	13.8	15.9	14.6	15.8-16.1	-1.40
187 a	12 48 07.0	+18 38 13	20	QSO B1245+189	S11001 Com	QSO		16.7	16.8	16.7	16.6-17.6	-0.39
188 a	12 41 17.0	+18 34 50	6					18.4	19.0	18.6		+1.39
b	12 41 17.5	+18 34 28	17	LEDA 42543		G		11.6	14.4	12.7		-0.97
c	12 41 16.1	+18 34 04	43			G		17.0	18.1	17.4		0.91
189 a	12 46 57.3	+18 33 40	33			AGN?		16.9	16.9	16.9		-0.64
190	—	—	—									—
191 a	12 21 54.9	+18 24 35	50	GSC 1445 562				11.3	12.4	11.7		-2.81
192 a	12 19 06.5	+18 21 23	8	GSC 1445 1087	FN Com	LB		10.1	12.1	11.2	10.7-11.0	-1.99
193 a	12 23 47.5	+18 20 47	40			AGN?		17.8	17.6	17.7		-0.24
b	12 23 51.0	+18 21 08	58					16.2	17.0	16.5		-0.72
c	12 23 46.2	+18 21 07	67	4C 18.34	V	QSO		17.6	17.1	17.4		-0.36
194 a	12 38 10.3	+18 21 47	16					18.8	19.4	—		—
b	12 38 10.9	+18 22 15	15			QSO		18.2	18.7	—		—
195 a	12 29 39.0	+18 17 55	70			QSO		15.3	17.3	16.1		+0.34
196 a	12 48 15.2	+18 16 53	18	GSC 1452 779	C?			10.4	11.8	10.9		-2.18
197 a	12 47 21.7	+18 14 40	6			AGN		15.3	16.5	16.1		0.13
b	12 47 22.5	+18 14 42	13	GSC 1446 1967				10.9	12.2	11.3		-1.79
c	12 47 24.1	+18 14 38	33		C	G		15.2	17.0	—		0.26
198 a	12 41 40.0	+18 10 47	33			G?		18.8	18.5	18.7		+0.20
199 a	12 35 50.3	+18 02 43	5		C?	QSO		16.4	16.7	16.5		-0.35
b	12 35 49.4	+18 02 26	23	GSC 1446 267	C		FG	14.1	15.5	14.6		-1.11
200 a	12 48 04.1	+18 00 09	10			QSO		13.3	15.6	14.2		-1.55
b	12 48 03.3	+17 59 51	14					16.1	18.4	17.0		-0.43
201 a	12 54 48.6	+17 56 04	19					18.2	19.7	18.8		+0.17
b	12 54 47.0	+17 56 24	18			AGN?		18.2	—	—		-0.01
202 a	12 50 36.2	+17 57 06	5		S11002 Com	QSO		16.2	16.3	16.2	16.0-17.0	-1.03
203 a	12 44 06.6	+17 48 31	34			BLL?		14.2	18.2	15.7		-0.89
b	12 44 06.9	+17 47 56	6			G		17.9	19.3	19.0		+0.44
204 a	12 44 58.8	+17 33 00	2	2E 1242.4+1749	S11003 Com	QSO		16.5	16.8	16.6	16.6-18.3	-0.17
205 a	12 34 58.7	+17 32 24	26	BD+18 2629			F8	10.2	11.0	10.2		-3.32
206 a	12 43 28.6	+17 29 58	11	GSC 1446 640			FG	12.7	14.1	13.2		-1.91
b	12 43 26.5	+17 29 37	47	1228+1738		QSO		18.1	17.5	17.9		-0.03
207 a	12 31 21.1	+17 21 50	2			QSO		18.3	17.9	18.2		-0.02
b	12 31 19.8	+17 21 43	21					17.1	18.8	17.7		-0.22
208 a	12 24 24.7	+17 20 06	33		C			16.8	17.2	16.8		-0.71
b	12 24 23.9	+17 20 22	40		C			16.9	17.3	17.1		-0.59
c	12 24 19.7	+17 20 21	58		V?	AGN?		14.8	15.5	15.1		-1.39
209 a	12 44 10.8	+17 21 05	6	2E 1241.6-1737		QSO		15.1	15.7	15.3		-1.02
b	12 44 10.8	+17 21 15	9					15.5	16.5	15.9		-0.78
210 a	12 38 48.4	+17 16 46	31			G		17.0	18.3	17.5		-0.08
211 a	12 42 25.9	+17 13 50	38		S11004 Com	AGN/CV		20.0	19.7		17.7-21:	+0.68
212 a	12 46 13.6	+17 10 13	50			G		11.9	15.2	13.1		-2.19
213 a	12 52 22.9	+17 06 31	13	HD 111892			F8	-	7.6	7.0		-4.51
214 a	12 28 29.1	+17 05 04	11	NGC 4450		AGN		11.2	12.0			-2.49
215 a	12 33 05.1	+17 01 33	4			BLL?		17.7	18.1	17.9		+0.77
216 a	12 35 28.8	+17 00 37	11	2E 1232.9+1716		BLL?		17.5	18.8	18.0		-0.01
217	—	—	—			GCl						—
218 a	12 32 04.1	+16 55 23	8	LBQS 1229+1711		QSO		17.4	18.0	17.6		-0.37
219 a	12 41 33.3	+16 55 13	29					15.8	18.1	16.7		-0.61
b	12 41 32.1	+16 55 02	21					19.4	20.6	—		+0.73
c	12 41 33.3	+16 54 16	28			G		17.3	18.8	17.9		-0.13
220 a	12 44 07.1	+16 52 47	56			GCl		15.5	18.2	16.5		-0.71
221 a	12 37 24.9	+16 50 30	25					17.9	18.5	18.1		+0.65

Table 5a. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R mag	B mag	V mag	Ampl. mag	$\log(f_X/f_V)$
	R.A	Decl.	"									
b	12 37 26.5	+16 51 12	39					16.4	18.5	17.2		+0.29
222 a	12 48 04.1	+16 48 21	22	GSC 1449 473	S11005 Com	E?	FG	13.7	14.5	14.0	14.4-14.7	-1.75
223 a	12 29 44.7	+16 40 18	21			AGN		19.3	19.1	19.2		+1.15
b	12 29 47.7	+16 40 22	58			AGN?		16.3	16.7	16.5		+0.07
c	12 29 44.5	+16 40 04	8			QSO		19.3	19.7	—		+1.12
224 a	12 19 07.4	+16 33 00	14	HD 107146			G2V	11.3	12.6	11.8		-1.63
225 a	12 59 52.1	+16 30 41	11					16.0	18.2	16.8		-0.32
b	12 59 53.2	+16 30 39	18			G		16.4	16.5	16.4		-0.48
226 a	12 19 47.0	+16 30 51	8	GSC 1445 1539	CN Com	RS		11.9	13.0	12.3	13.1-14.0	-2.32
b	12 19 46.8	+16 31 22	25		C		K7	16.0	18.1	16.8		-0.52
227 a	12 57 37.0	+16 30 49	15		S11006 Com	QSO		16.0	16.1	16.0	16.1-16.8	-0.85
b	12 57 34.4	+16 30 26	47			G		15.0	16.1	15.4		-1.09
228 a	12 41 48.6	+16 32 22	22		V?		K	15.7	16.6	16.0		-0.71
229 a	12 34 36.6	+16 31 56	6		C?		K	13.8	16.8	14.9		-0.63
b	12 34 37.2	+16 32 14	16			AGN?		17.2	17.5	17.3		+0.33
230 a	12 36 24.6	+16 32 23	19	4C16.33		G(E3pe)		15.1	16.6	15.7		+0.07
b	12 36 25.8	+16 32 18	27			G		13.8	15.5	14.4		-0.45
231 a	12 43 57.7	+16 22 54	5	2E 1241.4+1639	V	QSO		18.2	18.8	18.4		+0.26
232 a	12 55 03.2	+16 16 06	37					16.4	18.1	17.0		-0.44
233 a	12 42 51.0	+16 18 14	27			AGN?		18.4	19.4	18.8		+0.19
234 a	12 34 38.3	+16 05 42	26			AGN?		18.5	18.5	18.5		+0.00
235 a	12 35 44.3	+16 05 36	34			QSO		16.1	17.4	16.6		-0.51
236 a	12 39 46.7	+16 05 48	31	GSC 1446 2344			FG	13.3	14.5	13.8		-1.35
b	12 39 42.9	+16 06 12	30			G		16.6	17.6	16.6		-0.23
c	12 39 46.7	+16 05 19	50					15.1	16.4	15.6		-0.63
d	12 39 45.8	+16 05 57	15			AGN		17.2	20.1	—		+0.94
237	—	—	—									—
238 a	12 32 35.9	+16 02 29	17	GSC 1446 1511			FG	10.6	12.5	11.3		-2.98

Table 5b. Optical objects in or near the ROSAT X-ray error circle in the Sge field.

No	Coordinates (2000.0)		D	Designation	Name if	Type	Spec.	R	B	V	Ampl.	log(f_x/f_v)
	R.A	Decl.	"		variable			mag	mag	mag	mag	
001 a	19 37 43.5	+24 15 40	30	HD 185334			A0	7.8	7.9	7.8		-3.90
b	19 37 41.7	+24 14 57	20	GSC 2138 966				13.4	12.5	13.1		-1.78
c	19 37 44.8	+24 15 27	33					12.4	14.1	13.0		-1.82
002 a	19 56 29.0	+23 54 45	13	HD 345287	B0 Vul	EA	F0	10.8	10.5	10.7	10.5-13.3	-3.15
b	19 56 29.7	+23 54 12	47	GSC 2140 745				11.6	12.6	12.0		-2.63
003 a	19 58 00.1	+23 54 15	15	HD 189181			A0	8.1	12.8	9.9		-3.06
b	19 57 57.5	+23 53 50	39	GSC 2141 123				11.2	13.5	12.1		-2.18
004 a	20 07 02.6	+23 52 33	14	GSC 2154 555	S11007 Vul	CA		11.9	13.7	12.6	13.6-13.9	-2.53
b	20 07 03.6	+23 52 55	18		C			12.6	14.3	13.2		-2.29
c	20 07 00.5	+23 52 42	28		C			12.4	16.0	13.8		-2.05
005 a	19 53 14.6	+23 50 52	19		C			13.5	14.8	14.0		-1.62
006 a	20 20 03.7	+23 38 17	7	HD 193554			G5	8.2	9.1	8.5		-3.17
b	20 20 02.8	+23 38 44	37					14.8	16.8	15.6		-0.33
007 a	19 45 52.9	+23 30 12	20	GSC 2139 1736	S11008 Vul	CA		11.9	12.9	12.3	12.4-12.8	-1.79
008 a	19 57 26.3	+23 28 47	8	GSC 2141 1977	C			11.2	13.5	12.9		-2.00
b	19 57 24.3	+23 28 57	30		C			15.1	16.9	15.8		-0.84
009 a	19 38 38.9	+23 20 52	8	GSC 2138 2017	S11009 Vul	CA		13.9	15.8	14.6	15.8-16.4	-1.81
b	19 38 37.7	+23 20 05	43		C			13.0	14.8	13.7		-2.17
010 a	19 43 12.6	+23 19 39	8		C			16.8	18.0	17.3		-0.27
b	19 43 11.6	+23 19 47	15		V?			17.2	18.8	17.8		-0.07
c	19 43 13.6	+23 20 19	34		C			15.9	16.7	16.2		-0.71
d	19 43 10.9	+23 20 21	42	GSC 2139 2039	C			14.8	15.1	14.9		-1.23
e	19 43 14.5	+23 20 17	39		C			14.3	15.5	14.8		-1.27
f	19 43 11.7	+23 19 10	39		C			13.9	15.0	14.3		-1.47
g	19 43 15.1	+23 19 12	43	GSC 2139 2375	C		B0.5	-	12.0	11.8		-2.47
011 a	20 13 51.2	+23 17 49	21	HD 346128			G5	11.6	12.3	11.9		-2.43
b	20 13 54.1	+23 17 08	38		C			15.0	16.6	15.6		-0.95
012 a	20 15 34.6	+23 12 59	27		V?			12.7	13.8	13.1		-2.26
b	20 15 33.2	+23 13 27	32		C			15.4	16.9	16.0		-1.10
c	20 15 32.3	+23 13 32	37		C			13.0	15.6	14.0		-1.90
013 a	20 17 38.4	+23 10 01	20	HD 193078			F5	8.3	9.0	8.6		-3.48
b	20 17 37.6	+23 09 23	36					13.9	15.8	14.6		-1.08
c	20 17 37.0	+23 09 52	35					15.8	16.9	16.2		-0.44
014 a	20 18 05.3	+23 03 52	11	GSC 2155 1563	C			11.3	12.5	11.8		-2.72
b	20 18 06.8	+23 03 52	17		C			14.6	15.5	14.9		-1.48
c	20 18 07.4	+23 04 16	28		C			14.4	16.2	15.1		-1.40
d	20 18 04.5	+23 03 21	44		C			14.5	15.4	14.8		-1.52
015 a	20 10 05.9	+23 03 48	30	HD 345835			G5	10.7	12.2	11.3		-2.86
b	20 10 09.2	+23 04 13	32					14.3	15.4	14.7		-1.50
c	20 10 06.4	+23 04 54	40	GSC 2154 2253				10.8	13.0	11.6		-2.74
d	20 10 07.2	+23 03 41	35					14.0	15.3	14.5		-1.58
e	20 10 09.3	+23 04 44	43	GSC 2154 1455				13.1	14.2	13.5		-1.98
f	20 10 09.5	+23 03 46	46					14.2	15.4	14.7		-1.50
016 a	20 19 49.0	+22 56 33	13	GSC 2155 1614	S11010 Vul	UV?		10.4	13.0	11.4	12.2-13.3	-1.65
b	20 19 47.5	+22 56 47	35		C			15.4	16.6	15.9		+0.15
c	20 19 48.0	+22 57 00	43		C			13.1	14.7	13.7		-0.73
d	20 19 49.6	+22 55 36	44		C			14.8	15.9	15.2		-0.13
017 a	20 03 23.3	+22 59 32	3		C			17.0	17.8	17.3		-0.45
b	20 03 21.4	+22 59 42	26		C			14.5	18.1	15.9		-1.01
c	20 03 24.7	+22 59 56	33	GSC 2141 2428	C			13.0	14.8	13.7		-1.89
d	20 03 23.1	+23 00 13	42		C			15.1	16.3	15.6		-1.13
e	20 03 23.4	+22 58 50	41		C			15.0	16.2	15.5		-1.17
018 a	19 52 12.4	+22 52 47	9		S11011 Vul	CA?		14.5	15.6	14.9	15.2-15.7	-1.28
b	19 52 11.7	+22 53 11	34	GSC 2140 102	C			11.7	14.6	12.8		-2.12
019 a	19 59 36.1	+22 43 00	-	NGC 6853		PN		-	8.6			-3.36
020 a	20 00 43.7	+22 42 39	6	HD 189733			G5	-	8.9	7.8		-3.52
b	20 00 41.5	+22 42 35	25					16.0	16.7	16.3		-0.12
021 a	20 05 41.9	+22 39 59	14	1E 2003+2205	QQ Vul	AM		14.5	14.8	14.6	14.4-17.0	+0.62
b	20 05 41.6	+22 40 28	40	GSC 2141 3961				13.3	14.1	13.6		+0.22

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	log(f_x/f)
	R.A	Decl.	"					mag	mag	mag	mag	
022 a	20 07 31.0	+22 34 29	5		C			14.8	16.8	15.6		-1.30
b	20 07 31.7	+22 34 18	12		S11012 Vul	CA		13.9	14.9	14.3	14.9-15.2	-1.82
c	20 07 29.3	+22 34 10	33	GSC 2154 199	C			12.0	12.9	12.3		-2.62
d	20 07 30.6	+22 33 53	38		C			13.8	15.6	14.5		-1.74
023 a	19 47 34.9	+22 28 58	2	GSC 1627 1512	S11013 Vul	CA		10.9	13.5	11.9	13.4-14.0	-1.96
024 a	20 02 48.6	+22 28 27	7	2E 2000.6+2219	V362 Vul	QSO		13.1	15.9	14.2	16.0-17.7	-1.59
025 a	20 07 50.3	+22 26 19	30	HD 345865			A2	11.3	12.0	11.6		-2.76
b	20 07 49.2	+22 26 00	8					15.9	16.8	16.2		-0.92
c	20 07 48.3	+22 25 47	14					15.7	16.3	15.9		-1.04
d	20 07 49.3	+22 25 30	33					14.5	16.2	15.1		-1.36
026 a	20 10 49.0	+22 21 37	6	HD 346034			G5	11.1	12.1	11.5		-2.48
b	20 10 48.4	+22 21 47	7					16.0	16.9	16.3		-0.56
c	20 10 50.5	+22 21 34	26					15.6	16.3	15.9		-0.72
027 a	20 01 51.4	+22 16 49	11	HD 345604			F2	8.2	8.6	8.2		-4.24
b	20 01 49.4	+22 16 48	30					14.9	17.0	15.7		-1.24
028 a	20 14 04.5	+22 13 22	3	HD 192424	NSV 12927	CA?	A0	7.9	7.2	7.2		-3.90
b	20 14 02.3	+22 13 35	35					15.0	14.1	14.7		-0.90
c	20 14 03.5	+22 12 55	33	GSC 1630 2556				17.9	13.9	14.6		-0.94
029 a	20 16 17.6	+22 06 10	26	HD 346177	V?		K0	9.0	10.4	9.5		-3.39
b	20 16 18.7	+22 06 19	14		C			15.7	16.9	16.2		-0.71
030 a	19 49 58.6	+22 04 36	13	HD 345032			G0	11.3	11.9	11.5		-2.81
031 a	20 01 24.5	+22 03 38	24	HD 345492	S11014 Vul	CA	A2	10.8	12.0	11.3	11.9-12.1	-2.71
b	20 01 25.6	+22 03 31	10		C			16.3	16.9	16.5		-0.63
c	20 01 25.3	+22 03 19	22		C			15.0	16.9	15.7		-0.95
d	20 01 25.4	+22 03 05	34		C			14.5	16.8	15.4		-1.07
e	20 01 26.6	+22 03 08	30		C			13.5	14.1	13.7		-1.75
f	20 01 28.2	+22 03 42	29		C			15.4	16.3	15.7		-0.95
g	20 01 23.4	+22 03 44	39	GSC1629 3566	C			13.6	14.3	13.9		-1.67
032 a	20 11 35.4	+21 59 12	29	GSC 1630 3418	C			12.9	14.2	13.4		-2.26
b	20 11 38.1	+21 58 41	21		V?			14.2	15.1	14.5		-1.82
c	20 11 37.3	+21 58 08	44		C			14.3	15.3	14.7		-1.74
033 a	20 08 21.6	+21 47 33	22		S11015 Vul	CV?		14.4	14.8	14.6	14.5-14.8	-1.60
b	20 08 20.4	+21 47 58	23		C			15.9	16.8	16.2		-0.96
c	20 08 20.0	+21 48 13	34	GSC 1630 2694	C			13.1	14.6	13.7		-1.96
d	20 08 20.3	+21 47 35	31		C			15.8	16.5	16.1		-1.00
e	20 08 23.7	+21 47 35	31		C			14.8	16.7	15.5		-1.24
034 a	20 09 23.7	+21 43 02	10	HD 345887	C		A3	11.2	12.1	11.5		-2.91
b	20 09 26.4	+21 42 33	37		C			14.5	15.9	15.0		-1.51
035 a	19 59 30.2	+21 33 27	8	HD 345513			F7	9.3	10.1	9.6		-2.63
b	19 59 31.3	+21 33 59	28					13.3	15.1	14.0		-0.87
c	19 59 32.5	+21 33 21	32					15.0	15.6	15.2		-0.39
036 a	19 55 37.5	+21 29 35	17		V?			13.1	15.6	14.0		-1.78
b	19 55 37.4	+21 29 23	28		C			15.7	17.4	16.3		-0.86
c	19 55 40.0	+21 29 45	31		C			14.2	15.3	14.6		-1.54
d	19 55 37.7	+21 30 22	31		C			15.0	15.5	15.2		-1.30
037 a	20 19 07.0	+21 21 21	12	GSC 1643 998	S11016 Sge	CA		11.4	12.5	11.8	12.6-12.9	-2.03
b	20 19 06.9	+21 21 03	10	GSC 1643 448	C			12.5	14.2	13.1		-1.51
c	20 19 07.9	+21 21 13	21		C			15.6	16.8	16.2		-0.27
038 a	19 43 56.3	+21 18 19	33		C	AGN?		16.5	19.4			+0.45
039 a	20 07 54.6	+21 15 32	29	GSC 1629 3002	C			12.7	14.4	13.3		-2.20
b	20 07 53.4	+21 16 09	26	GSC 1629 2204	C			14.1	14.6	14.3		-1.80
c	20 07 56.6	+21 16 24	31	GSC 1629 2882	V?			13.1	14.4	13.6		-2.08
d	20 07 56.2	+21 15 41	24		C			15.0	15.9	15.3		-1.40
e	20 07 56.5	+21 15 29	36		C			16.1	16.9	16.4		-0.96
f	20 07 52.3	+21 16 14	42		C			15.1	16.6	15.7		-1.24
g	20 07 55.4	+21 15 30	30	GSC 1629 130	C			13.2	15.8	14.2		-1.84
h	20 07 54.9	+21 16 24	24		C			15.5	16.4	15.8		-1.20
i	20 07 53.8	+21 14 50	42		C			15.9	16.4	16.1		-1.08
j	20 07 53.8	+21 15 15	50		—			12.8	14.0	—		-2.15
040 a	19 53 04.9	+21 14 50	21		V405 Vul	UGSU		16.2	16.8	16.1	15.0-18.0	-0.76

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	$\log(f_x/f_y)$
	R.A.	Decl.	"					mag	mag	mag	mag	
b	19 53 04.9	+21 14 37	10		C			16.0	16.8	16.3		-0.68
c	19 53 05.8	+21 14 39	10		C			15.1	17.4	16.0		-0.80
d	19 53 05.6	+21 14 49	19		C			14.7	15.1	14.9		-1.24
e	19 53 07.0	+21 14 40	25		C			15.1	16.0	15.4		-1.04
f	19 53 06.8	+21 14 16	24		C			13.6	14.5	13.9		-1.64
g	19 53 03.6	+21 14 19	28		C			15.3	16.0	15.6		-0.96
h	19 53 04.6	+21 13 59	34		C			13.1	16.4	14.3		-1.48
041 a	20 13 42.2	+21 07 44	24	HD 346216			F2	8.2	9.0	8.5		-4.11
b	20 13 42.4	+21 07 22	13					12.2	13.2	12.6		-2.47
c	20 13 43.7	+21 07 42	18					11.9	13.7	12.6		-2.47
042 a	20 06 24.7	+21 07 02	9		S11017 Sge	CV?		15.2	15.5	15.3	15.8-16.5	-1.19
b	20 06 25.3	+21 07 00	13		C			15.6	16.3	15.9		-0.95
c	20 06 22.2	+21 07 13	37		C			14.5	15.7	15.0		-1.31
d	20 06 25.2	+21 06 28	43		C			15.4	16.7	15.9		-0.95
043 a	20 08 29.4	+21 05 29	34	GSC 1630 973	S11018 Sge	BY?		11.3	12.7	11.8	12.4-13.0	-2.67
b	20 08 30.4	+21 04 39	21	GSC 1630 968	C			12.4	14.0	13.0		-2.19
c	20 08 31.4	+21 05 06	13		C			15.5	15.5	15.5		-1.19
d	20 08 28.8	+21 04 54	26		C			14.6	15.3	14.9		-1.43
e	20 08 32.6	+21 05 20	35		C			15.9	16.6	16.2		-0.91
f	20 08 33.1	+21 05 00	36		C			14.5	15.3	14.8		-1.47
044 a	20 12 04.1	+20 57 52	12		S11019 Sge	?		17.2	18.7	17.8	16.5-19.0	-0.29
b	20 12 05.3	+20 57 43	25		C			15.0	16.1	15.4		-1.25
c	20 12 03.0	+20 58 11	29		C			15.5	15.9	15.7		-1.13
d	20 12 01.5	+20 57 39	28		C			15.6	16.5	15.9		-1.05
045 a	20 09 56.6	+20 54 53	12	HD 191570			F5IV	-	6.9	6.5		-4.31
b	20 09 56.4	+20 54 59	20	SAO 88275			F2	-	9.7	8.8		-3.39
046 a	19 59 21.4	+20 55 25	7		C			12.7	13.1	12.9		-2.26
b	19 59 21.1	+20 55 37	7	GSC 1628 1030	V?			12.5	13.9	13.0		-2.22
c	19 59 20.6	+20 55 38	9		C			15.6	14.5	15.2		-1.34
047 a	20 18 58.7	+20 51 52	38	HD 357949			K0	8.0	9.5	8.6		-3.70
048 a	19 53 12.9	+20 43 57	31	GSC 1628 3835				13.6	13.6	13.6		-1.93
b	19 53 12.3	+20 43 39	38	GSC 1628 1335				12.5	14.6	13.3		-2.05
c	19 53 09.0	+20 44 46	44	GSC 1628 1324				12.4	14.4	13.2		-2.09
d	19 53 08.8	+20 44 27	34					16.0	16.9	16.3		-0.85
e	19 53 10.9	+20 44 14	6		S11020 Vul	CV?		18.1	18.6		17.3-20.0	-0.06
f	19 53 09.3	+20 44 12	24					16.3	17.1	16.6		-0.73
g	19 53 10.1	+20 44 30	22					16.5	17.5	16.9		-0.61
h	19 53 10.7	+20 43 49	24			CV?		15.0	17.2	15.8		-1.05
049 a	20 06 48.2	+20 41 55	41	GSC 1629 566	HH Sge	SRB		11.3	12.4	11.7	11.1-12.5	-2.23
b	20 06 48.9	+20 41 05	13					14.6	15.7	15.0		-0.91
c	20 06 46.9	+20 41 09	19					12.5	14.5	13.3		-1.59
d	20 06 45.5	+20 41 15	39					16.0	16.5	16.2		-0.43
050 a	19 41 07.2	+20 08 15	39	HD 186001			F0	8.4	8.8	8.4		-3.79
b	19 41 08.2	+20 08 43	5	GSC 1610 0461	V?			10.2	11.5	10.5		-2.95
051 a	20 17 42.2	+20 08 10	11		C?			15.1	16.0	15.4		-1.21
b	20 17 43.3	+20 08 25	26		C			15.9	16.7	16.2		-0.89
c	20 17 41.4	+20 08 36	40		C			15.8	16.6	16.1		-0.93
d	20 17 39.8	+20 08 04	41					15.7	16.4	16.0		-0.97
052 a	19 57 03.1	+20 05 09	9	GSC 1624 703				10.7	12.1	11.2		-2.67
b	19 57 01.4	+20 05 45	35		NSV 12597, C			12.4	14.6	13.2		-1.87
c	19 57 01.1	+20 05 32	28		C			13.5	16.5	14.6		-1.31
d	19 57 03.6	+20 05 47	36		C			15.5	16.5	15.9		-0.79
e	19 57 01.9	+20 04 55	21		C			11.7	13.6	12.4		-2.19
f	19 57 02.5	+20 04 45	29		C			14.1	15.8	14.7		-1.27
g	19 57 01.1	+20 05 12	22		C			14.3	16.5	15.1		-1.11
053 a	19 53 54.8	+19 59 02	17	HD 350742	C		G0	10.1	10.7	10.0		-2.31
b	19 53 55.3	+19 58 40	15		C			14.3	16.1	15.0		-0.31
054 a	20 13 09.1	+19 57 00	12		V?			15.3	16.1	15.6		-1.19
b	20 13 08.6	+19 56 40	10		C			15.2	16.0	15.5		-1.23
c	20 13 09.2	+19 57 42	54					11.3	12.2	12.0		-2.75

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	log(f_x/f)
	R.A	Decl.	"					mag	mag	mag	mag	
055 a	20 02 01.0	+19 55 52	11	GSC 1625 1567	S11021 Sge	CA		11.2	12.3	11.6	12.3-12.5	-2.81
b	20 02 00.2	+19 55 18	25		C?			14.5	14.6	14.5		-1.65
056 a	19 40 28.2	+19 38 28	20	HD 350270	C?		G0	9.0	11.9	10.1		-3.33
057 a	19 56 59.3	+19 35 16	17	GSC 1624 2660	C			11.4	12.1	11.7		-2.60
b	19 56 57.4	+19 34 55	18		C			13.0	15.4	13.9		-1.72
c	19 56 57.6	+19 34 49	21		C			14.3	15.7	14.8		-1.36
d	19 56 58.2	+19 35 34	29		C			13.0	13.5	13.2		-2.00
e	19 56 55.2	+19 35 45	58					9.3	10.0	9.9		-3.51
058 a	19 44 33.1	+19 32 13	7		V?			15.4	16.9	16.0		-0.19
b	19 44 33.1	+19 31 42	35	HD 350379	C		G0	9.3	9.9	9.3		-2.87
c	19 33 34.7	+19 32 36	35		C			11.7	14.5	12.8		-1.47
059 a	19 40 37.5	+19 30 05	12		C			13.8	16.7	14.9		-1.34
b	19 40 36.5	+19 29 21	39	GSC 1610 800	C			12.3	14.2	13.0		-2.10
c	19 40 39.9	+19 30 20	39		C			14.9	15.8	15.2		-1.22
060 a	19 19 20.5	+19 24 01	27	GSC 1639 2264	V?			12.7	13.4	13.0		-2.19
b	19 19 18.2	+19 23 09	42		C			15.6	15.7	15.6		-1.15
061 a	20 11 13.0	+19 21 54	14	HD 191837			F5V	7.3	8.0	7.6		-4.19
062 a	20 17 43.8	+19 14 53	32	HD 193080			F8	8.2	8.8	8.4		-2.86
b	20 17 43.7	+19 14 31	10					15.8	12.9	14.7		-0.34
c	20 17 42.3	+19 14 00	28					15.2	16.0	15.5		-0.02
063 a	20 18 15.9	+19 04 17	9		C			15.9	16.6	16.2		-0.85
b	20 18 14.5	+19 04 52	49					16.3	15.9			-1.19
064 a	19 53 00.2	+18 51 17	5		C			16.8	17.4	17.0		-0.58
b	19 53 00.2	+18 51 35	20		C			16.6	17.6	17.0		-0.58
c	19 52 59.3	+18 51 31	24		C			16.5	18.0	17.1		-0.54
d	19 53 01.3	+18 51 31	17		C			17.1	18.7	17.7		-0.30
e	19 53 01.1	+18 50 45	32		C			13.4	14.4	-		-3.66
f	19 53 01.7	+18 51 34	24		V?			17.6	18.5	-		-0.28
g	19 53 03.5	+18 51 59	59					10.7	12.6	11.7		-2.70
065 a	20 07 41.1	+18 39 17	40	HD 351475	C			9.9	10.4	9.9		-3.21
b	20 07 39.1	+18 38 52	20		V?			14.4	14.9	14.6		-1.33
c	20 07 42.4	+18 38 24	35		C			13.6	14.7	14.0		-1.57
d	20 07 41.5	+18 38 12	33		C			14.3	16.0	14.9		-1.21
066 a	20 05 55.8	+18 38 48	20	GSC 1621 268	S11022 Sge	CA?		-	13.5	12.8	13.2-13.8	-2.16
b	20 05 56.8	+18 38 54	22		C			12.7	13.0	12.8		-2.16
067 a	19 47 23.3	+18 32 03	35	HD 187076	δ Sge	LB	M2II+B9V		5.5	3.8	3.75-3.83	-5.85
068 a	20 11 14.0	+18 30 52	22		C			14.3	14.6	14.4		-1.15
b	20 11 12.2	+18 30 48	21		C			15.0	15.9	15.3		-0.79
c	20 11 14.8	+18 30 39	23		C			15.0	16.3	15.5		-0.71
d	20 11 14.8	+18 30 34	23		C			14.6	15.2	14.8		-0.99
e	20 11 13.3	+18 30 21	12		C			13.2	15.3	14.0		-1.31
f	20 11 11.7	+18 31 07	41	HD 351621	C			11.3	10.8	11.1		-2.47
069 a	19 49 30.9	+18 28 53	15		C			14.4	16.5	15.2		-1.26
b	19 49 29.8	+18 28 58	12		C			14.3	16.7	15.2		-1.26
c	19 49 29.3	+18 28 34	14		C			15.5	17.0	16.1		-0.90
d	19 49 29.8	+18 28 19	27		C			14.2	16.2	15.0		-1.34
e	19 49 29.9	+18 28 07	39		C			14.9	16.7	15.6		-1.10
f	19 49 28.0	+18 29 08	35		C			14.9	16.4	15.5		-1.14
g	19 49 31.2	+18 29 07	28		C			13.5	14.6	13.9		-1.78
070 a	19 55 57.9	+18 28 04	13		C			14.0	16.2	14.8		-1.02
b	19 55 59.5	+18 28 22	39		C			11.9	13.5	12.5		-1.94
c	19 55 56.5	+18 28 19	16		V?			16.8	18.4	17.4		+0.02
d	19 55 57.0	+18 28 59	54					11.2	12.4			-2.21
071 a	20 05 08.3	+18 25 10	20		C			14.3	15.4	14.7		-1.65
b	20 05 06.8	+18 25 26	9		C?			15.7	17.6	16.4		-0.97
c	20 05 08.4	+18 25 35	17		-			18.9	18.6	-		-0.31
072 a	20 02 33.2	+18 22 06	17	GSC 1621 1090	C			12.1	13.7	12.7		-2.26
b	20 02 35.1	+18 21 49	16		C			13.8	16.0	14.6		-1.50
c	20 02 35.0	+18 21 37	27		C			15.2	16.7	15.8		-1.02
d	20 02 35.2	+18 21 45	30		C			15.6	16.6	16.0		-0.94

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	log(f_x/f_y)
	R.A	Decl.	"					mag	mag	mag	mag	
e	20 02 32.1	+18 22 19	37		C			15.3	16.4	15.7		-1.06
073 a	20 06 31.9	+18 17 59	10		C			15.3	17.5	16.1		-0.66
b	20 06 28.3	+18 17 46	46		C			14.9	16.8	15.6		-0.86
074 a	20 07 20.6	+18 10 12	13		S11023 Sge	CA?		11.4	12.6	11.9	12.6-12.8	-2.52
b	20 07 23.0	+18 10 36	37		C			13.1	14.4	13.6		-1.84
c	20 07 17.9	+18 10 21	37		C?			13.5	15.0	14.1		-1.64
075 a	19 54 55.9	+18 04 52	26		V?			17.7	18.1	17.9		-0.08
b	19 54 54.2	+18 04 31	12		C			16.4	17.5	16.8		-0.52
c	19 54 53.8	+18 04 51	28		C			14.4	16.7	15.3		-1.12
d	19 54 53.4	+18 04 10	30		C			16.3	16.6	16.4		-0.68
e	19 54 56.4	+18 04 02	34		C			16.4	16.7	16.5		-0.64
076 a	19 40 05.8	+18 00 50	12	HD 185758			G1II	4.5	5.4	4.5		-5.64
077 a	20 09 25.1	+18 00 51	34		C			11.2	13.0	11.9		-2.66
b	20 09 23.8	+18 01 06	29		C			16.0	16.4	16.2		-0.94
078 a	20 15 22.8	+17 58 42	11		C			15.3	16.5	15.8		-1.02
b	20 15 23.0	+17 58 40	8		C			15.4	16.4	15.8		-1.02
c	20 15 24.4	+17 57 56	20		C			14.0	15.0	14.4		-1.58
d	20 15 21.9	+17 59 05	37		C			15.7	16.5	16.0		-0.94
e	20 15 23.4	+17 57 56	40		C			13.8	14.5	14.1		-1.70
f	20 15 25.9	+17 58 22	37		C			15.9	16.8	16.2		-0.86
079 a	20 17 38.7	+17 56 36	9		V?			16.0	17.0	16.4		-0.72
b	20 17 37.6	+17 56 31	21		C			15.9	16.1	16.0		-0.88
c	20 17 39.0	+17 56 55	25		C			15.6	16.2	15.8		-0.96
080 a	20 07 36.0	+17 42 17	12	UBV 17292	WZ Sge	UGSU		12.5	13.6	12.9	7.0-15.5	-1.40
b	20 07 38.5	+17 42 30	36					13.4	14.9	14.0		-0.96
081 a	20 04 36.0	+17 40 42	22	GSC 1621 2087	S11024 Sge	CA		11.7	12.9	12.2	12.9-13.1	-2.17
b	20 04 35.1	+17 40 22	6		C			16.2	17.2	16.6		-0.41
c	20 04 36.1	+17 40 17	10		C			14.6	15.7	15.0		-1.05
d	20 04 33.2	+17 40 32	35		C			13.4	13.6	13.5		-1.65
e	20 04 38.0	+17 40 34	37		C			15.7	16.3	15.9		-0.69
f	20 04 38.3	+17 40 12	40		C			13.0	15.0	13.8		-1.53
082 a	19 42 58.0	+17 37 02	10		C			17.8	18.3	18.0		-0.17
b	19 42 58.0	+17 37 10	7		C			17.2	18.1	17.5		-0.37
c	19 42 56.5	+17 37 04	29		V?			16.1	16.8	16.4		-0.81
d	19 42 57.5	+17 37 29	25		C			14.8	16.9	15.6		-1.13
e	19 43 01.5	+17 36 36	53		C			14.3	16.0	14.9		-1.41
083 a	20 05 44.9	+17 35 54	27	HD 190720			K2	11.7	9.3	10.8		-2.30
b	20 05 42.5	+17 35 09	40					14.5	15.5	14.9		-0.66
084 a	19 50 41.7	+17 31 43	15		C			16.3	17.8	16.9		-0.60
b	19 50 42.6	+17 31 48	16		C			16.5	17.7	17.0		-0.56
c	19 50 43.8	+17 31 32	20		C			15.4	17.1	16.0		-0.96
d	19 50 40.2	+17 31 24	33		C			14.6	16.5	15.3		-1.24
e	19 50 43.5	+17 31 20	20		C			16.3	17.2	16.6		-0.72
f	19 50 41.5	+17 31 09	27		C			14.6	15.5	14.9		-1.40
g	19 50 39.8	+17 31 20	39		C			15.1	16.0	15.4		-1.20
h	19 50 43.7	+17 31 05	33		C			14.4	16.1	15.0		-1.36
085 a	20 14 17.2	+17 27 36	4	GSC 1622 826	S11025 Sge	CA		11.3	12.7	11.8	12.3-12.5	-2.40
b	20 14 18.3	+17 27 16	26					14.9	15.4	15.1		-1.08
c	20 14 19.2	+17 27 10	38					13.5	14.1	13.7		-1.64
d	20 14 19.5	+17 27 18	36					16.1	16.7	16.3		-0.60
086 a	19 48 49.0	+17 25 43	22	GSC 1619 3349	S11026 Sge	CA		11.7	13.2	12.3	14.2-15.0	-2.35
b	19 48 48.3	+17 26 12	9					11.7	13.2	12.3		-2.35
c	19 48 50.4	+17 25 56	27					15.4	16.5	15.8		-0.95
d	19 48 47.3	+17 26 27	30					15.5	16.3	15.8		-0.95
e	19 48 48.8	+17 26 36	31					13.1	15.2	13.9		-1.71
f	19 48 50.5	+17 25 41	35					14.8	16.3	15.4		-1.11
087	—	—	—	SNR 053.6-02.2		SNR						
088	—	—	—	SNR 053.6-02.2		SNR						
089	—	—	—	SNR 053.6-02.2		SNR						
090	—	—	—	SNR 053.6-02.2		SNR						

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R mag	B mag	V mag	Ampl. mag	$\log(f_x/f_y)$
	R.A.	Decl.	"									
091	—			SNR 053.6-02.2		SNR						
092 a	20 05 24.1	+17 10 05	17		C			13.8	14.8	13.9		-1.80
b	20 05 25.9	+17 10 01	17		C			14.7	16.1	15.0		-1.36
c	20 05 24.0	+17 10 19	16		C			14.1	16.3	14.9		-1.40
d	20 05 24.1	+17 10 24	18		C			15.9	16.5	16.1		-0.92
e	20 05 25.5	+17 10 13	5		S11035 Sge	CV?		17.4	17.8	17.6	16.6–>20.0	-0.32
093	—			SNR 053.6-02.2		SNR						
094	—			SNR 053.6-02.2		SNR						
095 a	20 03 29.8	+17 08 18	35		C			15.6	17.2	16.2		-0.86
b	20 03 34.6	+17 07 27	55		C			17.0	18.0	17.4		-0.38
096 a	20 09 27.3	+17 05 55	32		C			15.9	17.0	16.3		-0.87
b	20 09 30.0	+17 06 23	23					17.1	19.1	—		+0.02
097 a	20 04 06.5	+17 04 16	24	HD 190406	NVS 12757	CA?	G1V	5.9	6.5	5.9	0.03	-4.58
098 a	20 03 18.5	+17 02 52	13		RZ Sge	UGSU		17.2	16.7	16.8	12.2–17.4	-0.35
b	20 03 17.9	+17 02 26	15		C			14.7	15.5	15.0		-1.07
c	20 03 16.6	+17 03 09	35		C			15.1	16.0	15.4		-0.91
d	20 03 16.0	+17 02 39	24		C			15.2	16.1	15.5		-0.87
099 a	20 15 21.7	+16 58 22	20		C			13.8	13.9	13.8		-2.18
b	20 15 19.3	+16 58 37	38		C			15.0	15.3	15.1		-1.66
100 a	19 46 10.1	+16 50 56	20	HD 353578			M	10.4	12.2	11.1		-2.49
b	19 46 08.6	+16 51 25	22					15.2	16.7	15.8		-0.61
101 a	20 14 53.3	+16 46 24	8	HD 355199			F2	8.9	11.8	10.0		-2.92
b	20 14 55.3	+16 46 12	38					17.2	16.1	16.8		-0.20
c	20 14 51.5	+16 46 38	24					14.9	15.7	15.2		-0.84
102 a	19 41 57.1	+16 44 40	6	GSC 1602 1636	HM Sge	NC	M	11.2	12.8	11.1	11.1-18.0	-2.70
b	19 41 57.2	+16 44 29	13					13.9	14.7	14.2		-1.46
103 a	20 10 41.0	+16 42 29	13	HD 354910			K0	9.7	12.3	10.7		-3.06
b	20 10 40.0	+16 41 56	23					15.3	15.8	15.5		-1.14
104 a	19 59 57.5	+16 43 24	11	GSC 1616 1197	S11027 Sge	CA		12.1	14.0	12.8	14.0-14.5	-1.88
b	19 59 58.2	+16 43 29	19		C			14.6	16.1	15.2		-0.92
c	19 59 56.4	+16 43 05	21		C			13.9	16.0	14.7		-1.12
d	19 59 58.3	+16 42 55	21		C			14.7	16.6	15.4		-0.84
e	19 59 56.6	+16 43 33	24		C			16.0	16.7	16.3		-0.48
105 a	20 00 58.8	+16 36 24	15	HD 354463			K7	10.9	12.5	11.5		-2.78
b	20 01 00.4	+16 36 02	22					15.2	15.8	15.4		-1.22
c	20 01 00.7	+16 36 49	28					13.5	14.5	13.9		-1.82
106 a	20 18 37.0	+16 27 44	19	HD 355354			F5	10.9	12.6	11.5		-2.75
107 a	20 13 54.3	+16 25 25	3	HD 192362			K7	—	9.9	8.5		-3.61
108 a	20 04 40.5	+16 27 11	28	HD 354645	C		G5	10.8	12.1	11.3		-2.88
109 a	20 15 01.1	+16 22 31	28		C			16.7	17.4	17.0		-0.62
b	20 15 03.8	+16 22 27	30			AGN?		17.7	19.0	—		-0.05
110 a	19 56 41.5	+16 21 31	21		C			15.0	16.8	15.7		-1.02
b	19 56 39.0	+16 21 59	26		C			13.5	14.7	14.0		-1.70
c	19 56 40.5	+16 22 20	33		C			15.7	16.1	15.9		-0.94
d	19 56 43.4	+16 21 42	40		C			15.0	15.5	15.2		-1.22
e	19 56 39.6	+16 21 49	14					17.3	18.6	—		-0.09
111 a	19 56 06.3	+16 19 58	9		C			13.2	15.2	14.0		-1.76
b	19 56 06.4	+16 20 25	28	GSC1616 1465	C			12.0	12.7	12.3		-2.44
c	19 56 06.8	+16 19 18	40		C			12.8	15.1	13.7		-1.88
d	19 56 08.2	+16 19 42	24		C			15.5	16.8	16.0		-0.96
112 a	20 07 59.2	+16 09 58	6	HD 191179			G5	8.5	9.4	8.5		-2.36
b	20 07 59.6	+16 10 16	24	GSC 1617 992				12.0	13.3	13.4		-0.40
c	20 07 57.6	+16 10 15	34					12.8	15.9	14.0		-0.16
d	20 08 01.4	+16 09 33	35					14.1	14.4	14.2		-0.08
113 a	20 10 14.5	+16 08 37	26		S11028 Aql	CA?		15.3	16.7	15.8	15.6–16.7	-1.10
b	20 10 16.0	+16 08 46	41		C			14.5	16.1	15.1		-1.38
114 a	20 17 23.9	+16 03 21	23		C			15.4	17.3	16.1		+0.27
b	20 17 25.7	+16 03 11	9		C			15.2	15.8	15.4		-0.01
c	20 17 25.8	+16 03 36	10		C?			16.4	16.6	16.8		+0.55
d	20 17 25.8	+16 03 52	34		C			13.1	14.6	13.7		-0.69

Table 5b. continued.

No	Coordinates (2000.0)		D	Designation	Name if variable	Type	Spec.	R	B	V	Ampl.	log(f_x/f_y)
	R.A	Decl.	"					mag	mag	mag	mag	
e	20 17 26.8	+16 02 51	34		C			14.1	14.6	14.3		-0.45
f	20 17 23.7	+16 03 46	37		S11029 Aql	CV?		15.5	16.3	15.8	15.8–17.3	+0.15
115 a	19 54 17.6	+16 00 12	33	HD 354137	C		F8	8.9	12.0	10.1		-3.35
b	19 54 15.6	+16 00 22	31		S11030 Aql	CV?		14.5	14.6	14.5	13.3–14.1	-1.59
c	19 54 14.8	+16 00 14	27					16.7	16.5	16.6		-0.75
d	19 54 13.5	+16 00 11	38					15.6	16.5	15.9		-1.03
116 a	20 09 51.4	+15 57 34	7	GSC 1618 1655	S10947 Aql	RS		12.0	13.5	12.6	13.1–14.0	-2.27
b	20 09 53.9	+15 57 07	40	GSC 1618 1609				12.7	13.2	12.9		-2.15
c	20 09 49.4	+15 57 34	36					13.4	14.6	13.9		-1.75
117 a	20 18 44.4	+15 50 47	23	GSC 1631 97	C		M2.5	10.9	13.1	11.3		-2.93
b	20 18 43.4	+15 50 50	36		C			13.0	14.6	14.1		-1.81
c	20 18 44.4	+15 51 06	24		C			15.9	16.2	16.0		-1.05
d	20 18 43.8	+15 50 23	43	GSC 1631 497	V?			12.9	13.1	13.0		-2.25
118 a	20 04 46.8	+15 54 11	19	HD 190516			G0V	7.3	8.0	7.4		-4.43
119 a	19 46 13.6	+15 48 59	20	GSC 1615 2844	S11037 Aql	CA		11.9	13.0	12.3	12.7-13.2	-2.25
b	19 46 11.5	+15 48 29	35	GSC 1615 1836	C			11.9	13.7	12.7		-2.09
c	19 46 10.5	+15 49 12	27		C			14.3	14.9	14.5		-1.37
d	19 46 12.4	+15 49 01	3		C			15.8	16.4	16.0		-0.77
e	19 46 10.0	+15 49 19	36		C			15.5	16.6	15.9		-0.81
120 a	20 13 58.8	+15 44 28	19		S11038 Aql	CV?		17.3	17.2	17.3	16.5–18.0	-0.27
b	20 13 57.8	+15 44 56	24		C			14.1	15.4	14.6		-1.35
121 a	19 58 36.6	+15 43 56	18	HD 189281			G9	8.7	9.8	9.1		-2.60
b	19 58 37.5	+15 43 24	17					14.0	15.9	14.7		-0.36
c	19 58 36.6	+15 43 12	29					14.0	14.5	14.2		-0.56
d	19 58 37.7	+15 43 12	29					14.4	15.1	14.7		-0.36
122 a	20 08 27.0	+15 40 30	6	HD 191262	V1423 Aql	RS	G5	8.0	8.5	8.0	< ±0.03	-3.33
b	20 08 24.9	+15 40 23	26					13.6	15.9	14.5		-0.73
c	20 08 24.3	+15 40 28	34					14.7	15.3	14.9		-0.57
123 a	19 43 10.4	+15 28 29	25		S11031 Aql	CA?		14.6	15.2	14.8	14.4-15.0	-1.65
b	19 43 12.2	+15 28 51	11		C			15.9	17.1	16.4		-1.01
c	19 43 12.5	+15 28 23	22		C			16.4	17.4	16.8		-0.85
d	19 43 11.6	+15 28 36	6					16.8	18.9	18.0		-0.37
e	19 43 11.3	+15 28 33	11			AGN?		18.1	18.6	18.4		-0.21
124 a	19 42 50.1	+15 22 56	23	HD 353475	C?		K2	9.1	11.9	10.2		-3.40
b	19 42 47.9	+15 22 28	31					13.9	15.9	14.7		-1.60
c	19 42 51.7	+15 22 35	24					15.4	17.1	16.0		-1.08
d	19 42 52.2	+15 22 48	35	GSC 1602 335				11.9	14.1	12.7		-2.40
e	19 42 49.2	+15 22 03	32					14.4	16.7	15.3		-1.36
125 a	19 54 02.0	+15 17 32	6	HD 188328			F8III	7.3	7.6	7.0		-4.23
126 a	20 18 54.9	+15 07 10	8		C	AGN?		16.0	16.7	16.3		+0.34
b	20 18 56.2	+15 07 28	31		S11032 Aql	RS+E?		14.6	15.5	14.9	15.4-16.0	-0.22
c	20 18 55.3	+15 07 41	31		C			15.3	16.6	15.8		+0.14
127 a	20 17 05.9	+15 02 45	14		C			14.2	15.4	14.7		-1.35
b	20 17 05.0	+15 02 41	8		V?			17.2	17.8	17.4		-0.27
c	20 17 08.0	+15 02 28	40	GSC 1631 1345	C			11.7	12.9	12.2		-2.35
d	20 17 03.4	+15 55 40	28		C			15.6	15.7	15.6		-0.99
128 a	19 43 02.1	+14 55 40	18		C			15.0	15.8	15.3		-1.19
b	19 43 01.3	+14 55 33	29		C			15.5	16.7	16.0		-0.91
c	19 43 04.9	+14 56 12	41					12.0	13.9	–		-1.98
129 a	20 14 55.8	+14 51 00	6	HD 355274	C		K0	10.7	10.9	10.8		-3.05
130 a	20 19 24.9	+14 36 57	1		C			15.8	16.5	16.1		-0.73
b	20 19 24.6	+14 37 00	22		S11033 Aql	CV?		17.2	17.6	17.4	16.6-17.5	-0.21
c	20 19 24.6	+14 37 16	18					16.2	16.8	16.4		-0.61
131 a	19 52 54.9	+14 36 08	12	GSC 1070 1591	S11034 Aql	CA		11.1	10.9	11.0	11.1-11.4	-2.57
132 a	20 10 12.8	+14 32 05	11		C			15.6	16.2	15.8		-0.97
b	20 10 13.7	+14 31 57	18					18.6	19.2	–		+0.16

3.4. Notes to individual objects in the Com Field (Table 5a)

1a: On one Tautenburg Schmidt plate (1944 Apr. 6) and two overlapping Palomar prints (1950 Apr. 10 and 1955 May 21) the object is visible with nearly equal brightness.

2a: The large distance from the ROSAT source indicates that the X-ray source may be near the periphery of the galaxy NGC 4565. Chandra and XMM positions confirm this. Because of its large luminosity this source was classified as an ultra-luminous X-ray source (ULX-4; Wu et al. 2002). Thus, the identification is not with the galaxy, but with the ULX in that galaxy. The B magnitude is for the whole galaxy. Since the optical counterpart of the X-ray source is not known, no f_X/f_{opt} is given. See also Brinkmann et al. (1995).

3a: Because this faint object is near the border of the astrograph plates, it is faintly indicated only on some dozen plates. Variable within some weeks. Accidentally, this object is visible on 3 overlapping fields of the POSS (1955 Apr. 15 17^m3, 1955 May 21 17^m0, 1956 May 17 16^m6) and on 4 plates of the 2 m Schmidt telescope of Tautenburg, which shows the object at different brightnesses. A low-S/N, not flux-calibrated optical spectrum (courtesy S. Zharykov, ONAM Ensenada, Mexico) reveals this object to be an AGN at $z = 0.195$ (see Fig. 5), consistent with the very soft X-ray spectrum. A 2-hr V-band monitoring on March 14/15 1996 (23:58–2:05 UT) shows variability by 0.3 mag. The SDSS-III spectrum identifies it as QSO at $z = 0.1957$ and a “starburst broadline” sub-class.




Fig. 5. Not flux-calibrated optical spectrum of object Com003, showing broad Balmer lines, and narrow [O III], suggesting a narrow-line Sy1 ($z=0.195$). Courtesy S. Zharykov, ONAM Ensenada.

6a: HR 4707 $\equiv$ 12 Com.

7a: Very soft X-ray spectrum. A low-S/N, not flux-calibrated optical spectrum (courtesy S. Zharykov, ONAM Ensenada, Mexico) reveals only one emission line at 465 nm, so it is likely an AGN. The SDSS-III spectrum identifies it as broadline QSO at $z = 0.6680$.

8a: Faint on two Tautenburg plates (3999 from J.D. 2442094 and 8586 from J.D. 244 9449). This variability, the f_X/f_{opt} ratio and the blue colour suggest an AGN or CV nature, rather than a stellar member of the Coma Berenices open cluster as proposed by Randich et al. (1996). The SDSS-III spectrum identifies it as a AGN at $z = 0.0668$.

9a: Fainter companion is totally blended.

10a: On POSS print and one Tautenburg Schmidt plate of equal brightness. CV Com is 4' outside the ROSAT position.

The SDSS-III spectrum identifies it as a QSO at $z = 0.1598$, with “starburst broadline” sub-class.

11a,b: Investigation on Sonneberg astrograph plates not possible due to blending. The Swift XRT observation did not detect this source. The upper limit of <0.002 cts/s is about a factor 5 below the ROSAT flux.

12a: Probable active galaxy. The published USNO-A2 B magnitude of 17.2 is substantially different from that in USNO-B which gives $B1 = 16.11$ mag. This object is too faint for Sonneberg plates. The red colour of the galaxy does not suggest an AGN nature, and the f_X/f_{opt} is too high for normal, inactive galaxies. Besides the all-sky survey, this sky area was observed at two more occasions with the ROSAT PSPC: once for 12.01 ksec between Dec. 17, 1991 and Jan. 6, 1992, and again for 7.35 ksec on Jun 4, 1992. At the first occasion, the measured count rate is 0.084 ± 0.004 cts/s while at the second it is 0.067 ± 0.004 cts/s, thus this source is clearly variable when compared to the all-sky survey rate of 0.045 ± 0.010 cts/s. The Swift XRT position confirms the association of the X-ray source to this galaxy. Considering the soft spectrum, the XRT count rate is similar to the intensity during the ROSAT all-sky survey. This X-ray intensity pattern, in particular the rise between 1990 and 1992, is rather slow compared to what one would expect for a tidal disruption event (e.g. Komossa & Greiner 1999). The SDSS-III spectrum reveals broad emission lines, and identifies it as QSO at $z=0.1417$.

13a: UVB photometry by Zeilik et al. (1982). See also Fleming et al. (1989).

14b: Is quasar 4C 25.39 $\equiv$ PKS J1217+2529, and is a blue object. The f_X/f_{opt} ratio favours the identification of the ROSAT source with this quasar, rather than the brighter star 14a. The SDSS-III spectrum reveals broad emission lines, and identifies it as QSO at $z=0.6789$.

15a: Triple blend on Sonneberg plates. The $B - V$ colour and f_X/f_{opt} are consistent with a K or M spectral type and the object being the optical counterpart of the X-ray source. This is verified by the Swift XRT data. **15b:** Object is too faint for Sonneberg plates. On POSS (243 5550) about 19.0 mag, and on 2 Tautenburg plates (244 1400 and 244 2074) about 19.5 mag. Nevertheless, variability is uncertain. The blue colour suggests an AGN nature (the X-ray hardness ratio argues against a white dwarf nature).

16a: HD111813 is the obvious counterpart, which is also confirmed by the XMM position (2XMMp J125138.4+253033; it is plotted with larger uncertainty for visibility reasons in Fig. 40a). The two pointings mentioned under Com012a also cover this source, and the corresponding X-ray count rates are 0.040 ± 0.002 and 0.035 ± 0.003 , respectively.

17a: The ROSAT position coincides with the centre of the Sy2 galaxy NGC 4725, but XMM resolves this into 2 sources (2XMMp J125026.6+253003 and 2XMMp J125027.3+253026), and in addition finds three more nearby sources which may add to the ROSAT source (2XMMp J125024.2+252947, 2XMMp J125024.3+252941, and 2XMMp J125024.9+253053).

18a: Flare star GJ 3739. No X-ray flares observed. $B - V$ colour and f_X/f_{opt} are consistent with coronal emission from an M star.

19a: Probably galaxy pair; if interacting, or with one active member, it could explain the X-ray emission. The two pointings mentioned under Com012a also cover this source, and the corresponding X-ray count rates are 0.086 ± 0.006 and 0.078 ± 0.007 , respectively. Both rates are about a factor 2 higher than the rate seen during the all-sky survey.

20a: Invisible on Sonneberg plates. The red colour and the fuzzy appearance on the digitized sky survey suggest a galaxy. This is supported by the SDSS-III spectrum which identifies this object as galaxy at $z=0.1834$, though Kouzuma et al. (2010) identify it as AGN candidate due to the NIR colours from 2MASS. However, the X-ray hardness ratio hints more towards a cluster identity. The similar colours and fuzzy appearance of several nearby objects, most notably the other two sources North and north-west of 20a within the RASS error circle, suggest galaxies at the same redshift.

21a: Invisible on Sonneberg plates. The blue colour and large f_X/f_{opt} ratio suggests an AGN nature, in which case it could be the counterpart.

22a: BL Lac object at $z=0.135$; see Brinkmann et al. (1995) and Sowards-Emmerd et al. (2005). Not tested for variability; on POSS (243 5249) and on Tautenburg plate (244 9449) of equal brightness.

23a: The variability is not sure. Probably constant. If the FG spectral type is correct, then the f_X/f_{opt} ratio argues against this object being the counterpart of the X-ray source.

24a: On Tautenburg plate 8586 (TJD = 24 49449) invisible. The SDSS-III spectrum identifies it as QSO at $z=0.3408$ with “starburst broadline” sub-class.

25: The Swift XRT position favours object 25b; being too faint for the Sonneberg plates. It is listed as SDSS J124640.80+251149.5 and type “Galaxy” at $z=0.084$ in Rines et al. (2001), with the SDSS-III giving an identification as star-forming galaxy. The Swift XRT intensity is about the same as that seen with ROSAT.

26a: *Einstein* X-ray source (Stocke et al. 1983). Not tested for variability. The SDSS-III spectrum identifies it as QSO at $z=0.0637$ with “starburst broadline” sub-class.

27a: Very difficult because near the brightness limit and near the edge of the plate, but the variability seems to be sure. Brightness varies irregularly. The SDSS-III spectrum identifies 27a as broadline QSO at $z=0.5447$, and 27b as star.




Fig. 6. Blow-up of the light curve of Com027a (Fig. 41a) showing pronounced variability.

28a: Not visible on Sonneberg plates. Blue colour suggests an AGN nature (see also Chen et al. 2002). The Swift XRT observation did not detect this source; the upper limit is not constraining, given the soft X-ray spectrum.

29a: Invisible on Sonneberg plates. Due to the blue colour most likely a QSO, which is confirmed by the SDSS-III spectrum (broadline QSO at $z=0.5865$). 29b is identified by Kouzuma et al. (2010) as AGN candidate due to the NIR colours from 2MASS.

30a: Seyfert 1.5 galaxy (TON 0616, 4C 1834, FBQS J122539.5+245836) at $z=0.268$. On Tautenburg plates (244 2094 and 244 9449) faint. See also the optical catalogue of QSOs by Hewitt & Burbidge (1987), and Brinkmann et al. (1995). The

SDSS-III spectrum classifies it as QSO at $z=0.2679$ with “starburst broadline” sub-class.

31: Nothing visible on Sonneberg plates. The Swift XRT position suggests 31a as counterpart.

32a: Within the error circle is one radio source identified during the FIRST bright quasar survey: FBQS J1249+2452 = FIRST J124958.8+245233 (White et al. 2000). This coincides within 1'' with object 32a which we therefore classify as an AGN. The SDSS-III spectrum supports this, and gives $z=0.2465$.

33: No optical counterpart candidate brighter than $B = 19.5$ mag within the ROSAT error circle. The Swift XRT observation reveals clearly extended emission, centred around $12^{\text{h}}45^{\text{m}}15^{\text{s}}.0 + 24^{\circ}53'35''$ and with a diameter of about 1'.5. The ROSAT and Swift X-ray fluxes are comparable. The hard spectrum and the diffuse emission suggest a cluster origin. The ROSAT source is likely the detection of a brighter blob at the south-eastern rim of the extended emission.

34a: M star. Eclipse minima of short duration (1 day or shorter) seem to exist, but because of irregularities in the light curve no secure period could be found. A period length of 0.35 days seems to be indicated, but there are contradictions with bright states. Very probably chromospherically active star of RS CVn type. Observed minima: JD 2438 083.64 (15.3 mag; descent, Fig. 7), 088.47 (15.4; ascent, Fig. 7), 089.54 (15.2), 090.57 (15.2), 091.63 (15.2), 093.66 (15.3), 116.38 (15.3), 140.45 (15.3), 142.35 (15.3), 146.51 (15.3), 473.32 (15.3), 817.66 (15.2), 2445 812.51 (15.5:), 2447 205.55 (15.3:), 262.47 (15.4), 2448 682.51 (15.3).




Fig. 7. Blow-up of the light curve of Com034a (Fig. 41a) showing pronounced variability.

35a: Invisible on Sonneberg plates. Due to the blue colour most likely a QSO. The SDSS-III spectrum classifies it as QSO at $z=0.2184$ with “starburst broadline” sub-class.

36a: Other name HD 111 395.

37b: Blue object, thus possibly AGN; but 37a is variable and is therefore the better candidate for the optical counterpart of the ROSAT source. This is confirmed with the Swift/XRT position. The optical light curve shows waves with a length of about 20–30 days. May be a BY Dra star or a massive X-ray binary.

38a: Classified as QSO in Zickgraf et al. (2003); too faint on Sonneberg plates for variability assessment. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5471$.

39a: Classified as M star in Zickgraf et al. (2003); $B - V$ colour and f_X/f_{opt} are consistent with this. Most of the time at a bright (Fig. 9, top panel), sometimes at mean magnitude level (Fig. 9, left part of bottom panel), with only small variations. Sometimes fading with a duration of some days or several weeks (Fig. 9, right part of bottom panel).

40a: See also Brinkmann et al. (1995) for the BL Lac classification and redshift $z=0.218$. On Sonneberg plates invisible. On POSS print 135 (J.D. 243 3291) at $B \sim 16.6$ mag. On Tautenburg




Fig. 8. Blow-up of the light curve of Com037a (Fig. 41a) showing pronounced variability.




Fig. 9. Blow-up of the light curve of Com039a (Fig. 41a) showing pronounced variability.

Schmidt plates 3999 (J.D. 244 2094) $B = 16.6$ mag, and 8586 (J.D. 244 9449) $B = 17.8$ mag. Therefore variable. The SDSS-III spectrum classifies it as galaxy at $z=0.2187$, despite its very blue continuum.

41a: Faint object, on only 4 astrograph plates just visible. May therefore be sometimes brighter than 18 mag and probably variable. The blue colour and f_X/f_{opt} ratio suggest an AGN nature.

42a: This object has been spectroscopically identified as QSO at $z=0.438$ (Chen et al. 2002). This very blue object is just visible on the 6 best plates. Nothing can be said about variability. Object 42b also has a blue colour, suggesting also an AGN nature. The SDSS-III spectrum classifies 42b as broadline QSO at $z=1.2078$. The RASS position suggests that 42a is the optical counterpart of the X-ray source.

43a: Other name BD+25°2511. See also Raveendran (1984).

44a: Invisible on Sonneberg plates. The blue colour suggest an AGN nature. The SDSS-III spectrum classifies it as broadline QSO at $z=0.3710$.

45a: Already detected by the *Einstein* Observatory (Stocke et al. 1983). Narrow-line Sy1 galaxy at $z=0.186$, with ROSAT data and optical variability based on Sonneberg data already reported earlier (Greiner et al. 1996). The SDSS-III spectrum classifies it as QSO at $z=0.1856$ with “starburst broadline” sub-class.

46: Both objects invisible on Sonneberg plates. The Swift XRT position points to 46a as the counterpart, and the blue colour and f_X/f_{opt} ratio suggest an AGN nature.

47a: Within the error circle is one radio source, FBQS J122214.6+242420 at a spectroscopic distance of $z=0.12$ (White et al. 2000). This coincides within 1'' with object 47a which

we therefore classify as an AGN. On Tautenburg plates invisible (>18 mag). The Swift XRT observation finds this source at twice the ROSAT rate, and confirms this association through an accurate X-ray position. The SDSS-III spectrum classifies it as QSO at $z=0.1167$ with “AGN broadline” sub-class.

48a: Not tested for variability. The blue colour suggest an AGN nature. The SDSS-III spectrum classifies it as broadline QSO at $z=0.8282$.

49a,b,c: If the K spectral type for 49c is correct, then it is excluded as counterpart. 49a,b are possible counterpart candidates, with 49a having higher likelihood. The Swift XRT observation detects 3 photons from this source 49a; this low intensity is consistent with the moderately soft X-ray spectrum. The SDSS-III spectrum classifies 49a as broadline QSO at $z=1.0444$.

50: No optical counterpart candidate brighter than about $B = 20$ mag inside the ROSAT error circle. The Swift XRT observation suggests diffuse X-ray emission, with an extent of about 20-30'' (likely biased due to counting statistics). Also, the ROSAT and the Swift XRT X-ray flux measurements are the same.

51a: Only seen on the 8 best plates. On POSS print 1435 (J.D. 243 5249) about 1 mag fainter than on Sonneberg plates as can be seen in Figs. 10 & 41a. The SDSS-III spectrum classifies it as broadline QSO at $z=0.465$.




Fig. 10. Blow-up of the light curve of Com051a (Fig. 41a) showing pronounced variability.

52b: Already detected by the *Einstein* Observatory (Stocke et al. 1983). BL Lac object at $z=0.14$, with ROSAT data and optical variability based on Sonneberg data already reported earlier (Greiner et al. 1996). XMM detects three X-ray sources in the field (see Fig. 40a), one of which (2XMMp J125731.9+241240) coincides with 52b, thus confirming the earlier identification. The SDSS-III spectrum classifies 52b as galaxy at $z=0.1406$.

53a: Slightly outside the ROSAT error circle, and too faint for variability analysis on Sonneberg plates, but the XMM position (2XMMp J125638.7+241252) confirms the identification.

54a: Not tested for variability due to faintness. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5071$.

55a: Blue colour suggests an AGN nature. Not tested for variability, since too faint on Sonneberg plates. On Tautenburg plate 3999 (244 2094) about 18.0 mag. The SDSS-III spectrum classifies it as broadline QSO at $z=0.3795$.

56a: Spectroscopically classified as Seyfert galaxy at $z=0.188$ by Chen et al. (2002). Difficult because of faintness. At the beginning of the observations (J.D. 243 7696 - 7822) rather quick brightness changes between 16.4 and 17.2 mag, and the whole amplitude can be run through within 1 day. Short weakenings from a generally bright level in irregular time intervals. From J.D. 243 8085 to 8549 the brightness varies between 16.8 and >17.5 mag. Faint observations preponderate, so that there are relatively short maxima. From J.D. 243 8817 to 8902 the object is obviously always bright between 16.4 mag and 16.6 mag

(see Fig. 11). The few observations from 243 9144 - 244 2540 show the object between 16.6 mag and 17.0 mag. Since J.D. 244 4292 the object is nearly always invisible. An observation of $B = 17.2$ mag at 244 5818 must already be considered as a brightening. On the blue print of the Palomar survey (JD 243 5550) about 18 mag (Figs. 11 & 41a). The SDSS-III spectrum classifies it as QSO at $z=0.1884$, with “starburst broadline” sub-class. The RR Lyr star EL Com has a distance of about $210''$ to the ROSAT position and is not considered a counterpart candidate.




Fig. 11. Blow-up of the light curve of Com056a (Fig. 41a) showing pronounced variability.

57a: Not tested for variability.

58a: Not investigated on Sonneberg plates.

59: Cluster of galaxies NSC J122158+235426 at $z=0.149$ (Gal et al. 2003). 59a: Irregular galaxy 2MASX J12215975+2354425, seemingly not active. The SDSS-III spectrum classifies 59a as galaxy at $z=0.1318$, while Kouzuma et al. (2010) identify it as AGN candidate due to the NIR colours from 2MASS. The hard X-ray spectrum (hardness ratio) suggests that the cluster is the counterpart, not the galaxy 59a.

60: No optical counterpart candidate brighter than $B = 21$ mag in the ROSAT error circle. The Swift XRT observation did not detect this source; the upper limit is not constraining, given the very soft X-ray spectrum. The brightest SDSS (DR9) objects are three galaxies with $g'=22.4$ at $23''$ offset, $g'=22.6$ mag at $25''$ offset, and $g'=21.7$ at $31''$ offset. The latter implies a lower limit on $\log(f_X/f_{\text{opt}}) \gtrsim +1.2$.

61a,b: Both objects have blue colour, and are likely AGN, see also Micaelian et al. (2006). The Swift XRT observation suggests 61a as the counterpart. The SDSS-III spectrum classifies 61a as galaxy at $z=0.3984$.

62a-e: Several faint galaxies within the error circle; not tested for variability. Remarkable is the high X-ray intensity. A ROSAT HRI observation indicates that the X-ray source is possibly extended (Gliozzi et al. 1999), and thus the cluster of galaxies at $z=0.134$ could be the counterpart. However, the X-ray flux is centred on the quasar 62e, and, more importantly, the flux seen in the HRI is a factor 4 lower than that from the survey, suggesting that the quasar dominantly contributes. 62a is a galaxy (NGP9 F378-0239966) at $z=0.132$ (Gliozzi et al. 1999) which coincides with the radio source FIRST J123438.6+235013, thus probably is an AGN. 62c is a galaxy (NGP9 F378-0239961) at $z=0.135$ (Gliozzi et al. 1999) which coincides with the radio source FIRST J123438.6+235013, thus probably also is an AGN. 62d is a galaxy (NGP9 F378-0239941) at $z=0.133$ (Gliozzi et al. 1999) which coincides with the radio source FIRST J123437.1+235016, thus probably also is an AGN. 62e is a quasar at previously unknown redshift (Bade et al. 1998), for which the SDSS-III spectrum provides $z=0.1382$ together with a classification as broadline QSO. Based on the X-ray variability, the hardness ratio and f_X/f_{opt} we identify 62e as the counterpart.

63a: Potentially a member of the Coma Berenices open cluster (Randich et al. 1996).

64a: This object is about 21 mag on the Palomar print. It is not contained in the USNO catalogues, just in the APM (McMahon et al. 2000) from which the magnitudes are taken. It is invisible on all Sonneberg plates with the exception of J.D. 243 8085.459 ($B = 15.5$ mag); on the contrary, at J.D. 243 8085.501 it is invisible (>17 mag), implying a decline of >1.5 mag within 1 hr. Reflected light microscopy (see e.g. Greiner et al. 1990 for a similar application) gives no indication for a plate fault. Based on this, the object may be a flare star. Nevertheless, the optical variability remains to be confirmed. The arrival times of the 13 X-ray photons are not compatible with a single flare, but seem to be distributed equally.

65: Red star of 20 mag at a distance of $19''$. Much too faint for Sonneberg plates. The Swift XRT observation did not detect this source; the upper limit is not constraining, given the moderately soft X-ray spectrum. The SDSS-III spectrum classifies 65a as AGN at $z=0.1377$.

66a: Within a cluster of galaxies. 66b is the heavily disturbed galaxy pair IC 3314+IC3312, which therefore is probably the optical counterpart of the ROSAT source. The Swift XRT observation did not detect a point source; the upper limit suggests a factor of a few fading relative to the ROSAT measurement.

67a: Stellar object with UV excess (Green et al. 1986). The very soft X-ray spectrum ($\text{HR1}=-1.0$) clearly points towards the white dwarf PG 1232+238 as the counterpart, consistent with the identification by Zickgraf et al. (2003).

68a: The blue colour suggests an AGN nature, and the the SDSS-III spectrum provides $z=0.2583$ together with a classification as broadline QSO.

69a/b: Close pair of stars of equal brightness; the eastern component 69a is variable, and is identical with S 10937 $\equiv$ KX Com (Richter et al. 1995).

70: The SDSS-III spectrum classifies 70a as starforming galaxy at $z=0.1947$, but the f_X/f_{opt} ratio suggests the QSO 70b as the counterpart, for which the SDSS-III spectrum provides $z=1.1775$.

71: Some galaxies fainter than 21 mag are within the ROSAT error circle, but too faint for Sonneberg plates. The hard X-ray spectrum might suggest a cluster identity.

72: While 72a is a galaxy based on its colours and too faint for Sonneberg plates, 72b is an A0 star according to SDSS-III, and difficult on Sonneberg plates because of faintness. Due to the inconsistent f_X/f_{opt} ratio, the A0 star cannot be the counterpart.

73a: On 4 Tautenburg Schmidt plates and on POSS of equal brightness. The f_X/f_{opt} ratio excludes a stellar counterpart, but rather suggests a CV or AGN origin. However, the optical colours are not particularly blue.

74: Nothing visible on Sonneberg plates. The Swift XRT position identifies 74a as counterpart. The blue colour and large f_X/f_{opt} suggest a AGN or CV identification. Given the moderately soft X-ray spectrum, the ROSAT and Swift intensities are comparable. The SDSS-III spectrum classifies 74a as QSO at $z=0.2728$, with “starburst broadline” sub-class.

75a: Considering the faintness of all optical objects in question, the high X-ray intensity is remarkable. Possibly an optically faint X-ray binary or an X-ray bright AGN, based on f_X/f_{opt} . Zickgraf et al. (2003) report “BLUE-WK”, a moderately blue continuum, weak point-like object. The SDSS-III spectrum classifies it as broadline QSO at $z=0.1606$.

76a: Because of faintness variability not quite sure. The Tautenburg plate 3999 shows the object at 17.5 mag. 76a and 76b are a narrow pair.

77a: Because of faintness nothing can be said about variability. The blue colour suggests an AGN (or CV) nature which would be consistent with the f_X/f_{opt} ratio. The SDSS-III spectrum classifies it as broadline QSO at $z=0.6667$.

78b: Because of faintness only on a part of plates visible. Seemingly there are long flat waves (slowly oscillating brightness variations) with a timescale of some 100 days, standstills seem also to occur. During the intervals J.D. 243 7695 - 7850, 8415 - 8550, 8815 - 9100 about $B = 17.5$ mag. Thereafter $B = 17.8$ mag and fainter, but observations are only sporadic. J.D. 244 4700 again brighter, thereafter weakening. On one plate of Tautenburg Schmidt telescope at 244 2094 about $B = 18$ mag. The SDSS-III spectrum classifies it as broadline QSO at $z=1.3866$.

79: Both objects too faint for Sonneberg plates. The SDSS-III spectrum classifies 79a as broadline QSO at $z=0.3412$.

81a: At first glance seems to be variable between about 17.5 and 18.5 mag. But in spite of the seemingly large amplitude, the variability is not quite sure because of being near the plate limit. The SDSS-III spectrum classifies it as broadline QSO at $z=0.9669$. 81b: Though beyond the plate limit, marginally visible on some Sonneberg plates. Nevertheless, variability questionable.

82: No object brighter than 21 mag in the X-ray error circle. The SDSS-III spectrum classifies 82a as broadline QSO at $z=0.7227$ which we consider to be the counterpart despite its somewhat large distance to the X-ray centroid position.

83a: This faint object seems to be barely visible on some plates, but this may be artefacts. Nevertheless, a true variability cannot be fully excluded. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5299$.

84a: On POSS 1435 (1956 May 20) about $B = 19$ mag, on Tautenburg Schmidt plate 4005 (1974 Feb. 15) about $B = 17.5$: mag. Seems to be faintly indicated on some Sonneberg plates. Nevertheless, optical variability is not quite sure. The Swift/XRT observation reveals no detection, with an upper limit about a factor 4 below the RASS rate, so clearly X-ray variable. This and the f_X/f_{opt} ratio suggest an AGN nature. The SDSS-III spectrum classifies it as QSO at $z=0.2033$, with "starburst broadline" sub-class.

85: No optical counterpart candidate brighter than about $B = 20$ mag inside the ROSAT position. The brightest SDSS (DR9) objects are a galaxy ($g'=21.0$ mag, $35''$ offset) and a star ($g'=21.35$, $26''$ offset). The former implies a lower limit on $\log(f_X/f_{\text{opt}}) \gtrsim +1.1$.

86a,b: Both objects are outside the error circle. Inside the error circle no object brighter than 21 mag. 86b is of equal brightness on POSS (243 5249) and the Tautenburg plate (244 2094).

87a: Variable with a time scale of several days to months: possibly chromospherically active star. Spectral classification from Garcia Lopez et al. (2000) who discard Coma Berenices cluster membership which had been proposed by Randich et al. (1996), but find spectroscopic evidence for chromospheric activity.

88a: KZ Com = S 10939. See Richter et al. (1995).

89a+b: The SDSS resolves this into 4 objects, and all objects are consistent with the ROSAT X-ray position. The f_X/f_{opt} ratio does not indicate an AGN nature for any of those, although Kouzuma et al. (2010) identify both 89a+b: as AGN candidates due to the NIR colours from 2MASS. The hard X-ray spectrum argues against a tidal disruption event in an inactive galaxy. The SDSS-III spectrum classifies 89b as galaxy at $z=0.1405$. A Swift/XRT observation does not reveal an obvious point source. There is, however, a blob of diffuse emission slightly west of




Fig. 12. Blow-up of the light curve of Com087a (Fig. 41a) showing pronounced variability.

89a+b, and a further blob of diffuse emission is about $2'$ to the west. We thus propose a galaxy cluster identification.

90a: Seems to be a spiral galaxy, for which the SDSS-III spectrum provides a classification as starforming galaxy, at $z=0.0760$. While this could be the counterpart of the ROSAT X-ray emission, we note that (i) there is no point source in a Swift/XRT pointing (the one centred on 89), but (ii) the ROSAT position falls in the middle of two Swift/XRT sources at $\text{RA}(2000.0) = 12\ 18\ 25.6$, $\text{Decl.}(2000.0) = +22\ 50\ 22.5$ and $\text{RA}(2000.0) = 12\ 17\ 42.1$, $\text{Decl.}(2000.0) = +22\ 48\ 43.7$, thus raising the possibility that 89 and 90 form part of a cluster about $7'$ in diameter.

91: No optical counterpart candidate brighter than $B = 21$ mag in the ROSAT error circle. The Swift/XRT observation does not reveal the source, with an upper limit a factor 5 below the RASS rate.

92a: The SDSS-III spectrum classifies it as broadline QSO at $z=0.13724$.

93a: Observed minima: J.D. 243 7783.516 (16.2 mag), 7822.417 (16.3: mag, ascent), 8440.654 (16.2 mag, descent), 244 5021.546 (16.3 mag) (Fig. 13).




Fig. 13. Blow-up of the light curve of Com093a (Fig. 41a) showing pronounced variability.

94a: KW Com = S 10936. See Richter et al. (1995).

95a: Other name: NGP9 F378-0391299 (Odewahn et al. 1995). Brightest member of a cluster of galaxies. The Swift XRT observation reveals a point source, so the X-ray emission is unlikely from the cluster. The SDSS-III spectrum classifies it as broadline QSO at $z=0.1603$.

96a: Spectral classification from Garcia Lopez et al. (2000) who discard Coma Berenices cluster membership, which had been proposed by Randich et al. (1996), but find spectroscopic evidence for chromospheric activity.

97a,b: Invisible on Sonneberg plates. The SDSS-III spectrum classifies 97a as broadline QSO at $z=0.5538$, and 97b as F5 star.

98a: Because of the large distance to the ROSAT source, it is unlikely that this is the optical counterpart, despite its

optical variability. From a bright normal light we have short (about two hours) minima (in parentheses B magnitude): J.D. 243 7795.474 (16.1), 244 2452.491 (16.3), 2480.524 (16.1), 2887.358 (16.2), 4701.522 (16.2), 6173.434 (16.2), 6827.618 (16.1), 6876.473 (16.2), 7206.516 (16.1), 9445.386 (16.2). Strikingly, from 1962–1975 only one minimum was observed, but thereafter eleven minima until 1994. The duration of the minima is about 0.05 days. *98b*: This object coincides with a strong radio source (NVSS 122401+223939; Condon et al. 1998), and has therefore been proposed as a BL Lac candidate (Sowards-Emmerd et al. (2005). Thus, it is the more likely counterpart of the X-ray source. The SDSS-III spectrum classifies it (with a small χ^2 warning) as galaxy at $z=0.4821$.

99a: Not tested for variability; too bright for astrograph plates. The very soft X-ray spectrum and the f_X/f_{opt} ratio are consistent with a G star counterpart, despite the relatively large distance between optical and X-ray position.

100a: Invisible on Sonneberg plates. Spectroscopically identified as Seyfert galaxy at $z=0.086$ by Chen et al. (2002).

101a: Clearly visible only on some plates. Mostly fainter than 18 mag. Gets as bright as 17.3 mag between J.D. 243 7696 and 7820 and between 243 9557 and 9609. On Palomar print 1435 (1956 May 21) about $B = 18.5$ mag. On Tautenburg Schmidt plates from 1974 Feb. 15 and 1991 Apr. 9 both about $B = 17.9$ mag. The Swift XRT position clearly favours *101b* as counterpart which is also a radio source (FIRST J125232.6+223338). The SDSS-III spectrum classifies it as broadline QSO, at $z=0.2113$.

102: Both optical objects too faint for Sonneberg plates. The Swift XRT observation clearly detects this bright source, and identifies *102b* as the counterpart. The SDSS-III spectrum classifies *102b* as broadline QSO, at $z=0.1296$.

103: No optical counterpart candidate brighter than $B = 21$ mag within either the ROSAT or Swift XRT error circle.

104a: Variability with amplitude ≤ 0.2 mag is indicated, but is very uncertain. The Swift XRT position confirms this source as the counterpart.

105a: Difficult because of faintness. More frequently faint than bright. There are brightness changes up to 0.3 mag within some hours. Classified as AGN by Zickgraf et al. (2003). The SDSS-III spectrum classifies it as QSO at $z=0.1019$, with “starburst broadline” sub-class.

106: Within X-ray error circle no object brighter than $B = 21$ mag. *106a* is possibly variable, but very uncertain. The Swift XRT observation did not detect this source, but due to the short exposure this is only mildly hinting at X-ray variability.

107a: During the whole interval of observation the brightness declines. At the beginning (J.D. 243 7696 - 8530) 16.3–16.7 mag, then (- 244 5100) 16.5–17.0 mag, thereafter often fainter than 17.0 mag. Extreme value on a good plate (244 7945) at 18 mag. Light variations within one night seem questionable. One plate of Tautenburg Schmidt telescope (J.D. 244 2094) shows the object at 17.3 mag. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5230$.

108a: Fainter than the Sonneberg plate limit, therefore not tested for optical variability. Pair consists of a QSO at $z=0.436$ and a star (Oscosz et al. 1997). The SDSS-III spectrum classifies it as broadline QSO at $z=0.4365$. *108b*: Spectral classification from Garcia Lopez et al. (2000) who discard Coma Berenices cluster membership which had been proposed by Randich et al. (1996), and find no spectroscopic evidence for chromospheric activity; this and f_X/f_{opt} make it unlikely to be the optical counterpart.

109a: On plates of the Sonneberg astrographs always invisible. The SDSS-III spectrum classifies it as QSO at $z=0.3392$, with “starburst broadline” sub-class. *109b*: No brightness variations. *109c,d*: Two galaxies, possibly intrinsic pair. *109e* is invisible on all Sonneberg plates. Magnitude on POSS print from 1950 Jan 09 about $B = 18.5$, and from 1955 May 21 about $B = 20.5$. Therefore obviously variable, but too far from the ROSAT position to be considered as counterpart of the X-ray source. Thus, we consider *109a* as the most likely counterpart.

110a: The SDSS-III spectrum classifies it as broadline QSO at $z=1.004$.

111a: This very blue object is marginally seen on only a few plates, which show it at about 18.4 mag. On POSS 1435 blue print (1955 May 21) it is about 18.5 mag. On one Tautenburg Schmidt plate (J.D. 244 8356) scarcely visible at about 20 mag. Therefore very probably variable.

112a: Not investigated on Sonneberg plates.

113a: Single white dwarf PG 1254+223. Stellar object with UV excess (Green et al. 1986); identified by McCook & Sion (1987). UVB measurements reported by Cheselka et al. (1993), proper motion by Eggen et al. (1967). The XMM position (2XMM J125702.3+220151) is plotted with somewhat larger uncertainty radius for visibility reasons (Fig. 40a). This and the very soft X-ray spectrum leaves no doubt about the identification.

114a: The brightness varies irregularly within some days with amplitudes of about 0.15 mag. The mean value of magnitudes varies slowly: 14.0–14.1 mag in 1962–1967, 14.2 mag in 1972–1978, then brightening, 14.0 mag in 1991, thereafter slightly fading.

115a-c: Too faint for Sonneberg plates. The SDSS-III spectrum classifies *115a* as M3 star. Could be a flare star, though the spectrum shows only marginal H α emission. M star classification and f_X/f_{opt} is also marginally consistent with chromospheric emission. *115c* is a galaxy at $z=0.2257$ according to SDSS-III, and unlikely the counterpart because of the large distance of 50'' and the f_X/f_{opt} ratio.

116a: Brightness usually at about 16.7 mag. From J.D. 243 7764 - 7806 about 16.9 mag. In the interval 243 8501 - 8525 fainter (about 17.1 mag). From 244 4342 - 4367 brighter (about 16.5 mag), from J.D. 244 6113 - 6121 still brighter (16.4 mag), then until 244 6177 fading (16.7 mag). Occasionally fading to nearly 17.5 mag. The SDSS-III spectrum classifies it as QSO at $z=0.1798$, with “starburst broadline” sub-class.

119a: Sometimes the object seems to get slightly brighter than 18 mag. But this is questionable, because it is at the very plate limit. The blue colour suggests an AGN nature. The SDSS-III spectrum classifies it as QSO at $z=0.4363$.

120a: Visible only on 20 of the best plates. Within the error circle no object brighter than about $B = 21$ mag. The Swift XRT observation did not detect this source, and the upper limit implies a fading of about a factor three.

121a,b: Both objects have blue colour. In spite of the relatively large optical amplitude of *121a* no exact statement about the brightness variations can be given because of faintness. Small magnitudes are favoured. Brightness changes seem to occur within a single night. One Tautenburg plate (JD 244 2094) shows the object at $B=17.7$ mag. The SDSS-III spectrum classifies *121b* as A0 star. Due to the smaller distance to the X-ray position and the optical variability we prefer *121a* as counterpart over *121b*.

122a: Galaxy; the NED reports a redshift of $z=0.4$, but without reference. If the redshift is true, this should be an AGN, since X-ray emission from non-active galaxies is not luminous enough




Fig. 14. Blow-up of the light curve of Com116a (Fig. 41a) showing pronounced variability.

to be seen in the ROSAT all-sky survey up to $z=0.4$. The SDSS-III spectrum classifies it as starforming galaxy at $z=0.1114$.

123a: See also Brinkmann et al. (1995).

124a: Too faint for Sonneberg plates. Blue magnitudes: On POSS print No. 135 (J.D. 243 3291): 16.4 mag, on Tautenburg Schmidt plate No. 4004 (J.D. 244 2094): 17.8 mag. The SDSS-III spectrum classifies it as broadline QSO at $z=0.3478$.

126a: Southern (blue) component of a double galaxy inside a cluster of faint galaxies (NSCS J122935+213714) at $z=0.25$ (Lopes et al. 2004). The blue optical colour, X-ray hardness ratio and f_X/f_{opt} ratio suggest an AGN nature. The Swift XRT observation marginally detects this source, which is consistent with the very soft X-ray spectrum. The Swift XRT image suggests also large-scale diffuse emission. The SDSS-III spectrum classifies it as broadline QSO at $z=0.3484$.

127: Inside a cluster of faint galaxies. Within the error circle there are 3 galaxies fainter than 18 mag. The NED lists **127a** as QSO, but without reference. See also Mickaelian et al. (2006). The SDSS-III spectrum classifies **127a** as broadline QSO at $z=0.2336$.

128: A ROSAT HRI pointing improves the X-ray coordinate to an error circle of $10''$ (1RXH J125627.3+213117) and falling on top of the galaxy **128a**. The SDSS-III spectrum classifies it as QSO at $z=0.0757$, with “starburst broadline” sub-class.

129: No optical counterpart candidate brighter than $B = 20$ mag within the ROSAT error circle.

130a: Only on the best astrograph plates visible at about $B = 18$ mag. On some poor plates there seem to exist some brightenings to about 16 mag, but this must be regarded as uncertain. One Tautenburg Schmidt plate 7482 (J.D. 244 8356) shows the object at about $B = 18.5$ mag. The SDSS-III spectrum classifies it as broadline QSO at $z=0.2354$.

131a: Could not be tested for variability because of blend. The SDSS-III spectrum classifies it as starburst galaxy at $z=0.1156$.

132a: Galaxy (by extent). Zickgraf et al. (2003) reports “BLUE-WK”, a moderately blue continuum, weak point-like object. The f_X/f_{opt} ratio suggests an AGN nature. Too faint on Sonneberg plates for variability assessment. Even invisible on two Tautenburg plates. The SDSS-III spectrum classifies it as QSO at $z=0.1896$, with “starburst broadline” sub-class.

133a: Because of the blue colour probably an AGN (or less likely a CV), and thus more likely the counterpart than the

galaxy **133b**. The SDSS-III spectrum classifies it as broadline QSO at $z=1.4783$.

134a: 4C+21.35. Too faint for Sonneberg plates. On POSS print 135 (J.D. 244 3291) about 1.5 mag brighter than on Tautenburg Schmidt plate 4004 (J.D. 244 2094). See also the optical catalogue of QSOs by Hewitt & Burbidge (1987) and Brinkmann et al. (1995). The SDSS-III spectrum classifies it as broadline QSO at $z=0.4338$.

135a: S 10938 Com (Richter et al. 1995).

137a: Very slow variation with a time scale of months to years. At the beginning of the observations around JD 243 7705: ~ 13.05 mag. Later at 243 7605–7825: ~ 13.1 mag, 243 8083–8173: ~ 13.2 mag, 243 8410–9205: ~ 13.0 – 13.2 mag, 243 9530–244 4367: ~ 13.0 – 13.1 mag, 244 4634–4697: ~ 12.9 – 13.0 mag, 244 5000–9163: ~ 12.8 – 13.0 mag, 244 9410–9864: ~ 12.8 – 12.9 mag. The colour indicates that it may be a coronal active object.

138a: Non-stellar due to extent, coincident with near-infrared source 2MASX J12380987+2114014. The SDSS-III spectrum classifies it as galaxy at $z=0.1089$. **138b:** Not tested for variability. Spectral type and f_X/f_{opt} do not match, so not a counterpart candidate.

139: Zickgraf et al. (2003) identify the galaxy **139a** as the counterpart. The Swift observation clearly favours the galaxy as the counterpart. The X-ray intensity is about a factor of three less than during the ROSAT all-sky survey. The SDSS-III spectrum classifies **139a** as galaxy at $z=0.0514$ with “AGN broadline” sub-class, and **139b** as A0 star.

140a: The SDSS-III spectrum classifies it as QSO at $z=0.1391$, with “starburst broadline” sub-class.

141a: IR Com = S 10932. Atypical UG star with eclipses (see Richter & Greiner 1995b, Kroll & Richter 1996, and Richter et al. 1997). The heavy absorption at X-ray wavelengths is likely an artefact of the small photon number (24) statistics and/or an inappropriate X-ray spectral model, since the optical colour is pretty blue. The SDSS-III spectrum classifies it as CV, and also shows substantial He II emission.

142a: Other name: HIP 61204. Variability discovered by *Hipparcos* (*Hipparcos* & *Tycho* catalogues 1997). Two close components of nearly equal brightness. Obviously a chromospherically active star.

143a: It cannot be excluded that the object is slightly variable with an amplitude of about 0.1 mag. However, to verify such variability, photoelectric observations are required.

144a: Narrow-line Sy1 galaxy at $z=0.335$, with ROSAT data and optical variability based on Sonneberg data already reported earlier (Greiner et al. 1996). The XMM position (2XMMp J122541.9+205503) confirms the identification beyond doubt.

145a: Variability discovered by *Hipparcos* (*Hipparcos* & *Tycho* catalogues 1997) from which also the B amplitude is extracted. XMM finds two sources one of which (2XMMp J123209.9+205508 = 1XMM J123210.0+205507) coincides with **145a**. The other (2XMMp J123209.4+205553) has no visible optical counterpart on the POSS sky survey print.

146a: Outside the error circle. Not tested for variability. Inside the error circle no object brighter than 21 mag. A Swift/XRT observation did not reveal this source; the upper limit implies a fading by a factor of 5. The SDSS-III spectrum classifies **146a** as starforming galaxy at $z=0.0851$.

147a: Just outside the error circle, but spectral type and f_X/f_{opt} suggest this to be the counterpart. Too bright for astrograph plates.

148a: Blue object. Because of faintness only on the best 25 Sonneberg plates from 1962–1993 visible. The brightness varies irregularly between 17.2 mag and 18.0 mag. On some poor

plates the object seems to be slightly brighter than 17 mag, but this must be regarded as questionable. Two Tautenburg Schmidt plates give similar brightness: 17.6 mag at J.D. 244 2094 and 18.0 mag at 244 2453. On the other hand, POSS print No. 1435 (1955 May 21) shows the object at about 18.5 mag. The SDSS-III spectrum classifies it as broadline QSO at $z=1.3540$. *148b* is constant within the error limits.

149a: Galaxy (18.4 mag) just outside the error circle which the SDSS-III spectrum classifies as starburst galaxy at $z=0.1963$. Within the error circle are two faint (about 21 mag) galaxies which are much too faint for Sonneberg plates. The Swift/XRT observation reveals two blobs of diffuse emission about 2.5 away towards the north-west and south-east, respectively, but no point source at the ROSAT position. This and the hard X-ray spectrum suggest a galaxy cluster nature.

150a: Possibly AGN due to blue colour. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5678$.

151a-c: Magnitudes of all three objects in Table 5a are taken from the USNO-B1 catalogue. Too faint for Sonneberg plates. The Swift/XRT observation, despite only with 540 sec exposure, reveals a 2.5σ X-ray source at a rate consistent with that of the RASS rate, and the X-ray position clearly suggests *151c* as counterpart. The f_X/f_{opt} ratio and the relatively hard X-ray spectrum suggest an AGN nature. The SDSS-III spectrum classifies it as broadline QSO at $z=0.8401$.

152a: Very faint for Sonneberg plates. Seems to be variable within the limits 17.0–17.3 mag, in some cases also fainter. But because near the plate limit, these variations are not quite sure. The SDSS-III spectrum classifies it as broadline QSO at $z=0.6332$.

153a: On Sonneberg plates diffuse appearance (blend or galaxy). Classified as AGN by Zickgraf et al. (2003). The SDSS-III spectrum classifies it as QSO at $z=0.0756$ with “starburst broadline” sub-class.

155a: Too faint for Sonneberg plates. $B-V$ colour and f_X/f_{opt} ratio suggest an AGN nature, and Swift XRT localization proves this identification. The SDSS-III spectrum classifies it as broadline QSO at $z=0.4287$.

156a: Not tested for variability because diffuse appearance (galaxy).

157a: Obviously variable, but with small amplitude (15.6–15.8 mag). On two plates of Tautenburg Schmidt telescope similar values. Brightness cycles with a duration of about 12 days possibly exist. Superposed are some fainter values up to 16.0 mag which, if real, could be the result of eclipses in an RS CVn system. Faint observations: J.D. 243 8085.459 (15.95: mag), 243 8085.501 (15.9 mag), 243 8146.517 (16.0: mag), 244 4701.444 (15.95: mag), 244 5384.628 (16.0: mag), 244 5403.540 (15.9 mag), 244 6109.541 (15.9: mag), 244 6563.507 (16.0: mag), 244 7613.457 (16.0: mag); 244 8683.507 (16.0 mag). The X-ray spectrum reveals substantial absorption, implying an $A_V \sim 1.8$ mag, which is consistent with the very red optical colours.

158a: Not tested for variability, but possibly a coronal active star.

159a: Sy1 galaxy Mrk 771 $\equiv$ TON 1542 $\equiv$ PGC 41532 at $z=0.063$. Stellar object with UV excess (Green et al. 1986), see also optical catalogue of QSOs by Hewitt & Burbidge (1987), and high-energy spectrum by Malaguti et al. (1994). Diffuse appearance on Sonneberg plates, so not tested for variability.

162a: S 10935 (see Richter et al. 1995). Mean magnitude about 15.5 mag. According to Bade (priv. comm.) K spectral type.

163a: Irregular light changes. Coincidence with the radio source 7C 1237+2010 $\equiv$ 87GB 123717.5+201037 (Brinkmann




Fig. 15. Blow-up of the light curve of Com157a (Fig. 41a) showing pronounced variability.

et al. 1997). This and the blue colour suggest a QSO identification which is proven by the SDSS-III spectrum, giving also $z=0.2394$.

164a: Difficult because of the small amplitude. Irregular variable with waves from about some dozens to more than 100 days (Fig. 16), suggestive of chromospheric activity. The SDSS-III spectrum classifies it as F9 star. *164b* is constant on Sonneberg plates.




Fig. 16. Blow-up of the light curve of Com164a (Fig. 41a) showing pronounced variability.

165: In a 13.7 ksec observation with the ROSAT HRI in Jun 2-4, 1995 (Obs.-ID 800435), two nearby sources are detected: the northern at $12^h26^m43^s.7 +19^\circ50'46''$ (with 0.0018 cts/s), the southern at $12^h26^m43^s.0 +19^\circ50'11''$ (with 0.0007 cts/s). The northern source is a factor 4 fainter than the all-sky survey source, the sum of both sources is still fainter by a factor of 3 (accounting for the factor 3 difference in sensitivity between PSPC and HRI for spectrally hard sources). It is not clear whether one of the two HRI sources is related to the all-sky survey source. Also, none of these two sources coincides with object *165a* which the SDSS-III spectrum identifies as galaxy at $z=0.2232$. There are several radio sources within the all-sky survey source’s error circle, but none coincides with either *165a* or the two HRI sources. The Swift XRT observation reveals a blob of faint, diffuse emission which is centred between the two HRI positions, but extends much further, to about 40 arcsec. The total flux is identical to that measured during the ROSAT all-sky survey. This suggests diffuse emission, either from the galaxy *165a* or a larger structure.

166: Cluster of very faint ($B \sim 21$ mag) galaxies. The next named galaxy cluster is more than 3’ away. The Swift XRT observation reveals faint, diffuse emission which extends about 3 arcmin towards the south.

167a: Galaxy with number 041 in cluster Abell 1570 (Flin et al. 1995) for which the SDSS-III spectrum provides a redshift of $z=0.2139$. The X-ray hardness ratio is large, so the ROSAT emission is rather cluster gas emission than X-rays from this individual galaxy. The optical cluster has an extent of $10' \times 15'$,

and 167a is near its centre. The Swift XRT observation reveals large-scale, diffuse emission.

168a: Classified as AGN by Zickgraf et al. (2003), and as counterpart of the X-ray source. This is supported by the Swift XRT position. The SDSS-III spectrum classifies it as QSO at $z=0.1603$ with “starburst broadline” sub-class. **168b:** Suspected variable, but too faint for confirmation. Light variations are not certain. **168c:** The SDSS-III spectrum classifies it as F9 star.

169a: On $\sim 70\%$ of the Sonneberg plates invisible. Some observations on poor plates near 16 mag are questionable. On 4 Tautenburg Schmidt plates the B magnitudes are: 17.4 mag (1974 Feb. 16; JD 244 2095), 17.9 mag (1992 Mar. 1; JD 244 8683), 17.9 mag (also 1992 Mar. 1), and 17.9 mag (1992 Mar. 2; JD 244 8684). Zickgraf et al. (2003) report “EBL-WK”, an extremely blue continuum, weak point-like object. Kouzuma et al. (2010) identify it as AGN candidate due to the NIR colours from 2MASS. The SDSS-III spectrum classifies it as QSO at $z=0.22457$ with “starburst broadline” sub-class.

171a: Classified as possible AGN by Mickaelian et al. (2006). The Swift XRT position confirms this without doubt as the counterpart. The SDSS-III spectrum classifies it as QSO at $z=0.1313$ with “starburst broadline” sub-class.

172a: Galaxy in cluster ACO 1548 at $z=0.16$ (see Schneider et al. 1983). The $0.1\text{--}2.4$ keV X-ray luminosity of 1.3×10^{43} erg/s and spectral hardness suggests cluster emission rather than the galaxy as counterpart.

173b: This blue object is at the magnitude limit of Sonneberg plates. A variability of small amplitude near 18 mag therefore seemed initially doubtful. But the Tautenburg Schmidt plate from J.D. = 244 2453 shows the object distinctly fainter (B about 18.6) than on the POSS print 1576 (J.D. 243 5548, B about 18.0 mag). Therefore the object is probably variable. The SDSS-III spectrum classifies it as broadline QSO at $z=0.5417$. **173a** is likely also an AGN due to the blue colour, but based on the optical variability we prefer **173b** as counterpart.

174a: Narrow-line Sy1 galaxy at $z=0.28$, with ROSAT data already reported earlier (Greiner et al. 1996).

175a: Not in the USNO-A2, but in USNO-B1 catalogue, from which magnitudes are taken. Coincides within $1''$ with radio source NVSS 124538+192050 (Condon et al. 1998). We therefore identify this object as an AGN. This is confirmed by the SDSS-III spectrum: broadline QSO at $z=0.6930$.

176a: Bright galaxy pair NGC 4561 $\equiv$ KPG 346 $\equiv$ LEDA 42020 at $z=0.0047$.

177: No optical counterpart candidate brighter than $B = 21$ mag in the ROSAT error circle.

178a,b: Both are viable counterpart candidates, and both are too faint for Sonneberg plates. The SDSS-III spectrum identifies **178a** as broadline QSO at $z=0.4994$, and **178b** as broadline QSO at $z=0.9963$.

180a: There seem to be brightness changes within 0.3 days between 17.0–17.2 mag. But considering the faintness and the small amplitude, this is very uncertain. The USNO as well as 2MASS colours suggest a K5 star suffering no extinction, and thus the f_X/f_{opt} ratio is (marginally) consistent with this star to be the counterpart. The source is not detected in a Swift XRT observation, with an upper limit about a factor 3 below the RASS rate, so possibly the RASS detected the object during an active period, explaining the elevated f_X/f_{opt} ratio.

182a: Magnitudes taken from USNO-B1. **182b:** Variability cannot fully be excluded because of relatively high dispersion of the magnitudes. **182c:** Coincides with the 365MHz radio source TXS 1223+192 (Douglas et al. 1996) which most likely corresponds to GB6 J1225+1858. Given the radio emission and

f_X/f_{opt} , we identify this source as the optical counterpart of the ROSAT source.

183a: Difficult because of faintness. Only visible on good plates. The magnitude changes are characterized by a mean brightness level from which declines occur with a duration of several days. The mean brightness level is about 17.1–17.3 mag at the beginning of the observations J.D. = (243 7700 – 8902) and rises slowly to 16.8–17.0 mag (244 4663 – 8362), see light curve in Fig. 17. AGN identification from Zickgraf et al. (2003). The SDSS-III spectrum identifies it as broadline QSO at $z=0.2223$.




Fig. 17. Blow-up of the light curve of Com183a (Fig. 41a) showing pronounced variability.

184a: The brightness varies with a time scale of weeks and months. An isolated observation on Tautenburg Schmidt plate 4272 (J.D. 244 2453) shows the object at magnitude 17.4.




Fig. 18. Blow-up of the light curve of Com184a (Fig. 41a) showing pronounced variability.

184a: The SDSS-III spectrum identifies it as broadline QSO at $z=0.5992$.

185a: Difficult because of faintness. There are typically waves with a length of about one week and 0.4 mag amplitude. Superposed are slow irregular changes. Variations of 0.2 mag

within one night seem to occur, but this is unsure. On POSS print 89 (JD 243 3391) at 17.0 mag, and on Tautenburg plates 1576 (JD 243 5548) at 17.2 mag and on 4011 (JD 244 2095) at 18.0: mag. The SDSS-III spectrum identifies it as QSO at $z=0.1970$ with “starburst broadline” sub-class.




Fig. 19. Blow-up of the light curve of Com185a (Fig. 41a) showing pronounced variability.

186a: During some hours the whole brightness interval is passed through, but seemingly not quite regular. A possible relationship to eclipsing stars of the W UMa type is therefore questionable. A relationship to RS CVn stars cannot be excluded. The EW star SS Com is 6' apart from the ROSAT position, and not considered as counterpart.




Fig. 20. Blow-up of the light curve of Com186a (Fig. 41a) showing pronounced variability.

187a: QSO at $z=0.723$ (Wills & Wills 1976). See optical catalogue of QSOs by Hewitt & Burbidge (1987), and Brinkmann et al. (1995). Difficult to investigate because near the plate limit. Sometimes fadings of short duration (about 1 hour).

188b: Galaxy MCG +03-32-083 at $z=0.07$, brightest member of a cluster of galaxies. The X-ray emission as seen by ROSAT as well as XMM (2XMMp J124117.3+183430) is not only clearly extended, but elongated with the same orientation as the galaxy, thus the galaxy itself rather than the cluster is the most likely optical counterpart. This is supported by the XMM position which aligns well with the galaxy centre. Identification also proposed by Zickgraf et al. (2003). The SDSS-III spectrum identifies **188c** as galaxy at $z=0.0711$.

189a: At the border of a cluster of galaxies. The hard X-ray spectrum indicates a cluster nature, rather than that of an AGN.

190: No optical counterpart candidate brighter than $B = 20$ mag in the error circle. HD 106972 is 84" distant from the ROSAT position, too far to be considered the counterpart.

191: No optical counterpart candidate brighter than $B = 20$ mag in the error circle.

192a: FN Com = S 8058 (Hoffmeister 1964). The observations of Wenzel and Ziegler (see Meinunger & Wenzel 1968) in connection with the yellow colour and the X-radiation let us suppose that this is a star of the BY Dra type.

193: No optical counterpart candidate brighter than $B = 20$ mag within the error circle. The following objects all have a rather large distance to the X-ray source. **193a:** The blue colour suggests an AGN classification. **193c:** QSO at $z=1.401$, see optical catalogue of QSOs by Hewitt & Burbidge (1987). It is variable according to Uomoto et al. (1976). Variability is confirmed by observation on POSS (17.1 mag at JD 243 5548) and Tautenburg plates (>17.6 mag at 244 2095, 17.9 mag at 244 8683.60, and 17.8 at 244 8683.66). On Sonneberg plates beyond the plate limit.

194: No object brighter than 18.5 mag within the ROSAT error circle, both objects **194a** and **194b** are too faint for Sonneberg plates. **194b** is nearly one magnitude fainter on POSS 1576 than on Tautenburg plates, but it is difficult to say whether this effect is caused by variability or by the diffuse image of this object which may possibly be a galaxy. The SDSS-III spectrum identifies **194b** as QSO at $z=0.2502$ with “starburst broadline” subclass.

195: No object brighter than 21 mag within the error circle. Remarkable is its high X-ray luminosity during the ROSAT all-sky survey, which has declined by at least a factor of 8 to explain the non-detection in the Swift XRT observation. The SDSS-III spectrum identifies **195a** as broadline QSO at $z=0.1341$. But due to its large distance to the X-ray centroid the association remains open. Turner et al. (2010) list the X-ray source as “BL Lac-type object”.

196a: Triple star. In addition to the second component of M-type ($R = 13.9$, $B = 15.7$, $V = 14.6$), there is a faint red third component (B about 18.4). The second component is possibly variable, but because of blending this is not sure. **196a** is listed among the periodic variables coincident with bright ROSAT sources (Kiraga 2012).

197a is an AGN according to Zickgraf et al. (2003), despite not particularly blue colour in the USNO catalogue. It was not investigated on Sonneberg plates due to blending with the bright star **197b**. **197c** is a galaxy classified as “Red-WK” by Zickgraf et al. (2003), which is classified as starforming by the SDSS-III spectrum, at $z=0.0667$. It is constant on Sonneberg plates.

198a: On two overlapping POSS prints and two Tautenburg plates of nearly equal brightness.

199: The RRc-type star AN Com is 96" away from the ROSAT position and thus not considered as counterpart. The QSO **199a** is the likely counterpart, and the SDSS-III spectrum provides $z=0.2540$.

200a: Not tested for variability because of diffuse appearance on Sonneberg plates (blend or galaxy). The SDSS-III spectrum identifies it as QSO at $z=0.0753$, with “starburst broadline” sub-class.

201: Both objects too faint for Sonneberg plates. The Swift XRT observation suggests object **201b** as counterpart.

202a: Slow light changes (within years) are superposed by waves of small amplitude (0.1–0.3 mag) and short duration (several days to 50 days). Small variations probably also within one night. The mean brightness rises (with fluctuations) from 16.5 mag since the beginning of the observations (J.D. 243 7650) to

16.1 mag (J.D. 243 8080 - 9150) with a flat dip of 16.3 in J.D. 243 8500. From J.D. 243 9200 - 9620 variations between 16.2 and 16.4 mag. Sporadic observations between J.D. 244 2840 and 3250 give 16.5–17.0 mag. From J.D. 244 4345 - 6530 bright again between 16.0 and 16.5 mag. Thereafter (only few observations) 16.4–16.9 mag, only at J.D. 244 8350 a little brighter, about 16.3. The SDSS-III spectrum identifies it as broadline QSO at $z=0.3526$.

203: Object *203a* is coincident with the radio source FIRST J124406.5+174831. The SDSS-III spectrum identifies it as galaxy (no emission lines) at $z=0.1670$. Could possibly be a blazar. Alternatively a cluster of galaxies, as *203b* is another galaxy (though at unknown redshift), and the X-ray spectrum is pretty hard.

204a: On POSS print 1572 (1956 Mar. 14) $B = 16.6$ mag, on Tautenburg Schmidt plates No. 4027 (1974 Feb. 17) $B = 17.4$, No. 7869 (1992 Feb 9) $B = 18.3$ mag, No. 7871 (1992 Feb. 9) $B = 18.2$ mag. On Sonneberg plates invisible. The SDSS-III spectrum identifies it as QSO at $z=0.2630$ with “starburst broadline” sub-class.

205a: Close pair of stars. Not tested for variability.

206a,b: It is not easy to decide which of the two objects is the better candidate for the ROSAT source. While the FG star *206a* is consistent with the soft X-ray spectrum, its f_X/f_{opt} ratio is high, somewhat borderline (see Fig. 2). On the contrary, the QSO *206b*, being at $z=0.5478$ according to the SDSS-III spectrum is clearly outside the ROSAT error circle. We tend to think that *206a* is the counterpart.

207a: Too faint for Sonneberg plates. On POSS print and 4 Tautenburg Schmidt plates of equal brightness. The SDSS-III spectrum identifies it as broadline QSO at $z=0.4067$.

208c: Brightness changes seem to be indicated, but not certain. The Swift XRT observation does not detect this X-ray source; the upper limit is not constraining due to the very soft spectrum during the ROSAT all-sky survey.

209a,b: Blended on Sonneberg plates. Not tested for variability, no variability found by Hewitt & Burbidge (1987). On POSS prints Nos. 1572 and 1576 (1956 Mar. 14) and on Tautenburg Schmidt plate no. 4027 (1974 Feb. 17) of equal brightness. The SDSS-III spectrum identifies *209a* as broadline QSO at $z=0.12822$.

210a: The SDSS-III spectrum identifies it as galaxy (no emission lines) at $z=0.0720$.

211a: Not in USNO-A2 or USNO-B1, but in APM catalogue, from which the magnitudes in Table 5a are taken. Only on a single Sonneberg plate barely indicated: J.D. 243 8853, $B = 17.7$: mag. Visible also on two POSS prints: 1956 Mar. 14, $B = 21$ mag and 1956 Mar. 15, $B = 18.4$ mag. On three Tautenburg Schmidt plates invisible, fainter than 20.5 mag. Therefore variable. Type unknown, may be AGN or CV.

212a: Blended on Sonneberg plates. The Swift XRT position clearly favours the galaxy which according to the SDSS-III spectrum is a galaxy without emission lines at $z=0.0804$. The red colour, low f_X/f_{opt} and hard X-ray spectrum argue against an AGN. The Swift XRT flux is about a factor 4 lower than the ROSAT measured flux. The hard X-ray spectrum would also be unusual for a tidal-disruption event.

215a: Within a cluster of faint galaxies. Remarkable is the high X-ray luminosity. Too faint for Sonneberg plates. On POSS print and on five Tautenburg plates similar magnitudes. Coincides with radio source NVSS 123305+170132 (Condon et al. 1998) $\equiv$ FIRST J123305.1+170132, therefore likely AGN/Blazar (see also Laurent-Muehleisen et al. 1997). The SDSS-III spectrum identifies it as star with “WD magnetic”

sub-class, but without any clear sign of either absorption nor emission line this is doubtful. The featureless blue spectrum argues more in favour of a BL Lac type object.

216a: Coincides with the radio source FIRST J123528.8+170036. The SDSS-III spectrum identifies it as galaxy at $z=0.3806$. Likely a BL Lac object.

217: Within a cluster of faint galaxies. No optical counterpart candidate brighter than $B = 21$ mag within the ROSAT error circle. The Swift XRT observation did reveal extended emission, suggesting emission from cluster gas.

218a: QSO LBQS 1229+1711 at redshift 0.209 (Hewett et al. 1995).

219: Invisible on Sonneberg plates. The Swift XRT observation does not detect this source, suggesting about a factor 10 variability.

220: According to Gioia and Luppino (1994) rich loose cluster of galaxies. No dominant galaxy. Hard X-ray emission ($HR1=1$) consistent with cluster-emission. Distance to the *Einstein* source MS 1241.5+1710 is $1'.5$, which in turn had been associated to the galaxy cluster at $z=0.31$ (Stocke et al. 1991). Also ASCA source 1AXG J124359+1653 (Ueda et al. 2001). The Swift XRT emission indicates large-scale diffuse emission, with the maximum about $1'.7$ north, at $12^{\text{h}}44^{\text{m}}01^{\text{s}}.2 + 16^{\circ}53'51''$, consistent with the ASCA position.

221: Invisible on Sonneberg plates. Source not detected with a Swift XRT observation. Given the very soft spectrum of the ROSAT source, the upper limit is not constraining, however.

222a: Probably eclipsing binary. Difficult because of small amplitude. Supposed minima, in parentheses magnitudes: J.D. 243 7820.474 (14.6), 8085.668 (14.6), 8142.454 (14.7:), 9611.380 (14.6), 9618.461 (14.6), 244 4702.540 (14.7:), 6552.376 (14.6), 7206.536 (14.7), 7613.447 (14.6), 9482.403 (14.7). We did not succeed in finding a period. Some of the minima may be erroneous because of small amplitude. Presumably chromospherically active star.

223a,b: Possibly AGN due to very blue colour. The SDSS-III spectrum identifies it as broadline QSO at $z=2.4468$. **223c:** Coincides with radio source NVSS 122944+164002 (Condon et al. 1998) $\equiv$ FIRST J122944.5+164003, thus most likely a QSO.

225: Source not detected with a Swift XRT observation, indicating a fading of about a factor of 10, or very soft spectrum. The SDSS-III spectrum identifies *225b* as starforming galaxy at $z=0.0763$.

226a: The elements given by Meinunger & Wenzel (1968) need only small corrections (Richter & Greiner 1995a). They are: $m = 243\,7668.521 + 0.7354422 \cdot E$. The minima are given in Table 6. The SDSS-III spectrum identifies *226b* as a K7 star, thus unlikely the counterpart due to the high f_X/f_{opt} .

227a: QSO at $z=1.017$, on Sonneberg plates small irregular brightness changes, probably also within some nights. The SDSS-III spectrum identifies *227b* as galaxy at $z=0.0635$.

228: The Swift XRT observation reveals a bright, extended source. Together with the hard X-ray spectrum this suggests a cluster origin. Thus, *228a* is likely not the counterpart.

229a: Very close pair of stars with similar B magnitude. A supposed variability of the northern component is uncertain because of faintness and blending. Identification as K star from Micaelian et al. (2006). *229b* is probably a AGN due to its blue colour, and therefore a good counterpart candidate.

230: See Brinkmann et al. (1995). In the midst of a cluster of galaxies. The X-ray emission could also be associated to the Abell cluster ACO 1569 (see e.g. Gomez et al. 1997), rather than to the radio galaxy *230a* at $z=0.0684$.

Table 6. Minima of CN Com $\equiv$ Com226a

J.D. (2400000+)	B	E	B-R (d)
37668.503	13.85	0	-0.018
37737.649	14.0	94	-0.004
37749.416	13.9	110	-0.004
37752.377	14.0	114	+0.016
37821.502	13.9	208	+0.009
38090.665	13.7	574	0.000
38407.630	13.9	1005	-0.010
38471.633	13.7	1092	+0.009
39205.593	13.8	2090	-0.002
44704.501	13.9	9567	+0.004
45780.439	13.9	11030	-0.009
45822.379	13.7	11087	+0.010
47612.455	13.7	13521	+0.020
48329.487	13.8	14496	-0.004
48357.436	13.9	14534	-0.002
48682.506	13.8	14976	+0.003

231a: Other name 3C 275.1. Optical identification see Sandage et al. (1965). According to Uomoto et al. (1976) and Hewitt & Burbidge (1987) variable. See also Brinkmann et al. (1995). The SDSS-III spectrum identifies it as broadline QSO at $z=0.5552$.

232: Invisible on Sonneberg plates. Not detected in a Swift XRT observation.

233: Invisible on Sonneberg plates. Marginally detected in a Swift XRT observation. Due to the soft X-ray spectrum of the ROSAT source, the low count rate is consistent with the ROSAT rate. The f_X/f_{opt} ratio suggests an AGN nature.

234a: Invisible on Sonneberg plates. Likely an AGN due to its blue colour on POSS sky survey prints. The Swift XRT position identifies this object as counterpart. The radio source FIRST J123439.3+160611 (Becker et al. 1997, catalogue version of Apr 2003; positional error of $1''$) does not coincide with 234a.

235a: Other name: SDSS J123544.26+160536.3; QSO at $z=0.07$ (data release 5 as obtained Jun. 28, 2006; <http://www.sdss.org/dr5/products/spectra/getspectra.html>).

236b: Other name: SDSS J123942.92+160612.1; galaxy at $z=0.07$. (data release 5 as obtained Jun. 28, 2006; <http://www.sdss.org/dr5/products/spectra/getspectra.html>). See also Mickaelian et al. (2006). The SDSS-III spectrum identifies it as starforming galaxy at $z=0.0702$. 236c: slightly outside the error circle of ROSAT position, not tested for variability. 236d: Coincides with NVSS 123945+160600 (Condon et al. 1998) $\equiv$ FIRST J123945.8+160558 (Becker et al. 1997; catalogue version of Apr 2003), thus likely AGN. The SDSS-III spectrum identifies it as starforming galaxy at $z=0.2808$. The Swift XRT position suggests this object as the counterpart of the ROSAT source.

237: No optical counterpart candidate brighter than $B = 21$ mag in or near the error circle of the ROSAT position. The X-ray source is located near the border of our extracted X-ray data region, and its detection significance is near the detection limit, thus it could potentially be a spurious X-ray detection. It is not detected in a Swift XRT observation.

238: The soft X-ray spectrum and the f_X/f_{opt} ratio support the identification with the bright FG star.

3.5. Notes on individual objects in the Sge field (Table 5b)

1a: A0 star HD 185334, coincident with the radio source WSRTGP 1935+2408 (Taylor et al. 1996). The RXS position favours 1b as optical counterpart.

2a: Algol star with variable period length (see GCVS). Whether it is a chromospheric active star is still to be proved.

3a: The USNO-A2 magnitudes are incompatible with the spectral type; the B magnitude must be wrong. 3b: GSC 2141 123 is a very close pair of stars of nearly equal B magnitudes.

4a: Optically variable with short waves (4–10 days) of small amplitude (0.2 mag) and long waves (years) with 0.3 mag amplitude (Fig. 21).




Fig. 21. Blow-up of the light curve of Sge004a (Fig. 41b) showing pronounced variability.

5a: The Swift XRT position is marginally consistent with this object, so we suggest it as counterpart of the ROSAT source though an optical variability could not be established photographically.

6a: Soft X-ray spectrum is consistent with G spectral type. 6b: Close pair of stars. On Sonneberg astrograph plates totally blended by 6a.

7a: Variability near the detectability limit. Amplitude only 0.2 mag. Waves with a length of several days to some months. Standstills of several 100 days at about 12.6 mag (Fig. 22).

8a: On Sonneberg plate unresolved blend of four stars. The Swift XRT position falls on the centre of the top three, and excludes the southern-most object. Optical colours suggest late-spectral type, or counterpart is behind substantial foreground extinction.

9a: Brighter (eastern) component of a double star. Variability is difficult to assess because the object is near the edge of the plate. From time to time waves with a length of about one month. The X-ray emission is extremely soft, suggesting a small distance. Not detected in a 3.6 ksec Swift XRT pointing; the upper limit is consistent with the ROSAT flux, given the very soft spectrum.

10: Star cluster NGC 6823. 10b seems to be just visible on some plates though the USNO magnitude is about 1.5 mag fainter than the plate limit. Nevertheless, the variability could not be definitively confirmed. 10g is identical with Hoag 9, a B0.5V star (Massey et al. 1995) which is unlikely the counterpart due to the hard X-ray spectrum and too large f_X/f_{opt} for a




Fig. 22. Blow-up of the light curve of Sge007a (Fig. 41b) showing pronounced variability.




Fig. 23. Blow-up of the light curve of Sge016a (Fig. 41b) showing pronounced variability.

B star and the large distance from the X-ray position. The Swift XRT position suggests *10e* to be the counterpart.

11a: Not tested for variability. The G spectral type and the f_X/f_{opt} ratio are consistent with identification as counterpart. This is supported by the Swift XRT source position.

12a,b: On Sonneberg plates difficult because of blending. The components of the double star 12a cannot be clearly separated on Sonneberg plates. Therefore, the investigation of the combined light curve can be misleading. Thus, it cannot be ruled out that some observed flares of about 0.5 mag amplitude are only spurious, and, if real, no assignment can be made to one or the other component. The Swift XRT position suggests the south-western component of *12a* as the counterpart.

13: There is little doubt on *13a* being the counterpart which is also verified by the Swift/XRT position as this object is contained in the pointing on Sge014.

14: The Swift XRT position identifies *14a* as the counterpart.

15a: The G spectral type and the f_X/f_{opt} ratio are consistent with identification as counterpart. The X-ray source is not detected in two Swift/XRT pointings on 2007 Nov. 29 (2163 ksec) and 2008 Feb. 29 (1675 sec); the upper limits are consistent with the brightness of the ROSAT source, given its soft spectrum.

16a: The optical brightness varies quickly with small amplitude between 12.9 mag and 13.3 mag. There seem to exist also brightenings or flares (see Fig. 23 and 41b): J.D. 243 7936.471 (25), B = 12.9 mag, 8530.480 (22), B = 12.8 mag, 9238.585 (23), 12.2 mag (in parentheses: exposure time in min.). The X-ray source exhibits rather strong intensity, with practically no neutral hydrogen absorption, thus it must be at a rather small distance. The X-ray light curve shows a bright (2.2 ± 0.3 cts/s) flare during one scan, with levels of 0.25 ± 0.13 cts/s and 0.31 ± 0.17 cts/s before and after (96 min apart). This GSC star has already been proposed as counterpart of this ROSAT source (Motch et al. 1998), and is classified as high-proper motion star in Ivanov (2002).

17a,b: These objects are at no time brighter than the plate limit 17.5 mag. The Swift XRT position identifies *17c* as the counterpart which is difficult to measure on Sonneberg plates due to blending.

18a: Brighter component of a double star. Difficult to treat because of small amplitude and blend. The colour indicates that it may be a star of about F or G type. Minima of short dura-

tion seem to be indicated, which could be interpreted as eclipse minima of an RS CVn star, if real.




Fig. 24. Blow-up of the light curve of Sge018a (Fig. 41b) showing pronounced variability.

19: Central star of planetary nebula M27 $\equiv$ NGC 6853 (Dumbbell nebula). Supersoft X-ray spectrum (Kreysing et al. 1992).

20a: Soft X-ray spectrum consistent with identification with this bright G star.

21a: Discovered already by *Einstein* Observatory observations, and identified by Nousek et al. (1984). Lightcurve published by Fuhrmann (1984).

22a: Double star. *22b*: Short period eclipsing star, may be of RS CVn type. Observed minima (in parentheses B magnitude): J.D. 243 0848.512 (15.3), 6807.367 (15.2), 8268.493 (15.3), 8296.353 (15.2), 8323.332 (15.2), 244 4116.430 (15.2), 7848.262 (15.3), 8894.371 (15.2) (Fig. 25). We did not succeed in finding a period.

23a: Rapid magnitude changes, mostly between 13.5 and 13.8 mag, occasionally a little brighter (13.4) (Fig. 26). There is a single deep minimum (14.0 mag, J.D. 243 3561). Sometimes




Fig. 25. Blow-up of the light curve of Sge022b (Fig. 41b) showing pronounced variability.

brightness changes within some hours. Probably chromospherically active star.




Fig. 26. Blow-up of the light curve of Sge023a (Fig. 41b) showing pronounced variability.

24a: Discovered in X-rays with *Einstein* Observatory observations, it has been originally interpreted by Takalo & Nousek (1985) as an old nova, but based on higher-quality optical spectra, Shara et al. (1990) identifies it as starburst galaxy at $z=0.029$. Light changes on Sonnerberg plates already reported by Andronov (1988), with outbursts up to 1.7 mag amplitude at several occasions. IRAS data are given by Harrison & Gehrz (1991). The X-ray hardness ratio as well as the optical colours are consistent with a source experiencing the full galactic absorbing column.

25: Not detected in a 6.5 ksec Swift XRT pointing on 2008 Feb. 28/29; the upper limit of <0.001 cts/s is not consistent with the ROSAT flux despite the very soft spectrum, but implies about a factor 3 variability.

26a: Relatively hard X-ray spectrum is inconsistent with G star identification unless foreground extinction is considerable. However, Swift XRT position clearly favours 26a as counterpart. Possibly interacting binary?

28: For object α , the NSV catalogue (Kholopov et al. 1982) indicates that the variability of this star is doubtful. But the coincidence with a ROSAT source might suggest that it indeed may be variable with a small photoelectrically measurable amplitude, possibly a chromospherically active star. The position as derived from a pointed ROSAT HRI observation excludes the other two objects 28 b,c as counterpart candidates. 28a is in the list of double stars by Jevers & Vasilewskis (1978).

29a: Possibly variable at the detection limit of photographic plates. Amplitude smaller than 0.2 mag. Chromospherically active star? The Swift XRT position clearly favours 29a as counterpart.

30a: Relatively hard X-ray spectrum is inconsistent with G star identification. Furthermore, this G star is the brightest member of a blend of several fainter stars (see Fig. 40b), so other stars

of this blend may be potential counterpart candidates. The Swift XRT position uncertainty prohibits solving this ambiguity.

31a: Irregular variable with an amplitude of only 0.2 mag, therefore very difficult. May be chromospherically active (see also Fig. 27).




Fig. 27. Blow-up of the light curve of Sge031a (Fig. 41b) showing pronounced variability.

32: The 4.5 ksec Swift XRT observation on 2007 Dec. 28 does not detect the ROSAT source. Given the hard (or absorbed) spectrum, this implies an X-ray variability of a factor of ≈ 9 . The optical brightness dispersion of 32b is relatively high, but is not sufficient to confirm variability. However, since the other two optical sources within the ROSAT error circle are constant, we tentatively identify the ROSAT source with 32b.

33a: Flat waves with a length of about 50 days (Fig. 28). In addition slow changes within years. Ten exposures in different spectral regions of the Tautenburg Schmidt telescope could be used: two B plates (JD 243 8004 and 8005 show the object at 14.8 and 14.9 mag, respectively. In the U band the object is about 0.2 mag brighter with one exception at JD 243 8302 (having 15.3 mag), thus implying 0.7 mag amplitude. Whether this is an eclipse minimum (the only one observed) or a plate fault is difficult to decide.




Fig. 28. Blow-up of the light curve of Sge033a (Fig. 41b) showing pronounced variability.

34a: No brightness changes could be found - constant within 0.1 mag. Stars of spectral type A are usually very faint X-ray sources, thus the f_X/f_{opt} ratio is somewhat high to claim the identification with certainty. But the Swift XRT position supports this ID.

36: The 5.4 ksec Swift XRT observation on 2007 Jul. 24 does not detect the ROSAT source. Given the hard (or absorbed) spectrum, this implies an X-ray variability of a factor of ≈ 15 . 36a seems to be variable within 0.1 mag, but uncertain. The Mira star DD Vul is 5.5 arc apart from the ROSAT position, thus not considered as counterpart.

37a: Difficult because of blending by 37b. Brightness changes of small amplitude. The few (23) observations during

the time interval J.D. 242 9130 - 243 1320 show the object always bright (12.6–12.7 mag). In the interval J.D. 243 7570 - 244 1980 rapid variations between 12.7 and 12.9 mag within only a few days with growing tendency to smaller magnitudes. The few (12) exposures between J.D. 244 2370 - 4520 show the object bright again (12.6–12.7 mag), since J.D. 244 5265 mostly weak (12.8–12.9 mag). May be a chromospherically active star according to the yellow colour. 37c is invisible on Sonneberg plates because of blending by 37a and 37b.

38a: The ROSAT HRI position favours object 38a which is listed in USNO-B with $R_1 = 17.25$, $B_2 = 19.41$ and $R_2 = 16.48$. Besides suggesting variability, the colour is very red. 2MASS magnitudes are $J=12.85$, $H=11.85$, $K=11.45$, suggesting substantial extinction. No flares have been observed on Sonneberg plates. The radio source NVSS J194356+211826 (Condon et al. 1998) is consistent with the HRI position, but not with object 38a. The X-ray spectrum is strongly absorbed, (corresponding to $A_V \sim$ several magnitudes depending on the spectral model), consistent with the optical colours. This indicates a very distant source, possibly even an AGN shining through the Milky Way disk.

39: The Swift XRT position is outside the ROSAT error circle, so this may be a different source, though the X-ray intensity is similar to that of the ROSAT source. 39c may be variable with small amplitude (0.2 mag), but very uncertain.

40: A ROSAT HRI observation (6.3 ksec during Apr 22-25, 1998) improves the X-ray coordinates to $\pm 10''$ (1RXH J195305.2+211449), thus supporting the identification with the dwarf nova V405 Vul ($= 40a \equiv$ S 10943); Richter et al. (1998).

42a: Apart from flat waves (15.8–16.0 mag) with a timescale of weeks, occasionally fadings down to 16.4 mag of very short duration are observed, presumably about 1 hour. Because of the slightly blue colour of this object it may be more probably a cataclysmic object than an RS CVn star. Observed fadings (in parentheses B magnitude): J.D.243 1318.380 (16.2), 243 8243.541 (16.5:), 243 8255.393 (16.3), 243 8286.325 (16.2), 243 9686.474 (16.4), 244 1917.430 (16.2), 244 6355.344 (>16.4), 244 7365.493 (16.4:), 244 7763.381 (16.3), 244 9840.560 (16.6:). The XMM Slew Survey source XMMSL1 J200624.8+210709 is consistent with this object.




Fig. 29. Blow-up of the light curve of Sge042a (Fig. 41b) showing pronounced variability.

43a: For the present time only analysed since 1983. From time to time sinusoidal waves with a length of about 40 days and an amplitude of 0.1-0.4 mag. Also long-time brightness variations. The colour is yellow. May be a chromospherically active star.

44a: Though normally far below the plate limit, it is faintly visible on several plates. Therefore probably variable. Type unknown. The reddish colour excludes CV type. On the other hand, the optical behaviour is not quite typical for a chromospherically

active star. The following supposed brightenings can be seen (in parentheses approximate B magnitude): J.D. 242 9429.518 (16.5), 243 0848.512 (16.5), 243 8238.419 (17), 243 9765.381 (17), 244 2369.235 (17), 244 3777.369 (16.5), 244 4116.445 (17), 244 6640.430 (16.5), 244 7411.376 (16.5), 244 9163.470 (16.5), 244 9213 (17).

45b: On Sonneberg plates totally blended by 45a.

46: The X-ray source has faded by a factor of 30 after the all-sky survey: 4 ROSAT HRI pointings in May 1991 (total exposure time of 13.4 ksec) do not detect the source, and a 15.9 ksec PSPC observation in Oct 20–23, 1991 just recovers 8 photons from the source, corresponding to a vignetting-corrected count rate of 0.0005 ± 0.0002 cts/s. The error circle of $20''$ is fully contained within the all-sky survey positional error (see Fig. 40b), but does not help in distinguishing between the counterpart candidates. The Swift/XRT observation in Feb. 2008 also provides only an upper limit. 46b possibly is slightly variable, but difficult because of blending.

48: The X-ray spectrum is very soft, so given the galactic latitude of $b=-3.4^\circ$ the object must be nearby. A ROSAT HRI observation for 7.3 ksec during Apr 18-21, 1998 reveals 27 cts, thus allowing to improve the X-ray position to $\pm 10''$: $19^h53^m10^s.7 +20^\circ43'53''$. (The small count rate as compared to that during the all-sky survey is consistent with the factor 8 smaller sensitivity of the HRI for very soft X-ray spectra). This improved X-ray position hints toward 48h as counterpart. The Swift XRT position with $5''$ accuracy supports this ID. Moreover, the last of the three Swift observations finds the source a factor 5 brighter in X-rays, while the two earlier Swift XRT flux measurements are compatible with the ROSAT flux. USNO-B1 reports B magnitudes of $B_1=17.4$ and $B_2=15.9$, suggesting strong variability. All these properties are consistent with an AM Her variable. 48e: Is not compatible with the ROSAT HRI position, but found to be variable. Not in USNO-A catalogue. Marginally visible only on a few of the best plates. Between J.D. 243 8282 and 8287 rising from 19 mag to 17.5 mag. At J.D. 243 8323 still bright. Further bright at J.D. 243 9765 (18.4 mag), 243 9792 (17.5: mag), 244 1896 (17.3 mag), 244 5489 (17.5 mag), 244 9213 (17.5 mag), 244 9504 (18.4 mag). On POSS print No. 190 at $B=19.5$ mag, on POSS print No. 302 at $B=20.0$ mag.

49a: It is just outside of the $30''$ error circle, but consistent with the RXS X-ray position (Voges et al. 1999) as well as the XMM slew survey source XMMSL1 J200648.6+204155. Optical light curve see Häussler (1975). Might be a symbiotic star.

50a was originally favoured as counterpart due to its f_X/f_{opt} ratio. 50b: On Sonneberg plates fairly blended by 50a. Between J.D. 244 5460 and 5492, it may be a little brighter than at other times, but uncertain. The RXS X-ray position (Voges et al. 1999) favours this star as counterpart which is supported by the Swift XRT measurement.

51a: Seems to be variable with small amplitude, but because of faintness (on most plates invisible) not sure. The Swift XRT observation does not detect this source; the upper limit suggests a fading by about a factor of 8.

52: Baade (1928) and Prager (1934) proposed NSV 12597 to be variable, but Richter & Greiner (1996) could not decide whether or not NSV 12597 = AN 59.1928 is the optical counterpart of the ROSAT source, because the coordinates published in the NSV catalogue (Kholopov 1982) are questionable. In the meantime, Skiff (1997) has identified the southwestern star of the pair of stars 52b as NSV 12597. The coordinates in Table 5b are for the unresolved pair from USNO-A2, while Skiff (1997) reports $RA = 19^h57^m01^s.2$, $Dec. = +20^\circ05'43''$ for

NSV 12597. On Sonneberg plates no variability of 52*b* could be found. However, the possibility cannot be excluded that a small variability, if existing, will be masked because of blending by the northeastern component of this pair of stars. The Swift XRT observation identifies 52*b* as the counterpart of the X-ray source. While the XRT position favours the northeastern component, the southwestern component cannot be excluded.

53: Both objects are constant. Based on the ROSAT data, the identification with HD 350742 is not obvious since one would expect a softer X-ray spectrum for a nearby G0 star. Foreground X-ray absorption is negligible for all spectral models tested, and a power law model gives a significantly better reduced χ^2 than a bremsstrahlung (4.7) or blackbody model (2.6). However, the Swift XRT position clearly favours 53*a* $\equiv$ HD 350742 as the counterpart.

54*a*: Possibly variable between 16.2 and 16.8 mag. Very uncertain because near the plate edge. The Swift XRT position is clearly outside the ROSAT error circle, and coincides with the bright star 54*c*.

55*a*: Difficult because of small amplitude. Usually nearly constant for several years. From time to time fluctuations with small amplitudes up to 0.2 mag and a cycle length of about 1 month. May be of BY Dra type.




Fig. 30. Blow-up of the light curve of Sge055*a* (Fig. 41*b*) showing pronounced variability.

56*a*: May be variable with an amplitude of about 0.1 mag, but very uncertain. The Swift XRT position proves this object to be the counterpart.

57*a*: Very close double star; the Table 5*b* gives the brightness and coordinates of the northern component. The brightness of the southern component is $R = 11.3$ mag, $B = 12.6$ mag, $V = 11.8$ mag. The Mira star RR Sge (2.5) and the suspected variable 63.1928 = NSV 12599 (3') are too far from the ROSAT position to be considered as counterpart candidates. The position of the only bright Swift XRT source coincides with the bright star 57*e* which is 58'' away from the ROSAT position.

58*a*: On most of the astrograph plates heavily blended by 58*b*. Nevertheless it cannot be excluded that it is variable with an amplitude of even about 1 mag, but very questionable. If real, it could be an RS CVn star. 58*b* is a double star, even on POSS prints not fully separated. The Swift XRT position suggests it to be the counterpart.

59: Not detected in the Swift XRT pointing, though the upper limit is consistent with the ROSAT flux given the soft spectrum.

60: The 1.8 ksec Swift XRT observation on 2008 Apr. 23 does not detect this ROSAT source. 60*a* Seems to be variable, but the amplitude is too small to claim this with certainty.

62*b*: The USNO-A2 magnitudes/colours are likely incorrect, possibly due to blending with 62*a*.

63: While 63*a* is constant on Sonneberg plates, the Swift XRT position coincides with 63*b*.

64: This X-ray source is positionally coincident with 1E 1950.8+1844. Magnitudes of 64*e* and 64*f* are from the USNO-B1 catalogue. Though the B magnitude of 64*f* on the POSS is about 18^m, the object seems to be marginally visible (about 17 mag) on some Sonneberg plates. Nevertheless its variability is questionable. The Swift/XRT position clearly coincides with 64*g* $\equiv$ TYC 1624-1644-1, with a count rate which is compatible to that of the ROSAT source.

65: The Swift XRT position suggests 65*a* as counterpart. Variability of 65*b* with an amplitude near 0.1 mag cannot be fully excluded. QY Sge is more than 5' off the ROSAT position, and thus not considered to be the counterpart.

66*a*: Difficult because of blending by 66*b* for which magnitudes are taken from the USNO-B1 catalogue. Violent light changes, particularly before 1970 (see Fig. 31). From time to time constant over many years at about $B = 13.5$.




Fig. 31. Blow-up of the light curve of Sge066*a* (Fig. 41*b*) showing pronounced variability.

67*a*: See photoelectric catalogue of M stars by Neckel (1974). May be the correct counterpart of the ROSAT source in spite of the relatively large positional offset of 35''. Spectral type M2.5II-III + B9V according to GCVS. Probably a symbiotic star.

68: The Swift XRT position clearly favours 68*f* as counterpart.

69: The 4.1 ksec Swift XRT observation on 2007 Jul. 18 did not detect this source; the upper limit is a factor 3 below the ROSAT flux.

70: Though the USNO-A2 brightness of 70*c* is $B = 18.4$, the object seems to be sometimes just visible on Sonneberg plates. Therefore, variability is possible. The Swift/XRT position clearly coincides with 70*d* with a count rate which is compatible to that of the ROSAT source.

71: Object 71*b* is heavily blended by a star with $B = 18.0$. It appears on Sonneberg plates as one object. At first glance it seems to be variable, but being at the very plate limit, this is questionable. The Swift XRT position clearly favours 71*c* as counterpart. This object is listed as radio source MRC 2002+182 and candidate gravitational lens (King et al. 1999).

72: The Swift XRT position suggests 72*a* as counterpart.

73: The 3.0 ksec Swift XRT observation on 2008 Jun. 17 did not detect this source; the upper limit of <0.002 is a factor 15 below the ROSAT flux.

74*a*: Extremely slow brightness changes. In 1938–1944 at $B=12.80$, during 1961–1967 at $B=12.75$, during 1973–1978 at $B=12.70$, during 1979–1993 $B=12.65$, and since 1994 about $B=12.65$ – 12.70 . About 165'' away from the ROSAT position is the Mira star RT Sge, which surely has nothing to do with the X-ray source.

75a: Seems to be variable, but not quite sure because of faintness. The Swift XRT observations do not detect this source, implying a factor of about 4–6 variability.

76a: Is the well known star α Sge.

77: The Swift XRT observation does not detect this source, implying variability by about a factor of 4. The ROSAT spectrum as well as the f_X/f_{opt} ratio suggest 77a as a chromospherically active K/M star counterpart, but other identifications can not be excluded.

78: The 8.0 ksec Swift XRT observation on 2007 Dec. 20/21 did not detect this source; the upper limit of <0.001 is a factor 15 below the ROSAT flux. There is, however, a bright point source at about 1.3 distance, at RA = $20^{\text{h}}15^{\text{m}}21^{\text{s}}64$, Decl. = $+17^{\circ}59'50''.1$ ($\pm 4''.5$), in both pointings. However, its distance is substantially larger than the ROSAT localisation error, so we do not consider this source to be related to the ROSAT source.

79: The 4.3 ksec Swift XRT observation on 2007 Jul. 23 did not detect this source; the upper limit of <0.002 is a factor 10 below the ROSAT flux. We detect three other sources within $3'$ of the ROSAT position, all with count rates below the ROSAT all-sky survey limit. 79a seems to be variable between 16.4 and 16.9 mag, but very uncertain, because the object is near the plate edge, and thus the plate limit.

80a: See for example Vogt & Bateson (1982). WZ Sge is the obvious counterpart which is nicely confirmed by the XMM position (see small error circle in Fig. 40b). (This is the only XMM observation in the Sge field.) IRAS data are given by Harrison & Gehrz (1991).

81a: In spite of the small amplitude obviously really variable. Long waves with a cycle length of several 100 days. Due to variability preferred as optical identification. The Swift XRT position supports this ID. This is $38''$ from the RXS position.




Fig. 32. Blow-up of the light curve of Sge081a (Fig. 41b) showing pronounced variability.

82c: The dispersion of the magnitudes is relatively high, variability cannot be fully excluded. The Swift XRT position falls near to it, but is slightly off. The probable Algol star DO Sge is about $3'$ apart from the ROSAT position, too far to be considered a counterpart candidate.

83a: The published USNO-A2 R, B, V magnitudes (non-simultaneous!) are marked as “probably wrong”. The K2 spectrum as listed in Simbad can be traced back to the AGK3 catalogue, but Cannon & Mayall (1949) list it as M0 star with $V=7.7$ mag! The USNO-B1 magnitudes are $B1=9.38$, $R1=6.90$, $B2=8.44$, and $R2=6.83$. With a photometric accuracy of better than 0.3 mag, this might suggest variability (the object is too bright to check variability on our astrograph plates). The f_X/f_{opt} of HD 190720 is compatible with an K or M star to be the optical counterpart. The Swift XRT position supports this ID.

84: The two Swift XRT observations in 2007 did not detect this source; the upper limit of <0.003 is a factor ≈ 2 below

the ROSAT flux when accounting for the soft spectrum. 84f is a double star.

85a: The magnitudes show waves with a length of some 100 days and an amplitude smaller than 0.1 mag. The magnitude changed from 12.3 to 12.4 mag between J.D. 242 7570 and 242 9630 to 12.4 to 12.5 later. More rapid changes within several days were observed only from J.D. 243 8200 to 243 8400. Probably BY Dra type.

86a: Light changes in waves of some weeks or months with an amplitude up to 0.5 mag. Superposed are very slow variations. Mean magnitude: J.D. = 242 7500 - 244 0400: brightness fluctuates between 14.2 and 14.6 mag; J.D. = 244 1900 - 244 7850: slowly fading to about 15.0 mag; J.D. = 244 7850 - 255 0300: rising from 15.0 to 14.2 mag (Fig. 41b).

87-91,93,94: SNR 053.6-02.2 = 3C 400.2. This is an extended X-ray source (of order $20'$), and our source detection “finds” 7 spurious X-ray sources, as it was tailored for finding point sources. The stars in these regions were not tested for variability, and no finding charts are given. See Saken et al. (1995) on the ROSAT data of 3C 400.2.

92: Objects a–d constant on Sonneberg plates. 92e is invisible on most of the plates, but appears occasionally with clearly different magnitudes (though some of the brighter observations might be artefacts). Because of the relatively blue colour it may be a cataclysmic object. The Swift XRT position is consistent with this source, though we cannot distinguish from 92b. The combined evidence of variability, X-ray emission and f_X/f_{opt} ratio suggest object 92e to be the counterpart.

95: The 2.9 ksec Swift XRT observation on 2007 Dec. 17 did not detect this source; the upper limit of <0.001 is a factor 20 below the ROSAT flux. Both objects 95a,b: are only visible on the best plates. No variability could be found.

96: The Swift XRT observation suggests 96b as counterpart which is much too faint to be visible on Sonneberg plates.

97a: Brightness amplitude photographically not measurable. It may be a chromospherically active star. See also Blanco et al. (1972). The very soft X-ray spectrum as well as f_X/f_{opt} are compatible with the G spectral type.

98a: A pointed ROSAT PSPC observation yields a detection (2RXP J200318.5+170253) with a position which is accurate to $5''$, and thus leaves no doubt that RZ Sge is the counterpart of this X-ray source. For optical properties see for example Vogt & Bateson (1982) and Khruzina & Shugarov (1991). Curiously, Kouzuma et al. (2010) identify this as AGN candidate due to the NIR colours, but in addition also list 98c and two fainter objects just outside the $30''$ error circle as AGN candidates.

99: This source is not detected by a Swift XRT observation, with an upper limit consistent with the ROSAT flux.

100a: Blend of three reddish stars. Because of this blend and the small expected amplitude a photographic search for variability seems to be without chance of success. Therefore not tested for variability. The USNO coordinates and BVR magnitudes are the combined values of this blend.

101b: Extremely blue. On plates of the Sonneberg astrograph blended by 101a which is considered to be a viable counterpart due to the match of spectral type and f_X/f_{opt} .

104a: Brightness changes with small amplitude with a timescale of months or years (Fig. 33). Moreover, three distinct minima of short duration (some hours or perhaps one day) were found. These might be eclipse minima: J.D. 243 7584.308 (14.5 mag), 243 8371.224 (14.3 mag), 243 8371.266 (14.3 mag), 244 7850.238 (14.4 mag). The star may be of RS CVn type.

105a: X-ray spectral hardness, f_X/f_{opt} and optical colour all suggest this to be the counterpart.




Fig. 33. Blow-up of the light curve of Sge104a (Fig. 41b) showing pronounced variability.

109: The Swift XRT observation suggests **109b** as counterpart. No other optical object brighter than 18 mag within a distance of 40'' from the ROSAT source is visible on Sonneberg plates except **109a**, which is visible only on some of the best plates. No brightness fluctuations could be found. Object **109b** coincides with the radio source 4C +16.67 (Douglas et al. 1996), and has been identified by Kouzuma et al. (2010) as AGN candidate due to the NIR colours.

110: The semi-regular variable (sub-type A) star DY Sge is about 6' away from the ROSAT position and not considered to be the counterpart. The Swift XRT position excludes objects **110a,c,d**, but is not accurate enough to distinguish between **110e** which is invisible on all Sonneberg plates, or the blend of four objects **110b**.

111: The Algol star DV Sge is nearly 4' away from the ROSAT position and not considered to be the counterpart. The f_X/f_{opt} ratio suggests **111b** as counterpart, the brightness amplitude of which is constrained to less than 0.1 mag, if variable.

112b,c: On astrograph plates totally blended by **112a**.

113a: Difficult because of faintness. At the limits of detectability, but the variability seems to be sure. Long waves of small amplitude.

114c: Magnitudes are from the USNO-B catalogue. Very blue colour; may be variable, but uncertain due to faintness. **114f:** Difficult because of faintness. On two overlapping POSS plates 276 and 372 (JD 243 3835 and 243 3910) at nearly 17.3 mag. A CV nature is possible because of the positive f_X/f_{opt} ratio. Identification is ambiguous.

115a is constant within ± 0.1 mag, but the Swift XRT position suggests this to be the counterpart. **115b** is disturbed by **115a**. **115b** itself consists of two very close objects of similar brightness, on Sonneberg astrograph plates only the combined light can be seen. Therefore it cannot be decided which of the two objects is variable. Magnitude changes with a time scale of several weeks or even years (Fig. 34). Because of the blue colour possibly CV type. The USNO magnitude is most likely only for the western component.




Fig. 34. Blow-up of the light curve of Sge115b (Fig. 41b) showing pronounced variability.

116a: See Raveendran (1984) and Richter & Greiner (2000).

117: Dispersion of magnitudes of object **117d** relatively high. Small amplitude variability cannot be excluded. The Swift XRT observation marginally detects the source, and suggests **117a** to be the counterpart, consistent with the M2.5 spectral classification and the f_X/f_{opt} ratio.

119a: Additional plates of the 140 mm triplet and of the neighbouring field γ Aql could be used. Irregular light changes of small amplitude.




Fig. 35. Blow-up of the light curve of Sge119a (Fig. 41b) showing pronounced variability.

120a: Though usually fainter than the plate limit, this very blue object can be seen on some plates: Variable between about 16.5–18.0 mag. More frequently faint than bright. The Swift XRT position clearly identifies this as counterpart of the ROSAT source. There is about 50% X-ray variability between the two Swift observations spaced by 10 days, where the second observation shows the source about a factor 3 brighter than the ROSAT flux. There is also a XMM Slew survey source XMMSL1 J201359.1+154419 within 8'' which is likely the same source. The blue optical colour, the hard X-ray spectrum and the clear optical and X-ray variability suggest an intermediate polar type CV. Alternatively, Kouzuma et al. (2010) identify it as AGN candidate due to the NIR colours from 2MASS. However, the blue optical colour together with a galactic foreground $A_V = 0.52$ mag make an AGN shining through the disk of our Galaxy unlikely.

122a: Spotted star according to Hall & Henry (1992).

123a: Variations within weeks or months. Also more rapid brightness changes within days. Plates of the 170 mm triplet and of the neighbouring field γ Aql could also be used. **123b,c:** Just visible on good plates. No brightness variations could be found. The Swift/XRT observation targeted on **124** clearly detects this X-ray source at 0.027 cts/s, about a factor 3 brighter than during the RASS. The derived X-ray position favours **123d** or **e**. Both objects are invisible on Sonneberg plates, so nothing can be said about optical variability. Kouzuma et al. (2010) identify **123e** as AGN candidate.




Fig. 36. Blow-up of the light curve of Sge123a (Fig. 41b) showing pronounced variability.

Table 7. Summary of identification statistics, separated in secure, suggested (sugg.) and no identification (ID), separately for genuine RASS catalogue sources and our additional RASS sources (50%/97% more sources for the Com/Sge fields). The numbers in parentheses are the fractions for each group.

Field	RASS catalogue			additional RASS source			Total (as in Tables 1a, 1b)		
	secure ID	sugg. ID	no ID	secure ID	sugg. ID	no ID	secure ID	sugg. ID	no ID
Com	122 (77%)	23 (14%)	14 (9%)	54 (68%)	14 (18%)	11 (14%)	176 (74%)	37 (15%)	25 (11%)
Sge	45 (70%)	11 (17%)	8 (13%)	35 (51%)	19 (28%)	14 (21%)	80 (61%)	30 (23%)	22 (17%)

124a: Additional plates of the 170 mm triplet and of the neighbouring field γ Aql could also be used. Relatively high dispersion of magnitudes, nevertheless variability uncertain. The f_X/f_{opt} ratio supports it to be the counterpart, as does the Swift XRT position. The Mira star V 1019 Aql is more than 5' distant from the ROSAT position and not considered as counterpart candidate.

126a: Identified by Kouzuma et al. (2010) as AGN candidate due to NIR colours. **126b:** Difficult because at the extreme edge of the plates. Some fadings seem to occur from a relatively constant brightness level. Eclipsing variable of RS CVn type? Observed minima: J.D. 243 7560.524 (15.8: mag), 8371.224 (15.9: mag), 9385.370 (15.8 mag), 9792.287 (16.1 mag), 244 6683.403 (15.9: mag), 6763.219 (15.9: mag), 9213.422 (15.9: mag).

127b: Distinctly beyond the plate limit, but on two occasions faintly visible: J.D. 242 9846.473 (15.6: mag), 244 2713.261 (16.7 mag). It is difficult to say whether these are real brightenings or artefacts. However, the Swift XRT position is consistent with **127b** being the counterpart, lending more credit to this optical behaviour.

128: Both, **128a** and **128b** are constant on Sonneberg plates, and not considered as potential counterparts. The Swift XRT position suggests **128c** instead.

129a: Any light variations of this object are surely smaller than 0.03 mag.

130b: Though the brightness is clearly beyond the plate limit, this object is indicated on five of about 250 plates: J.D. 242 9844.531 (16.9: mag), 242 9845.502 (16.6 mag), 243 9702.411 (16.6: mag), 244 8096.490 (16.7), 244 8186.348 (16.3: mag). Therefore it is variable with an amplitude of about 1 mag. According to the blue colour it may be a cataclysmic variable.

131: The slowly variable star NSV 12500 is too far from the ROSAT position (11') to be the optical counterpart. **131a:** Small brightness variations within the limits 11.3–11.4 mag seem to be indicated. In single cases small flares (11.1 mag) which last about 1–2 days. Since the f_X/f_{opt} ratio is inconsistent with a CV, this is likely a chromospherically active star.

132: The Swift XRT position clearly favours **132b** as counterpart which is invisible on Sonneberg plates.

4. Results and Discussion

4.1. Identification results and global statistics

We are able to identify $\sim 70\%$ of the X-ray sources in the two selected fields, and suggest likely counterparts for another 20% of objects (see Table 7). This is a surprisingly high number given that the expectation for the optical brightness of our faintest X-ray sources through the f_X/f_{opt} ratio is $B \sim 22^m$ (see Fig. 2).

As mentioned in Sect. 2, our source lists for each field are larger than the published 1RXS catalogue (Voges et al. 1999) due to primarily the slightly lower S/N ratio required for the X-ray detection (Voges et al. 1999). This results in 50%/97% more sources in the Com/Sge field, respectively. The rate of source identification is very similar for the official vs. this extended source list (see columns 2–4 vs. 5–7 in Table 7), suggesting that the number of spurious X-ray sources in our lists is still negligible. This is also supported by the distribution of optical to X-ray position offsets (Fig. 38) which does not show any systematic errors (see below).

A breakdown of the identified sources is given in Table 8, showing the dominance of AGN in the high-galactic field, and bright and/or chromospherically active stars in the low-galactic field. Most probably, the majority of optical counterparts in the "empty" fields (as well as some of the faint galaxies) are still undiscovered AGN. Therefore, nearly 2/3 of all the ROSAT sources in the 26 Com field may be of extragalactic origin.

Somewhat surprisingly, we find 9 clusters of galaxies and another 8 candidates, despite none of the X-ray sources was detected as extended. If true, this implies a cluster content in the RASS about a factor 2 more than hitherto found (e.g. Böhringer et al. 2013). This has interesting implications for the upcoming eROSITA survey for which the expected number of clusters (100,000; Chon & Böhringer 2013) is based on the RASS count.

The Hamburg/RASS catalogue with nearly 4400 optical identifications of northern hemisphere (Decl. $> 30^\circ$), high-galactic latitude ($> 30^\circ$) X-ray sources yielded as largest source population $\approx 42\%$ AGN, followed by $\approx 31\%$ stellar counterparts, whereas galaxies and cluster of galaxies comprised only $\approx 4\%$ and $\approx 5\%$, respectively (Zickgraf et al. 2003). Another large effort was the identification of a complete flux-limited sample of northern RASS sources in six sub-fields covering a total of 685 square degrees, which resulted in the spectroscopic identification of 662 out of 674 sources (Zickgraf et al. 1997, Appenzeller et al. 1998, Krautter et al. 1999). These authors find relative source class fractions of 42.1% AGN, 40.7% stars, 11.6% cluster of galaxies, and 3.9% galaxies. The ROSAT galactic plane survey (Motch et al. 1998) achieved identification of 76 out of 93 sources, among them 40 bright/active stars, 25 AGN and 6




Fig. 37. Blow-up of the light curve of Sge131a (Fig. 41b) showing pronounced variability.


Table 8. Summary of the secure and probable optical counterparts of ROSAT sources.

Type	26 Com	γ Sge	remarks
variable AGN and AGN?	34	1	1
constant AGN and AGN?	88	3	2
cluster of galaxies	17	0	
CA and CA?	18	25	3
LB and SRB	2	1	4
bright stars	31	34	5
UV and UV?	6	1	6
eruptive binaries	2	14	7
magnetic stars (Sp. Am)	1	0	
DA	2	0	
PN	0	1	
supernova remnant	0	1	8
other	11	5	
not identified	25	21	9
not classified	1	19	10

- 1: Some of them may be eruptive binaries.
- 2: Variability photographically not established (too faint, blends or too small amplitudes).
- 3: Chromospherically active stars (RS CVn and BY Dra).
- 4: Slow irregular and semiregular variables of late spectral type (see GCVS, Kholopov 1985), some of which may possibly also be related to the chromospherically active stars.
- 5: Mostly overexposed on the photographic plates, and thus not investigated for variability. Because of the scarcity of bright stars as counterparts of ROSAT sources, these matches are likely real associations (see Fig. 39). These stars are therefore with high probability chromospherically active stars.
- 6: Flare stars of UV Cet type.
- 7: Mainly cataclysmic variables (CVs), but also X-ray binaries of AM Her type.
- 8: 7 X-ray sources close together, since source detection algorithm was optimized for point sources.
- 9: Out of the 25 unclassified objects in the Coma field (dash in the last column of Table 1a), the error circle of 9 objects is empty on the DSS2 (about $B \sim 20$ th mag), and for another 17 X-ray sources (totalling 26), no optical object is visible on the Sonneberg astrograph plates (about $B \sim 17$ th mag).
- 10: Positional coincidence implies secure correlation, but source type remains open.

cataclysmic variables (this survey selected according to X-ray brightness and various hardness ratio criteria, so did not aim at completeness in a given area). These relative fractions of source types are largely consistent with our results (Table 8), though some differences show up. The low fraction of AGN in the two high-galactic latitude identification programs might be due to the bias against fainter counterparts for which objective prism spectra were either not available or inconclusive – this would rise the fraction of bright-vs.-faint counterparts, and thus the ratio of stellar-vs.-extragalactic objects. The nearly complete lack of AGN in our low-galactic latitude field as compared to the nearly 20% fraction in Motch et al. (1998) is most likely due to their survey covering galactic latitudes up to $\pm 20^\circ$, and our optical material not being deep enough to recover absorbed AGN behind the Milky Way disk.

Fig. 38 shows the histogram of the distances of the secure optical counterparts to their respective centroid ROSAT X-ray position for the Coma field. The distribution for the Sge field looks very similar. The mean offset is $15''$, and 90% of all X-ray sources have their counterparts within $30''$.

**Fig. 38.** Distribution of positional offsets between centroid X-ray and optical position of the identified objects in the Com field (solid line), and the subset of 26 sources for which Swift/XRT positions have been obtained. The dashed line shows the expected Rayleigh distribution, with a best-fit of $\sigma = 10''.5$.

In the Coma field, 5 of our identifications have offsets relative to the X-ray centroid of more than $40''$, and constitute the very tail of the offset distribution in Fig. 38: these are Com025b, 82a, 99a, 147a, and 212a. We have double-checked these for possible mis-identifications, but find no systematic problem. Two of these sources (99a, 147a) are identified with bright stars, two others (25b, 212a) are driven by Swift/XRT positions, and one is a SDSS-III classified QSO. None of 82a, 99a, 147a have any other object brighter than $R \sim 20$ mag in their error circles. Only for two objects, potential concerns could be risen: for 147a the f_X/f_{opt} ratio is on the low side if the K0 stellar classification is correct, and for 212a the Swift/XRT flux is a factor 4 below the RASS rate, so would have been undetected in RASS, and thus being possibly a different source if the RASS source has faded by a factor of 10.

4.2. Variability as a classification criterion

In our study, variability has been used as an additional criterion of an object to be an optical counterpart of a ROSAT point source, besides the classical F_X/F_{opt} ratio. Obviously, this variability criterion has not been applied to extended sources, like supernova remnant SNR 053.6-02.2 and clusters of galaxies.

For the Sonneberg plate material used here, the best magnitude interval to detect variables on astrograph plates is about 13–15 mag. In this magnitude range we can discover brightness amplitudes of 0.2 mag or even smaller. At fainter magnitudes, near the plate limit at about 17 mag, no variations smaller than about 0.5 mag can be found.

From Tables 1a, 1b and 5a, 5b we can deduce the following (see Table 8): 40 (30%) of the supposed optical counterparts in γ Sge and 61 (26%) in 26 Com were found to be variable. Some of these variables are already known, most are newly detected, and have therefore been assigned a S number (for new Sonneberg variables) in column 5 of Table 5a, 5b. Moreover, during the search for optical variability a total of 5 variable objects were found which seem not to be the counterparts of ROSAT sources, either because there are better candidates for optical identification or because of the relatively large spatial separation

to the ROSAT source: Com 98a (E), Com 101a (CV?), Com 109e (AGN), Sge 48e (CV?), and Sge 115b (CV?). Ignoring these, but considering that our sensitivity to detect variability degrades rapidly beyond $B \sim 16.5$ mag, we find that 50% (Com) and 39% (Sge) of the identified sources are variable, respectively. If we also take into consideration that most HD stars correlating with ROSAT sources and have not been tested for variability may be variables of small amplitude, a good fraction of all of the optical counterparts could be variable. Exceptions are objects classified "C" (constant) which are either spurious counterparts, or their light amplitudes are too small to be detected photographically, or they are real counterparts without expected optical brightness changes (e.g. Seyfert-2 galaxies, hot white dwarfs, brown dwarfs, see e.g. Neuhäuser & Comeron 1998).

Optical variability is, not unexpectedly, tightly correlated with X-ray emission, and thus our new method for optically identifying X-ray sources is demonstrated to be feasible. Given the large number of optical plates used, this method was likely not more efficient than e.g. optical spectroscopy. However, it required no proposal writing and no telescope time, but just access to publicly available archival data. We have shown that an unequivocal optical identification of ROSAT sources is not possible in every case, but for a large fraction of objects (65–80%) we can estimate for each object the reliability of a supposed identification by including optical variability as additional criterion. It remains, however, true that optical spectroscopy will remain a highly demanded asset, in particular to determine redshifts or sub-types of the AGN-population: only then important physical parameters like the luminosity, binarity or presence of an accretion disk can be deduced.

Yet, optical variability can serve as an efficient tool in pre-selecting objects from specific source classes. Together with the optical colour as e.g. provided by the USNO catalogue, the f_X/f_{opt} ratio and details of the optical light curve often provides already sufficiently accurate clues on the source type. This is proven by the (even for us surprising) little impact which the SDSS-III DR9 spectra had (due to their late availability relative to the completion of this work) on our identifications. Thus, optical variability can be an efficient substitute for optical spectroscopy in the early identification process of many thousands or million of objects. This is particularly true as long as no systematic spectroscopy of all objects on the sky down to 20th mag is available – which might well apply even for the first years of *eROSITA* operation.

4.3. Impact of variability as reliable identification criterion

With the about 200.000 X-ray sources from the ROSAT mission, massive identification programs have been employed based on spatial correlations with other large catalogues. A very early attempt was a correlation with the Hamburg objective prism survey which provided identifications for nearly 4400 ROSAT Bright Source Catalog sources (Zickgraf et al. 2003). Another correlation was done with the IRAS Faint Source Survey, the NRAO VLA Sky Survey, the FIRST Radio Survey, the Two Micron All-sky Survey and the APM and COSMOS catalogues of digitized Schmidt plates, revealing over 1500 sources with an identification reliability greater than 90% (Lonsdale et al. 2000). And finally the correlation with the Sloan Digital Sky Survey (SDSS, data release 7) results in approximately 10.000 confirmed quasars and other active galactic nuclei that are likely RASS identifications (Binder et al. 2011).

A different identification method used multicolour CCD imaging with the goal of a classification via the spectral energy

distribution. For one particular medium-deep ROSAT pointing, 149 of the 156 detected objects within the error circles of 75 X-ray sources were classified, but despite a 15-filter dataset per object, no identifications are proposed (Zhang et al. 2004).

While these examples are far from exhaustive, they show the size of the identifications of the various programs. With about $\lesssim 20.000$ ROSAT sources identified, this also demonstrates the corresponding challenge in getting identifications for the remaining ≈ 180.000 ROSAT sources.

As such, the herewith proposed identification of 256 ROSAT sources is just another little step. However, our main goal was to prove optical variability as a new, efficient classification criterion, and thus pave the way for larger projects of this kind, both for ROSAT as well as the upcoming *eROSITA* all-sky survey (Predehl et al. 2006).

Despite the better positional accuracy of $1''$ – $5''$ of the presently used X-ray missions *Chandra*, *XMM-Newton* and *Swift/XRT*, the ROSAT survey is still considered a unique resource when it comes to large areas, or all-sky coverage. Ignoring the ~ 180.000 X-ray sources because of missing identifications might be a superficial decision. Our suggestion to use optical variability as an efficient criterium could be used in the near-term future in a two-fold manner: (i) the digitization of large photographic plate collections, e.g. at Harvard (Grindlay et al. 2009, Grindlay et al. 2013) or Sonneberg Observatories (Kroll 2009) is well advanced, and is going to provide a census of the optical variability of the sky over the last 120 years down to limiting magnitudes of order 14–16 mag. (ii) Digital sky surveys such as the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS; Tonry et al. 2012), the Palomar Transient Factory (PTF; Kulkarni 2012), the Catalina Real-Time Transient Survey (CRTS; Drake et al. 2012), the La Silla-QUEST variability Survey (LSQ; Hadjijska et al. 2012), or the Sloan Digital Sky Survey (SDSS; strip 82, MacLeod et al. 2012) already provide multi-epoch observations in various filter and cadence combinations, and promise to be a great resource in a systematic identification program of the RASS. The recent cross-correlation of periodic variables from the ASAS survey with bright ROSAT sources (Kiraga & Stepień 2013) is a promising first step. In the long-term, the Large Synoptic Survey Telescope (LSST; Ivezić et al. 2008) will provide the ultimate database in 5 filters with 3-day cadence to a depth which is fully sufficient for the identification of the majority of RASS sources.

The upcoming *eROSITA* all-sky survey (Predehl et al. 2006), a 4-yr survey providing 8 consecutive all-sky surveys in the 0.3–10 keV band, will have source position uncertainties similar to that of the ROSAT survey. While large (and expensive) multi-object fiber spectrographs are being planned and build, such as BigBOSS (Schlegel et al. 2010), MOONS (Cirasuolo et al. 2012), 4MOST (de Jong et al. 2012) or WEAVE (Dalton et al. 2012), it is well conceivable that these spectroscopic surveys will not be available during the first years of the *eROSITA* operation. Identification programs using optical variability could be a valuable approach.

4.4. Comments on specific source classes

4.4.1. Semiregular and irregular variables

Red semiregular and irregular variables are usually not X-ray sources. On the other hand, three such objects (FN Com, HH Sge and δ Sge) have a very small distance to the ROSAT position. Possibly, they are members of X-ray radiating symbiotic binaries. This remains to be checked spectroscopically.




Fig. 39. Number of bright stars in dependence of the offset from the X-ray centroid position (D_1 in $''$) for three different brightness intervals as labeled in the three panels for the Com (filled hexagon) and Sge (open triangles) fields, respectively. The overdensity within the $30''$ X-ray error circle. The percentage of real identifications (within $27''$) is larger than 90% (Coma) and 65% (Sge) fields, respectively.

4.4.2. Chromospherically active and flare stars

Chromospherically active (CA) stars and UV Ceti stars (flare stars) are known to exist not only isolated, but also in T associations and star clusters, representing a very young stellar population. According to Gershberg and Shakhovskaya (1974) and Rodono (1980), flare stars and chromospherically active stars are related and both should therefore be young. Thus, the large number of these objects in our Com field is not a feature of stellar population, but it reflects the fact that a large area of our Coma Berenices field is covered by the extended, young open cluster Melotte 111. Based on a literature comparison (Randich et al. 1996, Odenkirchen et al. 1998, Abad et al. 1999), we assume that some of our objects are cluster members. In order to verify this, proper motion or radial velocity measurements are necessary.

4.4.3. Bright stars

The above tables 5a,5b contain many bright stars ($B \lesssim 13.5$ mag) within or near the ROSAT X-ray positions. On plates of the Sonneberg field patrol those objects are overexposed. In the few cases, in which variability measurements and/or optical spec-

Table 9. Summary of known and candidate cataclysmic variables in our two fields.

Name	Src-ID	Type	B (mag) (USNO)	Comment
IR Com	Com141a	SU	13.4	
J1238+1946	–	WD	17.4	1
S10980 Com	Com101a		18.5	2
DO Vul	–	SU	20.7	
J1953+1859	–	SU	20.4	
V458 Vul	–	Nova		3
QQ Vul	Sge21a	AM	14.4	
V405 Vul	Sge40a	SU	19.0	
WZ Sge	Sge80a	SU	15.0	
HM Sge	Sge102a	Nova		
S11015 Vul	Sge33a		14.8	
S11017 Vul	Sge42a		15.5	
S11020 Vul	Sge48e		18.6	2
	Sge48h	AM	17.2	
S11035 Vul	Sge92e		17.8	
S11029 Vul	Sge114f		16.3	4
S11030 Vul	Sge115b		14.6	2
S11038 Aql	Sge120a	IP	17.2	
S11033 Aql	Sge130b		17.6	

¹⁾ WD = white dwarf. Double-degenerate binary (Brown et al. 2013).

²⁾ Is not the counterpart of the X-ray source.

³⁾ ROSAT observation was prior to outburst of 2007.

⁴⁾ Is possibly not the counterpart of the X-ray source.

troscopy are available, these stars are mostly chromospherically active stars. While an unequivocal identification is not possible in every case, a statistical estimate of the correlation and of the completeness and reliability of the samples can be made (Richter 1975). Separately for the Coma and Sge fields, we have subdivided our sample into three sub-groups of apparent magnitude $B < 9$ mag, $9 \text{ mag} < B < 12$ mag, and $12 \text{ mag} < B < 13.5$ mag, and determined the number of bright stars in rings of increasing distance D_1 (in $''$) around the X-ray centroid. The resulting numbers (Fig. 39) clearly show an excess of bright stars for distances smaller than about $27''$ (corresponding to $D^2 < 750$). The corresponding total number of bright star counterparts are 37 (Coma) and 54 (Sge). This is surprisingly close to the number of proposed counterparts as given in Table 8, the slight difference being caused by the fact that some objects with already known variability are present in the above statistics, but not in Table 8.

4.4.4. Cataclysmic variables

In the Coma field, one of the two known cataclysmic variables (CV) is detected in the RASS, and one new possible CV is discovered. In the Sge field, 4 of the 7 known CVs are detected in RASS, and 9 new possible CVs are discovered; see Table 9 for a summary of these sources. The non-detections of known CVs are readily explained, as none is expected to provide an X-ray flux large enough for the RASS sensitivity.

While we miss a more detailed sub-classification of these new possible CVs, we can at least make statistical estimates of the population density of various classes of objects at low and high galactic latitudes. In spite of their low luminosity ($M = +5...+12$ mag at minimum), the high concentration of CVs towards low galactic latitudes is obvious, indicating a young (disk) stellar population. This nicely corresponds to earlier results of Richter (1968) who derived a logarithmic density gradient perpendicular to the galactic plane as large as $\delta \log v / \delta z = 3.9$.

The identifications in our two fields, if all confirmed, would double the number of cataclysmic variables in the fields. Given the observed optical brightness of the new CV candidates, as well as their optical colours and the X-ray hardness ratios, for none of these sources a distance larger than ~ 1 kpc would be required to model the observed properties. This is particularly true for Sge48 $\equiv$ S11020 Vul and Sge120a $\equiv$ S11038 Aql. For the latter, WISE photometry suggests a distance of about 350 pc, while for the former an even smaller distance is mandatory due to the extremely soft, i.e. unabsorbed, X-ray spectrum. Thus, despite the lower X-ray luminosity both the optical magnitude range as well as the covered distance/volume are very similar to that used by Pretorius et al. (2013) when deriving the space density of magnetic CVs. While we are missing the orbital periods and more accurate distance estimates, the present results indicate that the space density of CVs might be a factor 2 larger than derived in Pretorius et al. (2013). This prospect may motivate more detailed studies of these new CV candidates in the future.

4.4.5. AGN variability

The field γ Sge is basically free of AGN and other galaxies, except Sge24a and possibly Sge38a $\equiv$ RXS J194356.1+211731, see text. The following therefore is deduced exclusively from the high-galactic latitude Coma field.

We start by noting that the X-ray and optical measurements were not taken simultaneously, but years to decades apart. Long-term flux variations of a factor of ten or even more are nothing unusual. Optical variability studies have been moving from monitoring a few dozen sources over weeks to years (e.g. Cristiani et al. 1990) to the statistical study of thousands of sources over at most a few years (e.g. Bauer et al. 2009, Zuo et al. 2012). Here, we have analysed the optical variability of two dozen AGN over timescales of 50 years, which provides a less biased census of the more rare, large-amplitude variations. These follow the anti-correlation with luminosity as has been found in several previous studies (e.g. Cristiani et al. 1990, Vanden Berk et al. 2004, Zuo et al. 2012).

5. Outlook - the quest for a massive identification program of RASS sources

The X-ray source population of a galaxy provides important clues to the nature of its constituents and their evolution over the lifetime of the galaxy, in particular also by tracing the endpoints of stellar evolution through accreting compact objects. The statistical analysis of X-ray source populations in nearby galaxies, e.g. number density, the X-ray luminosity function of various source types as well as variability patterns, have led to substantial progress in our understanding of the evolution of important underlying physical parameters like metallicity and star formation rate (Gilfanov 2004, Grimm et al. 2003/2005, Townsley et al. 2006, Pietsch 2008, Fabbiano 2010, Mineo et al. 2012). As we are moving from the study of just the most luminous sources ($L_X > 10^{37}$ erg/s) to fainter thresholds, sometimes already as low as a few times 10^{35} erg/s, it becomes increasingly important to understand the variety of sources contributing in this luminosity range. This is where more detailed studies of X-ray sources in our galaxy come into play. Admittedly, luminosity estimates are much more uncertain for galactic sources due to limitations in the distance estimates and the clumpiness of foreground extinction. However, the diversity of spectral and temporal (variability at large, flares/outbursts, duty cycle, pulsed fractions) pa-

rameters are much easier to investigate for galactic sources. All the sources reported here will be among the bright sources of the forthcoming eROSITA survey, allowing to do exactly those spectral and timing studies. Only when having their optical identifications at hand including the multi-wavelength properties can the full power of these methods towards constraining evolutionary models be realized.

Acknowledgements. We thank the 4pi Systeme - Gesellschaft für Astronomie und Informationstechnologie mbH, Sonneberg, Germany, especially P. Kroll and H.-J. Bräuer, for support of this project. We are indebted to the team of the Karl Schwarzschild Observatory Tautenburg for lending us plates from the 2 m Schmidt telescope. We are grateful to S. Nedyalkov (Uni. Bremen) and J. Brunschweiler (TU München) for their thorough help in preparing the finding charts and part of tables 1a,1b and 5a,5b, to S. Zharykov (ONAM Ensenada, Mexico) for the spectra of Com003 and Com007, and M. Salvato for discussion about X-ray properties of AGN. JG thanks N. Gehrels and the Swift team for accepting a substantial fraction of the sources to be observed as “fill-in” targets with the Swift X-ray telescope. The early work (1993–1996) of this project has been supported by funds of the German Bundesministerium für Forschung und Technologie under grant 05 2SO524 (GAR), and FKZ 50 OR 9201 (JG). The ROSAT project was supported by the German Bundesministerium für Forschung und Technologie (BMBF/DLR) and the Max-Planck Society. We appreciate the work of the XMM survey science centre, in particular the regular updates of the EPIC Source and XMM Slew Survey catalogues. This research has made use of data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. This research has also made use of the VizieR catalogue access tool, CDS, Strasbourg, France, and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, University of Cambridge, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

References

- Abad, C., Vicente, B., 1999, A&AS 136, 307
- Andronov I.L., 1988, MVS 11, 84
- Appenzeller I., Thiering I., Zickgraf F.-J. et al. 1998, ApJ Suppl. 117, 319
- Baade W., 1928, Astron. Abhandlungen 9, C1
- Bade N., Engels D., Voges W., et al. 1998, A&AS 127, 145
- Bauer A., Balay C., Coppi P., et al. 2009, ApJ 696, 1241
- Becker R.H., Helfand D.J., White R.L., Gregg M.D., Laurent-Muehleisen S.A., 1997, ApJ 475, 479
- Beuermann K., Thomas H.-C., Reinsch K., et al. 1999, A&A 347, 47
- Binder B.A., Voges W., Margon B. et al. 2011, AAS Meeting #217, #430.13, Bull. AAS vol. 43
- Blanco, C., Catalano, S., Godoli, G., 1972, Mem. Soc. Astron. Italiana 43, 663
- Böhringer H., Chon G., Collins C.A. et al. 2013, A&A 555, A30
- Bräuer H.-J., Häusele I., Löchel K., Polko N., 1999, in *Treasure-Hunting in Astronomical Plate Archives*, P. Kroll, C. la Dous, H.-J. Bräuer, Harry Deutsch: Frankfurt (Main), p. 70
- Brinkmann, W., Siebert, J., Reich, W., Fürst, W., Reich, P., Voges, W., Trümper, J., Wieblinski, R., 1995, A&AS 109, 147
- Brinkmann, W., Siebert, J., Feigelson E.D., et al. 1997, A&A 323, 739
- Brown W.R., Kilic M., Allende Prieto C., Gianninas A., Kenyon S.J., 2013, ApJ 769, 66
- Burrows, D.N. et al. 2005, SSR 120, 165
- Cannon A.J., Mayall M.W., 1949, Ann. Astron. Obs. Harvard Coll., 112, 1-295
- Chen Y., He X.-T., Wu J.-H., Li, Q.-K., Green R.F., Voges W., 2002, AJ 123, 578

- Cheselka, M., Holberg, J.B., Watkins, R., Collins, J., Tweedy, R.W., 1993, AJ 106, 2365
- Chon G., Böhringer H., 2013, AN 334, 478
- Cirasuolo M., Afonso J., Bender R. et al. 2012, SPIE Astron. Instr. and Telescopes Conf., Amsterdam, SPIE 8446, 8446OS
- Condon J.J., Cotton W.D., Greisen E.W., Yin Q.F., 1998, AJ 115, 1693
- Cristiani S., Vio R., Andreani P., 1990, AJ 100, 56
- Cruddace, R.G., Hasinger, G.R., Schmitt, J.H.M.M., 1988, in *Astronomy from Large Databases*, eds. F. Murtagh F. & A. Heck, (Garching, ESO publications) p. 177
- Dalton G., Trager S.C., Abrams D.C. et al. 2012, SPIE Ground-based and Airborne Instr. Conf., SPIE 8446, id. 84460P-12
- de Jong R.S., Bellido-Tirado O., Chiappini C. et al. 2012, SPIE Astron. Instr. and Telescopes Conf., Amsterdam, to appear in SPIE (arXiv:1206.6885)
- Douglas J.N., Bash F.N., Bozayan F.A., Torrence G.W., Wolfe C., 1996, AJ 111, 1945
- Drake A.J., Djorgovski S.G., Mahabal A. et al. 2012, in “New Horizons in Time Domain Astronomy”, eds. E. Griffin et al., Cambridge Univ. Press, IAU Symp. 285, p. 306
- Eggen, D.J., Greenstein, J.L., 1967, ApJ 150, 929
- Fabbiano G., 2010, AIPC 1314, 1337
- Fleming, T.A., Gioia, I.M., Maccacaro, T., 1989, AJ 98, 692
- Flin P., Trevese D., Cirimele G., Hickson P., 1995, A&A Suppl. 110, 313
- Fuhrmann B., 1984, Mitt. Veränderl. Sterne 10, 97
- Gal R.R., de Carvalho R.R., Lopes P.A.A., Djorgovski S.G., Brunner R.J., Mahabal A., Odewahn S.C., 2003, AJ 125, 2064
- Garcia Lopez R.J., Randich S., Zapatero Osorio M.R., Pallavicini R., 2000, A&A 363, 958
- Gehrels N., Chincarini G., Giommi P., et al. 2004, ApJ 611, 1005
- Gershberg, R.E., Shakhovskaya, N.I., 1974, Izv. Krymsk. Astrofiz. Obs. 49, 73
- Gilfanov M., 2004, MN 349, 146
- Gioia, I.M., Luppino, G.A., 1994, ApJ Suppl. 94, 583
- Glozzio M., Brinkmann W., Laurent-Muehleisen S.A., Takalo L.O., Sillanpää A., 1999, A&A 352, 437
- Gomez P.L., Ledlow M.J., Burns J.O., Pinkney J., Hill J.M., 1997, AJ 114, 1711
- Green, R.F., Schmidt, M., Liebert, J., 1986, ApJ Suppl. 61, 305
- Greiner, J., Danner, R., Bade, N., Richter, G.A., Kroll, P., Komossa, S., 1996, A&A 310, 384
- Grimm H.-J. et al. 2003, MN 339, 793
- Grimm H.-J. et al. 2005, ApJS 161, 271
- Grindlay J., Tang S., Simcoe R., et al. 2009, ASP Conf. Proc. vol. 410, p 101
- Grindlay J., Doane A., Los E. et al. 2013, ATEL #5544
- Haberl F., 2005, Proc. 2005 EPIC XMM-Newton Consortium Meet., Ringberg Castle, Apr. 2005, Eds. U.G. Briel et al., MPE Report 288 (arXiv:0510480)
- Hadjijska E., Rabinowitz D., Baltay C. et al. 2012, in “New Horizons in Time Domain Astronomy”, eds. E. Griffin et al., Cambridge Univ. Press, IAU Symp. 285, p. 324
- Hagen H.-J., Groote D., Engels D., Reimers D., 1995, A&AS 111, 195
- Hall, O.S., Henry G.W., 1992, AJ 104, 1936
- Harrison T.E., Gehrz R.D., 1991, AJ 101, 587
- Häussler, K., 1975, Mitt. Sternw. Hartha 5, 14
- Hewett, P.C., Foltz C.B., Chaffee F.H., 1995, AJ 109, 1498
- Hewitt, A., Burbidge, C., 1987, ApJ Suppl. 63, 1
- Hipparcos & Tycho catalogues, 1997, ESA SP-1200
- Hoffmeister C., 1964, AN 288, 49
- Hoffmeister C., Richter G.A., Wenzel W., 1985, in *Variable Stars*, chap. 7, Springer: Berlin, p. 261ff
- Ivezic Z., Tyson J.A., Acosta E. et al. 2008, arXiv:0805.2366; living document on www.lsst.org/lsst/science/overview
- Ivanov G.A., 2002, Kinematika Fiz. Nebesn. Tel., 18, 287
- Jevers, H.M., Vasilewskis, S., 1978, AJ 83, 411
- Kholopov, P.N., 1982, (Ed. in chief), New Catalogue of Suspected Variable Stars, Moscow, Nauka
- Kholopov, P.N., 1985, (Ed. in chief), General Catalogue of Variable Stars (GCVS), vol. I-IV, Moscow, Nauka
- Khruzina T.S., Shugarov S.Yu., 1991, in “*Atlas of cataclysmic variables*”, Ed. A.M. Cherepashchuk, Moscow Univ. Press
- King L.J., Browne I.W.A., Marlow D.R., et al. 1999, MN 307, 225
- Kiraga M., 2012, Acta Astron. 62, 67
- Kiraga M., Stepień K., 2013, Acta Astron. 63, 53
- Komossa S., Greiner J., 1999, A&A 349, L45
- Kouzuma S., Yamaoka H., 2010, MN 405, 2062
- Krautter J., Zickgraf F.-J., Appenzeller I. et al. 1999, A&A 350, 743
- Kreysing, H. C., Diesch, C., Zweigle, J., et al. 1992, A&A 264, 623
- Kroll, P., Richter, G.A., 1996, Astron. Ges. Abstr. Ser. 12, 35
- Kroll, P., 2009, in Proc. of the Int. ICOMOS Symp., Hamburg, Oct. 2008, Ed. G. Wolfshmidt, p. 311
- Kulkarni S.R., 2012, arXiv:1202.2381
- Laurent-Muehleisen, S.A., Kollgaard R.I., Ryan P.J., Feigelson E.D., Brinkmann W., Siebert J., 1997, A&A Suppl. 122, 235
- Lonsdale C.J., Voges W., Bolter Th., Wolstencroft R.D., 2000, AAS 197, 9005
- Lopes P.A.A., de Carvalho R.R., Gal R.R., Djorgovski S.G., Odewahn S.C., Mahabal A.A., Brunner R.J., 2004, AJ 128, 1017
- Maccacaro T., Feigelson E.D., Fener M., et al. 1982, ApJ 253, 504
- MacLeod C.L., Ivezić Z., Sesar B. et al. 2012, ApJ 753, 106
- Massey P., Johnson K.E., DeGioia-Eastwood K., 1995, ApJ 454, 151
- Malaguti, G., Bassani, L., Caroli, E., 1994, ApJS 94, 517
- McCook, G.P., Sion, E.M., 1987, ApJ Suppl. 65, 603
- McMahon R.G., Irwin, M.J., Maddox, S.J., 2000, online catalogue I/267 in VizieR (<http://vizier.u-strasbg.fr/viz-bin/VizieR>)
- Meinunger, L., Wenzel, W., 1968, Veröff. Sternwarte Sonneberg 7, 385
- Mickaelian A.M., Hovhannisyan L.R., Engels D., Hagen H.-J., Voges W., 2006, A&A 449, 425
- Mineo S. et al. 2012, MN 419, 2095
- Motch C., Guillout P., Haberl F. et al. 1998, A&A Suppl. 132, 341
- Neckel, H., 1974, A&AS 18, 169
- Neuhäuser, R., Comeron, F., 1998, Sci 282, 83
- Noskova, R.I., Savelyeva, M.V., Archipova, V.P., Goranskij, V.P., Dokuchaeva, O.D., 1979, VS Prilochenije 3, 755
- Nousek, J.A., Takalo, L.O., Schmidt, D.G., et al. 1984, ApJ 277, 682
- Odenkirchen, M., Soubiran, C., Colin, J., 1998, New Ast. 3, 583
- Odewahn S.C., Aldering G. 1995, AJ 110, 2009
- Osczo A., Serra-Ricart M., Mediavilla E., Buitrago J., Goicoechea L.J., 1997, ApJ 491, L7
- Pietsch W., 2008, AN 329, 170
- Prager R., 1934, AN 232, 65
- Predehl P., Hasinger G., Böhringer H. et al. 2006, SPIE 6266, 62660P
- Pretorius M.L., Knigge C., Schwöpe A.D., 2013, MN 432, 570
- Randich, S., Schmitt, J.H.M.M., Prosser, C., 1996, A&A 313, 851
- Raveendran, A.V., 1984, ApSS 99, 171
- Richter, G., 1968, Veröff. Sternw. Sonneberg 7, 229
- Richter, G., 1975, AN 296, 65
- Richter, G.A., Bräuer, H.-J., Greiner, J., 1995, in IAU Coll. 151, *Flares and Flashes*, Lecture Notes in Physics, eds. Greiner, Duerbeck, Gershberg. Springer:Berlin, p. 69
- Richter, G.A., Greiner, J., 1995a, in IAU Coll. 151, *Flares and Flashes*, Lecture Notes in Physics 454, eds. Greiner, Duerbeck, Gershberg. Springer:Berlin, p. 106
- Richter, G.A., Greiner, J., 1995b, Abano-Padova Conf. on Catacl. Variables, eds. Bianchini, Della Valle, Orto, ASSL 205, 177
- Richter, G.A., Greiner, J., 1996, IBVS 4319
- Richter, G.A., Greiner, J., Kroll, P., 1998, IBVS 4622
- Richter, G.A., Greiner, J., 1999, in *Treasure Hunting in Astronomical Data Archives*, (eds.: P. Kroll, C. la Dous, H.-J. Bräuer), Harry Deutsch:Frankfurt (Main), p. 132
- Richter, G.A., Greiner, J., 1999, A&A 361, 1005
- Richter, G.A., Greiner, J., 2000, A&A 361, 1005
- Richter, G.A., Kroll, P., Greiner, J., Wenzel, W., Luthardt, R., Schwarz, R., 1997, A&A 325, 994
- Rines K., Geller M.J., Kurtz M.J., et al. 2001, ApJ 561, L41
- Rodono, M., 1980, Mem. Soc. Astron. Italiana 51, 623
- Saken J.M., Long K.S., Blair W.P., Winkler P.F., 1995, ApJ 443, 231
- Sandage, A., Veron, R., Wyndham, J.D., 1965, ApJ 142, 1307
- Schmitt J.H.M.M., Liefke C., 2004, A&A 417, 651
- Schlegel D., Abdalla F., Ahn C. et al. 2010, Proposal to NOAO, http://bigboss.lbl.gov/docs/BigBOSS_NOAO_public.pdf
- Schneider D.P., Gunn J.E., Hoessel J.G., 1983, ApJ 264, 337
- Shara M.M., Potter M.M., Moffat A.F.J., 1990, AJ 100, 540
- Skiff B.A., 1997, IBVS 4459
- Sowards-Emmerd D., Romani R.W., Michelson P.F., Healey S.E., Nolan P.L., 2005, ApJ 626, 95
- Stoeckle, J.T., Liebert, J., Gioia, I.M., et al. 1983, ApJ 273, 458
- Stoeckle J.T., Morris S.L., Gioia I.M. et al. 1991, ApJS 76, 813
- Takalo L.O., Nousek J.A., 1985, PASP 97, 570
- Taylor A.R., Goss W.M., Colemann P.H., van Leeuwen J., Wallace B.J., 1996, ApJ Suppl. 107, 239
- Tonry J.L., Stubbs C.W., Lykke K.R. et al. 2012, ApJ 750, 99
- Townsley L.K., et al 2006, AJ 131, 2140
- Trümper, J., 1983, Adv. Space Res. 2, p. 241
- Turner M.L., Rutledge R.E., Letcavage R. et al. 2010, ApJ 714, 1424
- Ueda Y., Ishisaki Y., Takahashi T., Makishima K., Ohashi T., 2001, ApJ Suppl. 133, 1
- Uomoto, A.K., Wills, B.J., Wills, D., 1976, AJ 81, 905
- Vanden Berk D.E., Willite B.C., Kron R.G. et al. 2004, ApJ 601, 692
- Voges W., et al. 1999, A&A 349, 389
- Vogt, N., Bateson, F.M., 1982, A&A Suppl. 48, 383

Vogt, N., Kroll, P., Splittgerber E., 2004, A&A 428, 925
Wenzel, W., Richter, G.A. Luthardt, R., Schwarz, R., 1995, IBVS 4182
White R.L., Becker R.H., Gregg M.D., et al. 2000, ApJS 126, 133
Wills D., Wills B.J., 1976, ApJ Suppl. 31, 143
Wu H., Xue S.-J., Xia X.Y., Deng Z.G., Mao S., 2002, ApJ 576, 738
Zeilik, M., Batuski, R., Burke, S., Eisten, R., Smith, P., 1982, IBVS 2257
Zhang H., Xue S., Burstein D. et al. 2004, AJ 127, 2579
Zickgraf F.-J., Thiering I., Krautter J. et al. 1997, A&A Suppl. 123, 103
Zickgraf F.-J., Engels D., Hagen H.-J., Reimers D., Voges W., 2003, A&A 406, 535
Zimmermann H.U., Becker W., Belloni T., Döbereiner S., Izzo C., Kahabka P., Schwentker O., 1994, MPE report 257
Zuo W., Wu X.-B., Liu Y.-Q., Jiao C.-L., 2012, ApJ 758, 104




Fig. 40a. DSS blue finding chart of Com sources Com001 to Com020. The size is 2.5×2.5 , North is up and East to the left. The central circle is the $30''$ error circle as derived from our source detection on the RASS data. Shifted circles of similar (or slightly smaller) size are ROSAT source position errors taken from the 1RXS catalogue (see 4th column in Tables 1a, 1b), and very small shifted circles are source positions taken from the ROSAT HRI (1RXH) catalogue or XRT positions from our Swift follow-up (see the notes on individual objects for details). Many of the alphabetically labelled objects have been investigated for variability, and details are given in Table 5a,5b.




Fig. 40a. contd. Com sources Com021 to Com044.




Fig. 40a. contd. Com sources Com045 to Com068. Object *a* for Com064 = RX J1252.9+2343 is either strongly variable or very red, and not visible on the DSS plate.




Fig. 40a. contd. Com sources Com069 to Com092.




Fig. 40a. contd. Com sources Com093 to Com116.




Fig. 40a. contd. Com sources Com117 to Com140.




Fig. 40a. contd. Com sources Com141 to Com164.




Fig. 40a. contd. Com sources Com165 to Com188.




Fig. 40a. contd. Com sources Com189 to Com212.




Fig. 40a. contd. Com sources Com213 to Com236.




Fig. 40a. contd. Com sources Com237 to Com238.




Fig. 40b. DSS blue finding charts of Sge sources Sge001 to Sge024. The size is 2.5×2.5 (except for Sge019 = RX J1959.6+2243 which is three times as large to show the full extent of the underlying nebula), North is up and East to the left.




Fig. 40b. contd. Sge sources Sge025 to Sge048.




Fig. 40b. contd. Sge sources Sge049 to Sge072.




Fig. 40b. contd. Sge sources Sge073 to Sge103.




Fig. 40b. contd. Sge sources Sge104 to Sge127.




Fig. 40b. contd. Sge source Sge132.




Fig. 41a. Optical light curves of Com sources. For each source, the measurements are shown by crosses, and upper limits as arrows. Filled triangles (red) and hexagons (green) are measurements of the POSS and the Tautenburg Schmidt plates, respectively, and open stars refer to CCD observations at SAO Russia.




Fig. 41a. contd. Optical light curves of Com sources.




Fig. 41a. contd. Optical light curves of Com sources.




Fig. 41a. contd. Optical light curves of Com sources.




Fig. 41a. contd. Optical light curves of Com sources.





Fig. 41a. contd. Optical light curves of Com sources.



88

Fig. 41b. Optical light curves of Sge sources. Labels as in 41a.




Fig. 41b. contd. Optical light curves of Sge sources.




Fig. 41b. contd. Optical light curves of Sge sources.




Fig. 41b. contd. Optical light curves of Sge sources.




Fig. 41b. contd. Optical light curves of Sge sources.